

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/11, 15/00, 15/63, C07H 21/02, 21/04		A1	(11) International Publication Number: WO 00/04140 (43) International Publication Date: 27 January 2000 (27.01.00)
(21) International Application Number: PCT/US99/15849 (22) International Filing Date: 14 July 1999 (14.07.99) (30) Priority Data: 60/092,921 15 July 1998 (15.07.98) US 60/092,922 15 July 1998 (15.07.98) US 60/092,956 15 July 1998 (15.07.98) US (71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Olney, MD 20832 (US). KOMATSOUKIS, George [US/US]; 9518 Garwood Street, Silver Spring, MD 20901 (US). DUAN, Roxanne, D. [US/US]; 5515 Northfield Road, Bethesda, MD 20817 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonville, MD 20882 (US). MOORE, Paul, A. [US/US]; 19005 Leatherbark Drive, Germantown, MD 20874 (US). SHI, Yang-gu [CN/US]; Apartment 102, 437 West Side Drive, Gaithersburg, MD 20878 (US). LAFLEUR, David, W. [US/US]; 3142 Quesada Street, N.W., Washington, DC 20015 (US). EBNER, Reinhard [DE/US]; 9906 Shelburne Terrace #316, Gaithersburg, MD 20878 (US). OLSEN, Henrik, S. [DK/US]; 182 Kendrick Place #24, Gaithersburg, MD 20878 (US). BREWER, Laurie, A. [US/US]; Apartment 115, 410 Van Dyke Street, St. Paul, MN 55119-4321 (US). FLORENCE, Kimberly, A. [US/US]; 12805 Atlantic Avenue, Rockville, MD 20851 (US). YOUNG, Paul, E. [US/US]; 122 Beckwith Street, Gaithersburg, MD 20878 (US). MUCENSKI, Michael [US/US]; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US). ENDRESS, Gregory, A. [US/US]; 9729 Claggett Farm Drive, Potomac, MD 20854 (US). SOPPET, Daniel, R. [US/US]; 15050 Stillfield Place, Centreville, VA 22020 (US). (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).			
Published <i>With international search report.</i>			
(54) Title: 71 HUMAN SECRETED PROTEINS			
(57) Abstract <p>The present invention relates to novel human secreted proteins and isolated nucleic acids containing the coding regions of the genes encoding such proteins. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human secreted proteins. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating disorders related to these novel human secreted proteins.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

71 Human Secreted Proteins

Field of the Invention

This invention relates to newly identified polynucleotides and the polypeptides encoded by these polynucleotides, uses of such polynucleotides and polypeptides, and their production.

Background of the Invention

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic reticulum (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in secretory vesicles (or secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins include the commercially valuable human insulin, interferon, Factor VIII, human

growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent
5 medical disorders by using secreted proteins or the genes that encode them.

Summary of the Invention

The present invention relates to novel polynucleotides and the encoded polypeptides. Moreover, the present invention relates to vectors, host cells,
10 antibodies, and recombinant methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting disorders related to the polypeptides, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying binding partners of the polypeptides.

15

Detailed Description

Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

20 In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of
25 matter, or particular cell is not the original environment of the polynucleotide.

In the present invention, a "secreted" protein refers to those proteins capable of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released
30 into the extracellular space, the secreted protein can undergo extracellular processing

to produce a "mature" protein. Release into the extracellular space can occur by many mechanisms, including exocytosis and proteolytic cleavage.

In specific embodiments, the polynucleotides of the invention are less than 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, or 7.5 kb in length. In a further embodiment, polynucleotides of the invention comprise at least 15 contiguous nucleotides of the coding sequence, but do not comprise all or a portion of any intron. In another embodiment, the nucleic acid comprising the coding sequence does not contain coding sequences of a genomic flanking gene (i.e., 5' or 3' to the gene in the genome).

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the ATCC Deposit Number. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an

overnight incubation at 42° C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

- 5 Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or
- 10 temperature. For example, lower stringency conditions include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent
- 15 hybridization can be done at higher salt concentrations (e.g. 5X SSC).

- Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and
- 20 commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

- Of course, a polynucleotide which hybridizes only to polyA+ sequences (such as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a
- 25 complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

- The polynucleotide of the present invention can be composed of any
- 30 polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of

single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

The polypeptide of the present invention can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation,

gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

5 (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

10 "SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

Polynucleotides and Polypeptides of the Invention

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 1

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

30 PFCSGFFPSLWIYLPFIFNVSDLWMGSLSGCALPFCLXVFFLTVSPSAVGLLXF
AGGPLQTLFAWVSPVEAAEQRLLPVLSSGSFVSEGTCQMPARALLYEVSVG

PYWEIPPSQDTRRSGLTYLRRQSDP (SEQ ID NO: 195) . Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in pancreas islet cell tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as
5 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the pancreas, including cancer and diabetes. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For
10 a number of disorders of the above tissues or cells, particularly of the pancreas, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., endocrine, cancerous, or wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the
15 standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in tumors of pancreatic islet cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis, treatment and intervention of such tumors, in addition to other endocrine or
20 gastrointestinal tumors where expression has been indicated. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or
25 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:11 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
30 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or

more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1099 of SEQ ID NO:11, b is an integer of 15 to 1113, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:11, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 2

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the

10 following amino acid sequence:

HEGSCRAPGFSAHKGRGCPSPRMTLPSRALASLGVGWGMRLRNQVTVSCG
GSRWSSRVALGAFSWVCGVALVLQPSGGGLTSPSEGCWEGELALAVLRA
PGGSPS (SEQ ID NO: 196). Polynucleotides encoding these polypeptides are also provided.

15 This gene is expressed equally in in .

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic disorders, particularly leukemia. Similarly,

20 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the vascular and immune systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hemolymphoid,

25 cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 104 as residues: Gly-29 to Ser-35, Ser-63 to Cys-68. Polynucleotides encoding said polypeptides are also provided.

5 The tissue distribution in hemangiopericytoma, breast lymph node, and bone marrow indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in
10 Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency
15 etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions,
20 in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
25 related to SEQ ID NO:12 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general
30 formula of a-b, where a is any integer between 1 to 969 of SEQ ID NO:12, b is an

integer of 15 to 983, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:12, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 3

5 The translation product of this gene shares sequence homology with the *Drosophila melanogaster* slit protein, a secreted protein that contains both an EGF domain and Leucine Rich Repeat domains. It is thought to be important in the development of midline glia and commissural axon pathways (See e.g., Rothberg et al. Genes Dev. 4:2169-87 (1990); which is hereby incorporated by reference herein).

10 This gene is expressed primarily in human hippocampus.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological, and developmental disorders. Similarly, polypeptides and
15 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neurological system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neurological, cancerous, or wounded tissues) or
20 bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 The tissue distribution within human hippocampus combined with the
25 homology to the *Drosophila* slit protein, indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and
30 elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease,

Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:13 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 959 of SEQ ID NO:13, b is an integer of 15 to 973, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:13, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 4

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by

the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

IPLTLPGIFLLIRLFWRLGQSICGPGKLVLPQFCCGCAVISGHCVPRGMPSSW
LPGCFVLLCLVAVGCQLREWGVGVS AVGLLALPHLQVLGMRGRGLISGG

5 (SEQ ID NO: 197) . Polynucleotides encoding these polypeptides are also provided.

The gene encoding the disclosed cDNA is believed to reside on chromosome 16. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 16.

This gene is expressed in KMH2 cells, osteoblasts, fetal spleen, Jurkat
10 membrane bound polysomes, breast, and cerebellum.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cancer, immune, and skeletal disorders. Similarly, polypeptides and
15 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, skeletal, cancerous, or wounded tissues) or bodily
20 fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in KMH2 cells, osteoblasts, and fetal spleen indicates
25 that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Expression of this gene product in fetal spleen and T-cells indicates a role in the regulation of the
30 proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved

in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene
5 product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory
10 bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity
15 disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, scleroderma and tissues. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that
20 recruits hematopoietic cells to sites of injury. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify
25 agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
available and accessible through sequence databases. Some of these sequences are
25 related to SEQ ID NO:14 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
30 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1444 of SEQ ID NO:14, b is an

integer of 15 to 1458, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:14, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 5

- 5 The translation product of this gene shares sequence homology with phospholipase A2 which cleaves fatty acids from carbon 2 of glycerol (ref. Prosite pattern documentation for PS2_HIS). Many snake venoms contain phospholipase A2, which prevents transmission of nerve impulses to muscles by blocking the release of acetylcholine from the neuron. Therefore, included in this invention as preferred
- 10 domains are Phospholipase A2 histidine active site domains, which were identified using the ProSite analysis tool (Swiss Institute of Bioinformatics). Phospholipase A2 is an enzyme which releases fatty acids from the second carbon group of glycerol. Structurally, PA2's are small and rigid proteins of 120 amino-acid residues that have four to seven disulfide bonds. PA2 binds a calcium ion which is required for activity.
- 15 The side chains of two conserved residues, a histidine and an aspartic acid, participate in a 'catalytic network'. Two different signature patterns for PA2's were developed. The first is centered on the active site histidine and contains three cysteines involved in disulfide bonds. The consensus pattern is as follows: C-C-x(2)-H-x(2)-C [H is the active site residue].
- 20 Preferred polypeptides of the invention comprise a Phospholipase A2 histidine active site domain selected from the following amino acid sequences: CCNQHDRC (SEQ ID NO: 199), SLTKCCNQHDRCYET (SEQ ID NO: 200) , and/or LTKCCNQHDRCYETCG (SEQ ID NO: 201) . Polynucleotides encoding these polypeptides are also provided. Further preferred are polypeptides comprising the
- 25 Phospholipase A2 histidine active site domain of the sequence listed in Table 1 for this gene, and at least 5, 10, 15, 20, 25, 30, 50, or 75 additional contiguous amino acid residues of this referenced sequence. The additional contiguous amino acid residues is N-terminal or C- terminal to the Phospholipase A2 histidine active site domain. Alternatively, the additional contiguous amino acid residues is both N-terminal and
- 30 C-terminal to the Phospholipase A2 histidine active site domain, wherein the total N- and C-terminal contiguous amino acid residues equal the specified number. The

above preferred polypeptide domain is characteristic of a signature specific to Phospholipase A2 proteins. Based on the sequence similarity, the translation product of this gene is expected to share at least some biological activities with Phospholipase A2 proteins. Such activities are known in the art, some of which are described
5 elsewhere herein, or see, for example, McIntosh, et al. J. Biol. Chem. 270 (8), 3518-3526 (1995), incorporated herein by reference.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the
10 following amino acid sequence:

GPAGKEAWIWSWLLPSPGPAPLPSASWGLCGDAPR
AAARGPVEPGAARMALLSRPALTLALLMAAVVRCQEQAQTTDWRATLKTI
RNGVHKIDTYLNAALDLLGGEDGLCQYKCSGSKPFPRYGYKPSPPNGCGSP
LFGXHLNIGIPSLTKCCNQHDRCYETCGKSKNDCDEEFQYCLSKICRDVQKTL
15 GLTQHVQACETTVELLFDSEVIHLGCKPYLDSQRAACRCHYEETDL (SEQ ID
NO: 198) . Polynucleotides encoding these polypeptides are also provided.

The gene encoding the disclosed cDNA is believed to reside on chromosome 4. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 4.

20 This gene is expressed in a variety of cell types with no single type predominating.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
25 not limited to, neurological disorders, or metabolism disorders, specifically phospholipase A2 deficiencies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neuromuscular system, expression of this gene at
30 significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., pancreas, cancerous and wounded tissues) or bodily fluids (e.g., bile, lymph,

serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 107 as residues: Gln-23 to Asp-30, Lys-66 to Cys-87. Polynucleotides encoding said polypeptides are also provided.

The ubiquitous tissue distribution and homology to phospholipase A2 indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neuromuscular disorders. Alternatively, considering the activity of
10 phospholipase A2 as a block for neuro- transmission may suggest that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette
15 Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked
20 disorders, or disorders of the cardiovascular system. Alternatively, the homology to Phospholipase A2 proteins may indicate a potential use for the protein product of this gene in diagnosis, treatment and/or prevention of metabolism disorders, specifically deficiencies in Phospholipase A2. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate
25 ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
30 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:15 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general
 5 formula of a-b, where a is any integer between 1 to 1991 of SEQ ID NO:15, b is an integer of 15 to 2005, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:15, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 6

10 In a specific embodiment polypeptides of the invention comprise the following amino acid sequence:

GTSSARPRGALPGGSAPSAPHGQLPGRAQPAPVSGPPPTSGLCHFDPAAPWPL
 WPGFWQLPPHPQDWPAPHDIPQDWVSFLRSFGQLTLCPRNGTVTGKWRGSH
 VVGLLTTLNFGDGPDRNKTRTFQATVLGSQMGLKGSSAGQLVLITARVTTER
 15 TAGTCLYFSAVPGILPSSQPPISCSEEGAGNATLSPRMGEECVSVWSHEGLVLT
 KLLTSEELALCGSRLLVLGSFLLFCGLLCCVTAMCFHPRRESHWSRTRL
 (SEQ ID NO: 202). Polynucleotides encoding these polypeptides are also provided.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by
 20 the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

ARAPPGPEGLSPEAQPLLPMGNCQAGHNLHLCLAHHPPLVCATLILLGLS
 GLGLGSFLLTHRTGLRT LTSPRTGSLF (SEQ ID NO: 203). Polynucleotides
 encoding these polypeptides are also provided.

25 This gene is expressed in a wide variety of tissue types including testes, cerebellum, dendritic cells, breast cancer, umbilical vein endothelial cells, epididymus, corpus colosum, chronic synovitis, liver hepatome, normal breast, osteoblasts, melanocytes, B cell lymphomas, and to a lesser extent in other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as
 30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, cancer, particularly of endothelial tissues. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system,

5 expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., endothelial, cancerous, or wounded tissues) or bodily fluids (e.g., lymph, seminal fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in

10 healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 108 as residues: Thr-52 to Gly-57. Polynucleotides encoding said polypeptides are also provided.

Expression within embryonic tissue and other cellular sources marked by

15 proliferating cells indicates that the protein product of this gene may play a role in the regulation of cellular division and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Additionally, the expression

20 in hematopoietic cells and tissues indicates that this protein may play a role in the proliferation, differentiation, and/or survival of hematopoietic cell lineages. In such an event, this gene is useful in the treatment of lymphoproliferative disorders, and in the maintenance and differentiation of various hematopoietic lineages from early hematopoietic stem and committed progenitor cells. Similarly, embryonic

25 development also relies on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain

30 neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have

applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:16 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 929 of SEQ ID NO:16, b is an integer of 15 to 943, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:16, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 7

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

RFLSVXPQXEVPFLLHPCVCFXGGHPSLLPDPCRAVGGGWEAPRCCLHEALC
 QSLGCKAEEIVSVSESSSAQRCWYLLRGRKAGGRGPASPVLFALMRLESLCH
 LCLACLFFRLPATRTVYCMNEAEIVDVALGILIESRKQXKACEQPALAGADNP
 EHSPPCSVSPHTSSGSSSEEDSGKQALXPGLSPSQRPGGSSSACSRSPEEEE

- 5 EEDVLKYVREIFFS (SEQ ID NO: 204) . Polynucleotides encoding these polypeptides are also provided. Polynucleotides of the invention do not consist of the nucleic acid sequences shown as GeneSeq Accession Nos: V59595 and V59744, which are hereby incorporated herein by reference.

This gene is expressed primarily in a variety of immune cell types, including
 10 stromal cells, dendritic cells, leukocytes, activated T-cells, macrophages, monocytes, neutrophils and to a lesser extent in a variety of other adult and fetal tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
 15 not limited to, cancer and other proliferative disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain
 20 tissues or cell types (e.g., immune, cancerous, or wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

25 The tissue distribution in immune cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Expression of this gene product in fetal
 30 tissue and various hematopoietic cancers indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all

hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

5 Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatou's Disease, inflammatory
10 such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, scleroderma and tissues. Moreover, the protein may represent a secreted
15 factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity,
20 raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

 Many polynucleotide sequences, such as EST sequences, are publicly
25 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:17 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
30 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1489 of SEQ ID NO:17, b is an

integer of 15 to 1503, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:17, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 8

5 When tested against Jurkat T-cell lines, supernatants removed from cells containing this gene activated the NF-kB (Nuclear Factor kB) pathway. Thus, it is likely that this gene activates T-cells through the NF-kB signal transduction pathway. NF-kB is a transcription factor activated by a wide variety of agents, leading to cell activation, differentiation, or apoptosis. Reporter constructs utilizing the NF-kB
10 promoter element are used to screen supernatants for such activity. Preferred polypeptides of the invention comprise the following amino acid sequence: VPGWPRACSPCQADSPRAHPPKLRGILRWAPVPLXCAALCPPLDSG MSMAACPEAPEPSFLREVPS SPASTQWHRPCNFRQVEANPRKEPKNLVWRD VSLGQXSRTPRGSGLELVRVCGGGMQRD KTVVEERVGEERERERERESLGG
15 AGKHGEMRCVYVRESVGAPGRAGGGGNGVNSVGCVRTVHSGSXPPPSAGV S (SEQ ID NO: 205). Polynucleotides encoding these polypeptides are also provided.

 This gene is expressed primarily in parts of the brain such as cerebellum and frontal lobe.

 Therefore, polynucleotides and polypeptides of the invention are useful as
20 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of
25 the above tissues or cells, particularly of the central nervous system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, cancerous, or wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene
30 expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in cerebellum and frontal lobe indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, prevention and/or treatment of neurodegenerative disease states and behavioural disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:18 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1498 of SEQ ID NO:18, b is an

integer of 15 to 1512, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:18, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 9

5 In a specific embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

TRPGKELNLVFGLQLSMARIGSTVNMNLMGWLYSKIEALLGSAGHTTLGITL
 10 MIGGITCILSLICALALAYLDQRAERILHKEQGKTGEVIKLTVDKDFSLPLWLIF
 IICVCYYVAVFPFIGLGKVFTEKFGFSSQAASAINSVVYVISAPMSPVFGLLV
 DKTGKNIIWVLCA (SEQ ID NO: 206) . Polynucleotides encoding these
 polypeptides are also provided.

The gene encoding the disclosed cDNA is believed to reside on chromosome
 15 3. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 3.

This gene is expressed primarily in fetal tissue, and to a lesser extent in a variety of adult human tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as
 20 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, fetal abnormalities, particularly developmental disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For
 25 a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., developing, or cancerous and wounded tissues) or bodily fluids (e.g., amniotic fluid, lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having
 30 such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 111 as residues: Lys-30 to Thr-35. Polynucleotides encoding said polypeptides are also provided.

- 5 The tissue distribution in fetal tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of cancer and other proliferative disorders. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.
- 10 Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Expression within embryonic tissue and other cellular sources marked by proliferating cells
- 15 indicates that this protein may play a role in the regulation of cellular division. Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating,
- 20 detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in
- 25 proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Additionally, the expression in hematopoietic cells and tissues indicates that this protein may play a role in the proliferation, differentiation, and/or survival of hematopoietic cell lineages. In such an event, this gene is useful in the treatment of lymphoproliferative disorders,
- 30 and in the maintenance and differentiation of various hematopoietic lineages from early hematopoietic stem and committed progenitor cells. Furthermore, the protein

may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:19 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1641 of SEQ ID NO:19, b is an integer of 15 to 1655, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:19, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 10

The translation product of this gene shares sequence homology with human histiocyte-secreted factor (HSF) which is a novel cytokine that shows in vivo antitumour activity without the cytotoxicity associated with tumour necrosis factor. Furthermore, The translation product of this gene also shares sequence homology with the human endogenous virus S71 gag polypotein, the sequence of which is believed to represent a transformation locus for several cancers (See Genebank Accession No. pir|A46312|A46312; all references available through this accession are hereby incorporated by reference herein). Similarly, The translation product of this gene also shares homology with B219, a sequence that is expressed in at least four isoforms in very primitive hematopoietic cell populations which may represent a novel hemopoietin receptor (See, e.g., Cioffi, et al. Nat. Med. 2:585-589 (1996), which is hereby incorporated by reference herein). In a preferred embodiment polypeptides of the invention comprise the following amino acid sequence:

CKDLCSR VYLLT LSP LLSYDPATSHSPRNTQ (SEQ ID NO: 207) . Also preferred are the polynucleotides encoding these polypeptides.

This gene is expressed primarily in tonsil, and colon, and to a lesser extent in a wide variety of human tissues.

5 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, hematopoietic, and gastrointestinal disorders, particularly tumors of the colon and tonsil. Similarly, polypeptides and antibodies directed to
10 these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hematopoietic, digestive and immune systems, expression of this gene at significantly higher or lower levels is routinely detected in
15 cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

20 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 112 as residues: Met-1 to Cys-6. Polynucleotides encoding said polypeptides are also provided.

 The tissue distribution in tonsil and colon, combined with the homology to human histiocyte growth factor, the human endogenous viral protein, and B219
25 strongly indicate that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis, treatment and/or prevention, of a variety of hematopoietic and immune system disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19,
30 20, and 27, and elsewhere herein. Expression of this gene product in tonsils indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation

of potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g., by boosting immune responses).

5 Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, 10 such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, scleroderma and tissues. Moreover, the protein may represent a secreted 15 factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, 20 raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

 Many polynucleotide sequences, such as EST sequences, are publicly 25 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:20 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or 30 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2511 of SEQ ID NO:20, b is an

integer of 15 to 2525, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:20, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 11

5 The gene encoding the disclosed cDNA is believed to reside on chromosome 7. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 7.

 In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

10 IICECWEEECQSCRLKITQPREICRMDFLVLFLFYLASVLMGLVLICVCSKTHS
 LKGLARGGAQIFSCIPECLQRAXHGLLHYLFHTRNHTFIVLHLVLQGMVYTE
 YTWEVFGYCQELELSLHYLLLPYLLLGVNLFFFTLTCGTNPGIITKANELLFLH
15 VYEFDEVMPKPNVRCSTCDLRKPARSKHCSVCNWCVHRFDHHCVWVNCCI
 GAWNIRYFLIYVLTLTASAATVAIVSTTFLVHLVVMSDLYQETYIDDLGHLHV
 MDTVFLIQYLFTFPRIVFMLGFVVLSFLLGGYLLFVLYLAATNQTNEWYR
 GDWAWCQRCPLVAWPPSAEPQVHRNIHSHGLRSNLQEIFLPAFPCHERKKQE
 (SEQ ID NO: 208) . Polynucleotides encoding these polypeptides are also provided.

20 This gene is expressed primarily in colon and brain and to some extent in all tissues.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological and digestive disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system and digestive system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neurological, gastrointestinal, or cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine,

synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly,
10 the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive
15 compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation,
20 neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Alternatively, expression of this gene in colon may indicate a role in the detection, prevention and/or treatment of colon disorders such as colon cancer, Crohn's Disease, ulcers, and digestive tract disorders in general. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to
25 isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
30 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:21 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general
5 formula of a-b, where a is any integer between 1 to 1382 of SEQ ID NO:21, b is an integer of 15 to 1396, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:21, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 12

10 When tested against Reh cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activation site) pathway. Thus, it is likely that this gene activates B-cells through the Jaks-STAT signal transduction pathway. GAS is a promoter element found upstream in many genes which are involved in the Jaks-STAT pathway. The Jaks-STAT pathway is a large, signal
15 transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. This gene maps to chromosome 7, and therefore, is used as a marker in linkage analysis for chromosome 7.

20 This gene is expressed primarily in brain, and in the developing embryo.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological, behavioral, immune, and developmental disorders.
25 Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous and developmental systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural,
30 developing, immune, or cancerous and wounded tissues) or bodily fluids (e.g., lymph, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue

or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic
5 epitopes shown in SEQ ID NO: 114 as residues: Lys-60 to Asn-67. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions.
10 Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia,
15 trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, the tissue distribution in developing
20 embryo indicates that the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Alternatively, the biological activity within B-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of
25 immune system disorders. Activation of genes within B-cells indicates a role for this protein in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of
30 cancer (e.g., by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions; in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:22 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1055 of SEQ ID NO:22, b is an integer of 15 to 1069, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:22, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 13

The gene encoding the disclosed cDNA is believed to reside on chromosome 6. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 6.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

LLSFKIRGLRTEDAGWAQSSSGGLCVRGDAFWMPSSSSGLGSPSRPPSSFLCL
LLLLLPPAALALLFFLDFFPPRAAVSPFLPDHCSARQPRVWRRETLNRSASGL

GCWARSTEQGA VGVATGTVLDI SLPASCLSLWPPGPSGGI (SEQ ID NO: 209)

. Polynucleotides encoding these polypeptides are also provided.

In a specific embodiment polypeptides of the invention comprise the following amino acid sequence:

5 QLGLCLTSASLPPASRCGHQAPLGASDLSAHSAPGFSDSYFTMSCQSSLSRA
EILQCPLVPSVSPPTHLPQGRANKSSRASLPLLPQTHWCLFPSARGWRRGIQSG
LPPGGSCTSPRSPPQTLHQHITLVNHNTSYWQSPST (SEQ ID NO: 210) ,
HQPCLLPLAVATRPLWGHLTCLPIHLVSVTLTSPCLANQAFQGQRSYNAL
WCPLFLLPTSPKGEQTNHPEACPCFPKLTGVFSLQHVVGAEEFSQVFLDVD
10 PVPVLDHLLKLFTSTSHLLIIPHIGKAPAPDSLL EELSLSLATHCKVAVARFT
(SEQ ID NO: 211) . Also preferred are the polynucleotides encoding these
polypeptides. Polynucleotides of the invention do not consist of the nucleic acid
sequence shown as GeneSeq Accession No. X04377, which is hereby incorporated
herein by reference.

15 This gene is expressed primarily in brain.

Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, behavioral and neurological disorders. Similarly, polypeptides and
20 antibodies directed to these polypeptides are useful in providing immunological
probes for differential identification of the tissue(s) or cell type(s). For a number of
disorders of the above tissues or cells, particularly of the central nervous system,
expression of this gene at significantly higher or lower levels is routinely detected in
certain tissues or cell types (e.g., neural, or cancerous and wounded tissues) or bodily
25 fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another
tissue or cell sample taken from an individual having such a disorder, relative to the
standard gene expression level, i.e., the expression level in healthy tissue or bodily
fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic
30 epitopes shown in SEQ ID NO: 115 as residues: Pro-2 to Gly-7, Ser-10 to Ser-16,

Pro-52 to Val-62, Arg-64 to Ser-73. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:23 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1644 of SEQ ID NO:23, b is an

integer of 15 to 1658, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:23, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 14

5 The translation product of this gene was shown to have homology to the lysosomal mannosidase alpha-B protein (See Genebank Accession No. P34098) which is thought to be important in protein metabolism. One embodiment of this gene comprises polypeptides of the following amino acid sequence:

MAAEGSRFSSQSPGLVDRQGPKCDPSRLVSPWGRHGLRILQIGHHHGRDGGQH
10 EATHHLLRVLRAPRVGKADEGAVDSDPSTPLQLKHEAAHAEDHAQQVHVVR
RRVVQGRVTFARRGLVPQHFVRPPWVRHIVSGHSESKARSRLFRCRNRSFRR
AS (SEQ ID NO: 212) , and/or

RLVSPWGRHGLRILQIGHHHGRDGGQHEATHHLL RVLRA (SEQ ID NO: 213) .

15 An additional embodiment is the polynucleotides encoding these polypeptides. This gene maps to chromosome 19, and therefore, is used as a marker in linkage analysis for chromosome 19.

 This gene is expressed primarily in brain, placenta, fetal liver, and to a lesser extent in most tissues.

 Therefore, polynucleotides and polypeptides of the invention are useful as
20 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological, reproductive, and hepatic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For
25 a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, hepatic, or cancerous and wounded tissues) or bodily fluids (e.g., bile, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a
30 disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 116 as residues: Asn-34 to Lys-42, Leu-60 to Trp-70. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution predominantly in brain indicates a role in the
5 detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder. Alternatively, the tissue distribution in liver indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of
10 liver disorders and cancers (e.g., hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). In addition the expression in fetus would suggest a useful role for the protein product in developmental abnormalities, fetal deficiencies, pre-natal disorders and various wound-healing models and/or tissue trauma.

15 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:24 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is
20 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1063 of SEQ ID NO:24, b is an integer of 15 to 1077, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:24, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 15

This gene is expressed primarily in spinal cord, Merkel cells, and adipose tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as
30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, disorders of the nervous and immune systems, particularly those disorders relating to the CNS involving lipid metabolism disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For
5 a number of disorders of the above tissues or cells, particularly of the nervous and immune systems and adipose tissue, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, immune, or cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an
10 individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in spinal cord, Merkel cells and adipose tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the
15 treatment and/or diagnosis of diseases the nervous systems, such as spinal cord injury, neurodegenerative diseases, muscular dystrophy or obesity. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
20 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:25 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
25 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1191 of SEQ ID NO:25, b is an integer of 15 to 1205, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:25, and where b is greater than or equal to a + 14.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 16

The translation product of this gene shares sequence homology with the human uncoupling protein-2 which is thought to be important in energy metabolism, obesity, and the predisposition of hyperinsulinemia (See Genebank Accession No. gi|1857278). Recently, another group published on this gene, designating it brain mitochondrial carrier protein-1 (BCMP1) (J Biol Chem 1998 Dec 18;273(51):34611-5). One embodiment of this gene comprises polypeptides of the following amino acid sequence: PTDVLKIRMQAQ (SEQ ID NO: 214) , and/or TYEQLKR (SEQ ID NO: 215) . An additional embodiment is the polynucleotides encoding these polypeptides. This gene maps to the X chromosome, and therefore, is used as a marker in linkage analysis for the X chromosome.

This gene is expressed primarily in manic depression brain tissue, epileptic frontal cortex, human erythroleukemia cell line, T-helper cells, and to a lesser extent in endothelial and amygdala cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the central nervous system or hematopoietic/immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system or hematopoietic/immune systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, hemolymphoid, or cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 118 as residues: Ser-34 to Thr-39, Gln-198 to Leu-205. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in neural tissues combined with the homology to the human uncoupling protein indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and/or treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Alternatively, given the homology to uncoupling proteins, the gene and/or its translation product may also be used in the diagnosis, treatment, and/or prevention of thermogenesis disorders such as obesity, cachexia, and hyperinsulinemia. Uncoupling proteins dissipate the proton gradient created from the oxidation of fuels by the electron transport chain, thus releasing stored energy as heat. Dysfunction of thermogenesis can induce disorders such as obesity and cachexia. It is thought that obesity may result from decreased thermogenesis in humans. Alternatively, cachexia is a metabolic state in which energy expenditure exceeds food intake, for example in anorexia nervosa. Uncoupling proteins is useful for the treatment and/or prevention of diseases and/or disorders involved with aberrant metabolic and thermogenic pathways. The following method provides for the determination of respiration uncoupling activity of the polypeptides of the present invention, including fragments and variants of the full length proteins.

Briefly, yeast are transfected with an expression vector expressing polypeptide of the present invention as previously described by Bouillaud et al., EMBO J., 13:1990 (1994) (incorporated by reference herein in its entirety). Rates of growth in liquid medium of transformed yeast are measured in the presence of galactose, which induces expression, as described in International Publication No. WO 98/31396

(incorporated by reference herein in its entirety). Instantaneous generation times are compared between the polypeptide of the present invention and appropriate controls. An in vivo decrease of membrane potential associated with uncoupling of respiration is analyzed by flow cytometry of yeast labeled with the potential sensitive probe

5 DiOC6 (3) (3,3'-dihexyloxacarbocyanine iodine, Molecular Probes, Eugene, OR). The ability of a polypeptide of the present invention to influence mitochondrial activity and uncouple respiration is thus determined.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are

10 related to SEQ ID NO:26 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

15 formula of a-b, where a is any integer between 1 to 1660 of SEQ ID NO:26, b is an integer of 15 to 1674, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:26, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 17 .

20 The translation product of this gene shares sequence homology with 55 kD deglycosylated zona pellucida protein which is known to be important in egg fertilization (See Genebank Accession No.R39356). Preferred polypeptides of the invention comprise the following amino acid sequence:

25 RPRPSASSLARSASLLPAAHGXGVGGAGGGSSXLRORYQQQLQNEEESGEPEQ
AAGDAPPPYSSISAESAHXFDYKDESGFPKPPSYNVATTLPSYDEAERTKAEA
TIPLVGRDEDFVGRDDFDDADQLRIGNDGIF (SEQ ID NO: 216) ,
RYQQQLQNEEESGEPEQAAGD (SEQ ID NO: 217) , and/or
PGRDEDFVGRDDFDDADQLRIG (SEQ ID NO: 218) . Polynucleotides encoding

30 these polypeptides are also provided.

Preferred polypeptide fragments of the invention comprise the following amino acid sequence: MLTFFMAFLFNWIGFFLSFCLTTSAAGRYG AISGFGLSLIKWILIVRFSTYFPGYFDGQY WLWWVFLVLGFLFLRGFINYAKVRKMPET FSNLPRTRVLFIY (SEQ ID NO: 219). Polynucleotides encoding these polypeptides are also provided.

Preferred polypeptide variants of the invention comprise the following amino acid sequence: MKKSLENLNRLQVMLLHLTA AFLQRAQHXYDYKDESGFPKPPSYNVATTLP SYDEAERTKAEATIPLVPGRDEDFVGRDDFDDADQLRIGNDGIFMLTFFMAFLF NWIGFFLSFCLTTSAAGRYGAISGFGLSLIKWILIVRFSTYFPGYFDGQYWLW WVFLVLGFLFLRGFINYAKVR KMPETFSNLPRTRVLFIY (SEQ ID NO: 220), MLLHLTA AFLQRAQFSTYFPGYFDGQYWLWWVFLVLGFLFLRGFINYAKV RKMPETFSN LPRTRVLFIY (SEQ ID NO: 221), MLTFFMAFLFNWIGFFLSFCLT TSAAGRYGAISGFGLSLIKWILIVRFSTYFPAFMNSLSRSKRTPAGSESR CRTQ RNNHLL (SEQ ID NO: 222), and/or MKKSLENLNRLQVMLLHLTA AFLQRAHXIL TTRMSLGFSQSPH LTM (SEQ ID NO: 223). Polynucleotides encoding these polypeptides are also provided.

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells, and to a lesser extent, other immune and hematopoietic cells and JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

This gene is expressed primarily in adult kidney, colon adenocarcinoma, and fetal brain, and to a lesser extent, ubiquitously expression in many tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, disorders of kidney, colon cancers, and central nervous system.

Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the renal and neural systems, and cancers, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., renal, neural, urogenital, or cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 119 as residues: Cys-15 to Gly-36. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution adult kidney, colon adenocarcinoma, and fetal brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of kidney diseases, colon cancers, and disorders of the central nervous system. Additionally, the homology to the zona pellucida protein indicates that the gene product is used for male contraceptive development, and infertility diagnosis etc. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:27 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1951 of SEQ ID NO:27, b is an

integer of 15 to 1965, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:27, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 18

5 The translation product of this gene shares sequence homology with the chicken transferrin receptor in addition to a human prostate-specific protein homolog (See Genebank Accession Nos. *pir*|JH0570|JH0570 and *gi*|2565338 (AF026380), respectively). This gene also shares significant homology with both the murine and the rat hematopoietic lineage switch 2 proteins (See Genbank Accession Nos. 10 *g*3169729 and *g*3851632, respectively), which are induced during an erythroid to myeloid lineage switch.

A preferred polypeptide fragment of the invention comprises the following amino acid sequence: MTVMDPKQMNVA^{AA}VWAVVSYVVADMEEML PRS (SEQ ID NO: 224). Polynucleotides encoding these polypeptides are also provided.

15 This gene is expressed primarily in fetal tissues, such as liver/spleen and brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, pre-natal disorders, anomalies, deficiencies. Similarly, polypeptides 20 and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the developing fetus, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., developing, cancerous and wounded tissues) or bodily fluids 25 (e.g., amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic 30 epitopes shown in SEQ ID NO: 120 as residues: Arg-31 to Lys-37, Lys-58 to Glu-65,

Asp-157 to Gly-168, Ile-219 to Gly-225, Ala-260 to Ser-268, Thr-276 to Glu-282.
Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of pre-natal disorders, anomalies and deficiencies. The homology to the hematopoietic lineage switch 2 proteins indicates that The translation product of this gene is useful for the detection and/or treatment of immune system disorders. In addition, the homology to the transferrin receptor indicates that the translation product of the present invention may have utility in the regulation of iron metabolism as well as the numerous genes under the stringent control of physiologic iron levels. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:28 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1849 of SEQ ID NO:28, b is an integer of 15 to 1863, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:28, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 19

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

PRVRSREPVAGAPGCGTAGPPAMATLWGGLRLRLGSLLSLSCLALSVLLLAHC
QTPPSDCLHVVEPMPVRGPDVEAYCLRCECKYEERSSVTIKVTIIYLSILGLLL
LYMVYLTLEVILKRRLFGHAQLIQSDDDIGDHQPFANAHDVLARSRSRANV

LNKVEYAAQQRWKLQVQEQRKSVFDRHVVLS (SEQ ID NO: 225).

Polynucleotides encoding these polypeptides are also provided.

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 72 - 88 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 89 to 167 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins.

A preferred polypeptide variant of the invention comprise the following amino acid sequence:

MATLWGGLRLGSLLSLSCLALSVLLLAHCQTPPRISMSDVNVSALPIKKNS
GHIYNKNISQKDCDCLHVVEPMPVRGPDVEAYCLRCECKYEERSSVTIKVTIII
YLSILGLLLLYMVYLTLEVILKRRRLFQHAQLIQSDDDIGDHQPFANAHDVLA
RSRSRANVLNKVEYGTAALASSPRAAKSLSLTGMLSSANWGIEFKVTRKKQ
ADNWKGTDWVLLGFILIPC (SEQ ID NO: 226). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in infant brain tissue, and to a lesser extent in other cell types and tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental and neurodegenerative diseases of the brain and nervous system, such as depression, schizophrenia, Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, mania, dementia, paranoia, addictive behavior, sleep disorders, epilepsy, transmissible spongiform encephalopathy (TSE), Creutzfeldt-Jakob disease (CJD). Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, developmental, or cancerous and wounded tissues) or bodily fluids (e.g., amniotic

fluid, lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 121 as residues: Gln-110 to Pro-120, Val-152 to Val-159. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in infant brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis
10 of developmental, degenerative and behavioral conditions of the brain and nervous system. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of schizophrenia, Alzheimer's Disease, Parkinson's
15 Disease, Huntington's Disease, Tourette Syndrome, transmissible spongiform encephalopathy (TSE), Creutzfeldt-Jakob disease (CJD), mania, depression, dementia, paranoia, addictive behavior, obsessive-compulsive disorder and sleep disorders. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to
20 identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
25 related to SEQ ID NO:29 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general
30 formula of a-b, where a is any integer between 1 to 1612 of SEQ ID NO:29, b is an

integer of 15 to 1626, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:29, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 20

5 The translation product of this gene shares sequence homology with a recently published gene Dysferlin, which is thought to be a skeletal muscle gene that is mutated in Miyoshi myopathy and limb girdle muscular dystrophy (See Genbank Accession No. g3600028, and Nat. Genet. 20 (1), 31-36 (1998)).

 This gene is expressed primarily in fetal liver, fetal heart tissue, and T-cells.

10 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immunodeficiency, tumor necrosis, lymphomas, auto-immunities, cancer, inflammation, anemias (leukemia) and liver disorders, vascular disorders, and
15 cancers (e.g., hepatoblastoma, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For
20 a number of disorders of the above tissues or cells, particularly of the liver and immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., hepatic, developmental, vascular, or cancerous and wounded tissues) or bodily fluids (e.g., amniotic fluid, bile, lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or
25 cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders including: leukemias, lymphomas, auto-immunities, immunodeficiencies
30 (e.g., AIDS), immuno-suppressive conditions (transplantation) and hematopoietic disorders. In addition this gene product is applicable in conditions of general

microbial infection, inflammation or cancer. Expression in liver may suggest a role for this gene product in the treatment and detection of liver disorders and cancers (e.g., hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). Alternatively, the tissue distribution in fetal heart tissue indicates that the protein product of this gene is useful for the diagnosis and treatment of conditions and pathologies of the cardiovascular system, such as heart disease, restenosis, atherosclerosis, stoke, angina, thrombosis, and wound healing. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Additionally, the homology to the dysferlin gene indicates that polynucleotides and polypeptides corresponding to this gene are useful for diseases related to degenerative myopathies that are characterized by the weakness and atrophy of muscles without neural degradation; such as Duchenne and Becker's muscular dystrophies. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:30 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 591 of SEQ ID NO:30, b is an integer of 15 to 605, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:30, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 21

This gene is expressed primarily in haemopoietic cells and tumor cells, such as pancreatic tumor tissue, and to a lesser extent in bladder cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, haemopoietic disorders, diseases of the renal and pancreatic systems, and cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the haemopoietic, pancreatic, and renal systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., pancreas, renal, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of disorders of the renal, pancreatic and haemopoietic systems, and cancers thereof. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:31 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 917 of SEQ ID NO:31, b is an integer of 15 to 931, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:31, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 22

This gene is expressed primarily in liver tissue, cancer cells and fetal lung tissue, and to a lesser extent in dendritic cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, metabolic disorders, diseases of developing systems and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the fetus, metabolic systems and cancers, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., developing, metabolic, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 124 as residues: His-44 to Gly-49. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of disorders of the fetus, metabolic systems and cancers. The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of liver disorders and cancers (e.g. hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). In addition the expression in fetus would suggest a useful role for the protein product in developmental abnormalities, fetal deficiencies, pre-natal disorders and various wound-healing models and/or tissue trauma. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:32 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1393 of SEQ ID NO:32, b is an integer of 15 to 1407, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:32, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 23

This gene is expressed primarily in central nervous system tissues and cancers, such as endometrial tumors, and to a lesser extent in other tissues and organs.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the CNS and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system and cancerous tissues, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 125 as residues: Tyr-16 to Ser-22, Asp-209 to Leu-215. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in central nervous system tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of diseases of the central nervous system, as well as cancers of tissues where expression of this gene has been observed, such as in
5 endometrial tumors. The tissue distribution in central nervous system tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive
10 disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, or sexually-linked disorders. Protein, as well as, antibodies directed against the protein
15 may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:33 and may have been publicly available prior to conception of
20 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1512 of SEQ ID NO:33, b is an
25 integer of 15 to 1526, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:33, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 24

The translation product of this gene shares sequence homology with low-
30 density lipoprotein receptor (See Genbank Accession No. >dbj|BAA24580.1), which is thought to be important in the pathogenesis of atherosclerosis and other disorders.

The translation product of this gene also shares sequence homology with a rat homolog of the human CD94 (See Genbank Accession No. gb|AAC10220.1).

This gene is expressed primarily in macrophages, eosinophils, neutrophil and other cells of the haemopoietic and immune system.

5 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the immune and haemopoietic systems and diseases of the endothelial and vascular system. Similarly, polypeptides and antibodies directed to
10 these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune, haemopoietic and vascular system, expression of this gene at significantly higher or lower levels is routinely detected in
15 certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic
20 epitopes shown in SEQ ID NO: 126 as residues: Lys-9 to Ala-17, Met-55 to Leu-61, Tyr-105 to Cys-127, Asp-132 to Lys-141, Ser-165 to Tyr-172, Pro-178 to Met-186, His-222 to Gln-227. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution and homology to LDL receptor and rat CD94 homolog indicates that polynucleotides and polypeptides corresponding to this gene are useful
25 for the treatment and/or diagnosis of disorders of the immune, haemopoietic and vascular systems. The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or
30 differentiation of hematopoietic lineages. Expression of this gene product in eosinophils and macrophage also strongly indicates a role for this protein in immune

function and immune surveillance. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:34 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1723 of SEQ ID NO:34, b is an integer of 15 to 1737, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:34, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 25

A preferred polypeptide fragment of the invention comprises the following amino acid sequence:

MAAAGRLPSSWALFSPLLAGLALLGVGPVPARALHNVTAELFGAEAWGTLA
AFGDLNSDKQTDLFVLRERNDLIVFLADQNAPYFKPKVKVSFKNHSALITSVV
20 PGDYDGDSQMDVLLTYLPKNYAKSELGAVIFWGQNQTLDPNMNTILNRTFQ
DEPLIMDFNGDLIPDIFGITNESNQPIQLLGGNLSWHPALTTTSKMRIPHSHAFI
DLTEDFTADLFLTTLNATTSTFQFEIWENLDGNFSVSTILEKPQNMMVVGQSA
FADFDGDGHMDHLLPGCEDKNCQKSTIYLVRSGMKQWVPVLQDFSNGKTL
WGFVFPVDEQQPTEIPIITLHIGDYNMDGYPDALVILKNTSGSNQQAFLLENV
25 PCNNASCEEARRMFKVYWELTDLNQIKDAMVATFFDIYEDGILDIVVLSKGY
TKNDFAIHTLKNFEADAYFVKVIVLSGLCS NDCPRR (SEQ ID NO: 227).

Polynucleotides encoding these polypeptides are also provided.

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter
30 element. Thus, it is likely that this gene activates myeloid cells, and to a lesser extent, other immune and hematopoietic cells and tissue cell types, through the JAK-STAT

signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the
5 GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

This gene is expressed primarily in infant brain and placental tissues, and to a lesser extent in several other tissues including cancers.

Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, brain disorders and diseases of developing systems and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell
15 type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system and fetal systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, developing, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken
20 from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 127 as residues: Leu-56 to Thr-62, Gln-80 to Pro-87,
25 Gly-106 to Gln-113, Pro-122 to Lys-127, Gln-138 to Asn-146. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in neural tissues and developing tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of disorders of the central nervous system, disorders of
30 developing systems, and cancers. The tissue distribution in infant brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful

for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered
5 behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, or sexually-linked disorders.

The tissue distribution indicates that polynucleotides and polypeptides
10 corresponding to this gene are useful for the diagnosis and/or treatment of disorders of the placenta. Specific expression within the placenta indicates that this gene product may play a role in the proper establishment and maintenance of placental function. Alternately, this gene product is produced by the placenta and then transported to the embryo, where it may play a crucial role in the development and/or
15 survival of the developing embryo or fetus. Expression of this gene product in a vascular-rich tissue such as the placenta also indicates that this gene product is produced more generally in endothelial cells or within the circulation. In such instances, it may play more generalized roles in vascular function, such as in angiogenesis. It may also be produced in the vasculature and have effects on other
20 cells within the circulation, such as hematopoietic cells. It may serve to promote the proliferation, survival, activation, and/or differentiation of hematopoietic cells, as well as other cells throughout the body. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

25 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:35 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is
30 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

formula of a-b, where a is any integer between 1 to 2228 of SEQ ID NO:35, b is an integer of 15 to 2242, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:35, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 26

Preferred polypeptides of the invention comprise the following amino acid sequence:

MTKREDGGYTFTATPEDFPKKHKAPVIDIGIANTGKFIMTASSDTTVLIWSLK
GQVLSTINTNQMNNTAAVSPCGRFVASCFTPDVKVWEVCFGKKGEFQEV
10 VRAFELKGHSAAVHSFAFSNDSRRMASVSKDGTWKLWDTXVEYKKKQDPY
LLKTGRFEEAAGAXPCRLALSPNAQVLALASGSSIHLYNTRRGEKEECFERVH
GECIANLSFDITGRFLASCGDRAVRLFHNTPGHRAMVEEMQGHLKRASNEST
RQRLQQQLTQAQETLKS LGALKK (SEQ ID NO: 228). Polynucleotides encoding
such polypeptides are also provided.

15 The gene encoding the disclosed cDNA is thought to reside on chromosome 7. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 7.

When tested against U937 Myeloid cell lines, supernatants removed from cells containing this gene activated the GAS assay. Thus, it is likely that this gene activates
20 myeloid cells through the Jak-STAT signal transduction pathway. The gamma activating sequence (GAS) is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS
25 element, can be used to indicate proteins involved in the proliferation and differentiation of cells. Recently, another group published this gene, naming it WS beta-transducin repeats protein (See Human Genetics 103 (5), 590-599 (1998); which is hereby incorporated herein by reference), in which it was suggested that the protein is involved in William's Disease.

30 The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 12 - 28 of the amino acid sequence referenced in

Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ib membrane proteins.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

VIRHEGSTNMELSQMSXLMGLSVLLGLLALMATAAVXRGWLRAGEERSGRP
ACQKANGFPPDKSSGSKKQKQYQRRKEKPQQHNFTHRLAAALKSHSGNIS
CMDFSSNGKYLATCADDRTIRIWSTKDFLQREHRSMRANVELDHATLVRFP
10 DCRAFIWVLANGDTRLRVFKMTKREDGGYTFTATPEDFPKKHKAPVIDIGIAN
TGK
FIMTASSDTTVLIWSLKGQVLSTINTNQMNNTAAVSPCGRFVASCFTPDVK
VWEVCFGKKGEFQEVVRAFELKGHSAAVHSFAFSNDSRRMASVSKDGTWK
LWDTXVEYKKKQDPYLLKTGRFEEAAGAXPCRLALSPNAQVLALASGSSIHL
15 YNTRRGEKEECFERVHGECIANLSFDITGRFLASCGDRAVRLFHNTPGHRAM
VEEMQGHKLRASNESTRQLQQLTQAQETLKS LGALKK (SEQ ID NO: 229).

Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in testes, synovial sarcoma and fetal tissues, and to a lesser extent in several other tissues.

20 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the reproductive and developing systems and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in
25 providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive and developing systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., reproductive, testicular, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum,
30 plasma, seminal fluid, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene

expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in testes tissue, synovial sarcoma, and fetal tissues, indicates that polynucleotides and polypeptides corresponding to this gene are useful
5 for the treatment and/or diagnosis of disorders of the reproductive and developing systems, and cancers. The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of conditions concerning proper testicular function (e.g. endocrine function, sperm
10 maturation), as well as cancer. Therefore, this gene product is useful in the treatment of male infertility and/or impotence. This gene product is also useful in assays designed to identify binding agents, as such agents (antagonists) are useful as male contraceptive agents. Similarly, the protein is believed to be useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene
15 expression of transcripts that is expressed, particularly at low levels, in other tissues of the body. Therefore, this gene product is expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications.

Furthermore, the tissue distribution indicates that polynucleotides and
20 polypeptides corresponding to this gene are useful for the diagnosis and treatment of cancer and other proliferative disorders. Expression within embryonic tissue and other cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation of cellular division. Additionally, the expression in hematopoietic cells and tissues indicates that this protein may play a role in the
25 proliferation, differentiation, and/or survival of hematopoietic cell lineages. In such an event, this gene is useful in the treatment of lymphoproliferative disorders, and in the maintenance and differentiation of various hematopoietic lineages from early hematopoietic stem and committed progenitor cells. Similarly, embryonic development also involves decisions involving cell differentiation and/or apoptosis in
30 pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. This protein is useful for

the treatment, detection, and/or prevention of William's Disease. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:36 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2221 of SEQ ID NO:36, b is an integer of 15 to 2235, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 27

Preferred polypeptides of the invention comprise the following amino acid sequence positions 1-363, 2-363, 4-363, 5-363, 30-363, 31-363, 32-363, 75-363, 76-363 and 82-363 of the following amino acid sequence:

MSVMVVRKKVTRKWEKLPGRNTFCCDGRVMMARQKGIFYLTLFLILGTCTL
FFAFECRYLAVQLSPAIPVFAAMLFLFSMATLLRTSFSDPGVIPRALPDEAA
MEIEATNGAVPQGQRPPRIKNFQINNQIVKLKYCYTCKIFRPPRASHCSICDN
CVERFDHHCPCWVGNCVGKRNRYFYFLFILSLSLTIYVFAFNIVYVALKSLKI
GFLETLKETPGTVLEVLCFFTLWSVVGLTGFHTFLVALNQTNEDIKGSWTG
KNRVQNPYSHGNIVKNCCEVLCGPLPPSVLDRRGILPLEESGSRPPSTQETSSS
LLPQSPAPTEL NSNEMPEDSSTPEEMPPEPPEPPQEAAEAEK (SEQ ID NO:
230). Polynucleotides encoding such polypeptides are also provided.

A preferred polypeptide variant of the invention comprises the following amino acid sequence: MLFLFSMATLLRTSFSDPGVIPRALPDEAA

FIEMEIEATNGAVPQGQRPPRIKNFQINNQIVKLKCYTCKIFRPPRASHCSIC
DNCVE RFDHHCPWVGNCVGKRNRYFYLFILSLSLTIYVFAFNIVYVALK
SLKIGFLETLKGNS WNCSRSPHLLLYTLVRRGTDWISYFPRGSQ PDNQ (SEQ
ID NO: 231). Polynucleotides encoding these polypeptides are also provided.

5 This gene is expressed primarily in ovarian and endometrial tumors, fetal liver, spleen and brain tissues, and to a lesser extent in several other tissues and organs.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
10 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the developing systems, and cancers of the female reproductive system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above
15 tissues or cells, particularly of the developing, female reproductive and fetal systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., reproductive, developing, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a
20 disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 129 as residues: Pro-44 to Lys-54, Cys-88 to His-95, Val-103 to Tyr-108, Gln-181 to Ser-190, Thr-192 to Ile-206, Glu-233 to Ser-245, Ser-
25 252 to Ala-286. Polynucleotides encoding said polypeptides are also provided.

 The tissue distribution in developing systems indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of disorders of developing and fetal systems, and cancers. Furthermore, the tissue distribution in ovarian and endometrial tumor tissues indicates that the
30 translation product of this gene is useful for the detection, diagnosis, and/or treatment of cancers of the female reproductive system. Accordingly, preferred are antibodies

which specifically bind a portion of The translation product of this gene. Also provided is a kit for detecting these tumors. Such a kit comprises in one embodiment an antibody specific for The translation product of this gene bound to a solid support. Also provided is a method of detecting these tumors in an individual which

5 comprises a step of contacting an antibody specific for The translation product of this gene to a bodily fluid from the individual, preferably serum, and ascertaining whether antibody binds to an antigen found in the bodily fluid. Preferably the antibody is bound to a solid support and the bodily fluid is serum. The above embodiments, as well as other treatments and diagnostic tests (kits and methods), are

10 more particularly described elsewhere herein.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:37 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

15 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2957 of SEQ ID NO:37, b is an integer of 15 to 2971, where both a and b correspond to the positions of nucleotide

20 residues shown in SEQ ID NO:37, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 28

This gene is expressed primarily in normal and cancerous colon tissue, macrophages, endothelial cells and placental tissue, and to a lesser extent in several

25 other tissues and organs.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, colon cancer and gastrointestinal disorders, immune disorders, vascular

30 diseases and disorders of developing systems. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for

differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune, vascular and developing systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, gastrointestinal, developmental, 5 vascular, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

10 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 130 as residues: Thr-27 to Ser-33. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in macrophage, endothelial and placental tissues, and normal and cancerous colon tissues, indicates that polynucleotides and polypeptides 15 corresponding to this gene are useful for the treatment and/or diagnosis of immune, gastrointestinal and vascular disorders and diseases. Expression of this gene product in colon tissue indicates involvement in digestion, processing, and elimination of food, as well as a potential role for this gene as a diagnostic marker or causative agent in the development of colon cancer, and cancer in general. Accordingly, preferred are 20 antibodies which specifically bind a portion of the translation product of this gene. Also provided is a kit for detecting colon cancer. Such a kit comprises in one embodiment an antibody specific for The translation product of this gene bound to a solid support. Also provided is a method of detecting colon cancer in an individual which comprises a step of contacting an antibody specific for The translation 25 product of this gene to a bodily fluid from the individual, preferably serum, and ascertaining whether antibody binds to an antigen found in the bodily fluid. Preferably the antibody is bound to a solid support and the bodily fluid is serum. The above embodiments, as well as other treatments and diagnostic tests (kits and methods), are more particularly described elsewhere herein. Alternatively, the tissue 30 distribution in placental tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of disorders

of the placenta. Specific expression within the placenta indicates that this gene product may play a role in the proper establishment and maintenance of placental function. Alternately, this gene product is produced by the placenta and then transported to the embryo, where it may play a crucial role in the development and/or survival of the developing embryo or fetus. Expression of this gene product in a vascular-rich tissue such as the placenta and endothelial cells also indicates that this gene product is produced more generally in endothelial cells or within the circulation. In such instances, it may play more generalized roles in vascular function, such as in angiogenesis. It may also be produced in the vasculature and have effects on other cells within the circulation, such as hematopoietic cells. It may serve to promote the proliferation, survival, activation, and/or differentiation of hematopoietic cells, as well as other cells throughout the body. Additionally, expression of this gene product in macrophage also strongly indicates a role for this protein in immune function and immune surveillance. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:38 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1149 of SEQ ID NO:38, b is an integer of 15 to 1163, where both a and b correspond to the positions of nucleotide
5 residues shown in SEQ ID NO:38, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 29

The translation product of this gene shares homology with HNK-sulfotransferase, which directs glycan synthesis (see Genbank Accession no.
10 AF033827).

This gene is expressed primarily in activated T cells, osteoclastoma, and glioblastoma, and to a lesser extent in various other normal and transformed cell types.

Therefore, polynucleotides and polypeptides of the invention are useful as
15 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, inflammation, immune defects, cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of
20 disorders of the above tissues or cells, particularly of the immune and hemopoietic systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative
25 to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 131 as residues: Pro-32 to Gly-48, Gln-63 to Thr-69, Pro-77 to Trp-84, Val-88 to Leu-94. Polynucleotides encoding said polypeptides are
30 also provided.

The tissue distribution in T-cells and various types of neoplasms indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, study and/or treatment of inflammatory and general immune defects, and various types of neoplasms. Expression of this gene product in T cells strongly
5 indicates a role for this protein in immune function and immune surveillance. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the gene or protein, as
10 well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility
15 in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Alternatively, the tissue distribution in various cancerous tissues indicates that the translation product of the gene is useful for the detection, diagnosis, and/or treatment of these cancers, as well as cancers of other tissues where expression has been observed. Protein, as well
20 as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:39 and may have been publicly available prior to conception of
25 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1918 of SEQ ID NO:39, b is an
30 integer of 15 to 1932, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:39, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 30

Preferred polypeptides of the invention comprise the following amino acid sequence:

5 LHECLPGSISYLHP RTPWLCLPPQHLSFSTFSPPWQPAMSPVPGTGGPPCGL
(SEQ ID NO: 232), and/or
MLPLLICLLPAIEGKNCLRCWPELSALIDYDLQILWVTPGPPTELSQSIHSLFLE
DNNFLKWPYLD RDHLEEEETAKFFTQVHQA IKT LRDDKT VLL EEIYTHKNLFT
ERLNKISDGLKEKGAPPLHECLPGSISYLHP RTPWLCLPPQHLSFSTFSPPWQP

10 AMSPVPGTGGPPCGL (SEQ ID NO: 233). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in infant brain, testes and activated T cells, and to a lesser extent in various other normal and transformed cell types.

Therefore, polynucleotides and polypeptides of the invention are useful as
15 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological, reproductive and inflammatory conditions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For
20 a number of disorders of the above tissues or cells, particularly of the neural, immune and male reproductive systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, immune, reproductive, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken
25 from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 132 as residues: Gly-41 to Leu-46, Asp-67 to Thr-75,
30 Ile-114 to Ala-123. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in infant brain tissue, testes tissue, and activated T-cells, indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study, diagnosis, and/or treatment of neurological, reproductive and immune system disorders. Expression of this gene product in T-cells indicates a role
5 in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses).

10 Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it is also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease,
15 sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell type. Alternatively, the tissue distribution in testes tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of conditions
20 concerning proper testicular function (e.g. endocrine function, sperm maturation), as well as cancer. Therefore, this gene product is useful in the treatment of male infertility and/or impotence. This gene product is also useful in assays designed to identify binding agents, as such agents (antagonists) are useful as male contraceptive agents. Similarly, the protein is believed to be useful in the treatment and/or diagnosis
25 of testicular cancer. The testes are also a site of active gene expression of transcripts that is expressed, particularly at low levels, in other tissues of the body. Therefore, this gene product is expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications.

30 Furthermore, the tissue distribution in infant brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the

detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, or sexually-linked disorders. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:40 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 867 of SEQ ID NO:40, b is an integer of 15 to 881, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:40, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 31

The translation product of this gene shares sequence homology with some human and rodent melanoma and leukocyte specific antigens (see, for example, Genbank accession nos: gi|189384, gi|205898 and gi|180926). In addition, The translation product of this gene shares sequence homology with Tetraspan protein (see, for example, Genbank accession number: GI 3152703). Therefore, it is likely that the polypeptide of this gene shares some biological functions, such as cell-to-cell signaling, adhesion, proliferation, and differentiation with Tetraspan.

The polypeptide of this gene has been determined to have two transmembrane domains at about amino acid position 52-68 and 197 - 213 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed

that the protein product of this gene shares structural features to type IIIa membrane proteins.

The transmembrane 4 superfamily (TM4SF) or tetraspan superfamily has at least 16 members (including CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8), is the second biggest subfamily among CD antigen superfamily. and activation antigen of T- cells. All TM4SF member contains four putative transmembrane domains, two extracellular loops, and two short cytoplasmic tails. They are variously expressed on Immature, early, mature, activated lymphocytes, monocytes, macrophages, granulocytes, platelets, eosinophils, basophils, certain leukemic and lymphoma cells, and a variety of other cells and tissues. CD9 cell surface protein is expressed by both hematopoietic and neural cells, and may play a role for CD9 in intercellular signaling in the immune and nervous system. CD63 is a 53-Kd lysosomal membrane glycoprotein that has been identified as a platelet activation molecule, which play important role in cell adhesion of platelets and endothelial cells. Increased mRNA for CD63 antigen was found in atherosclerotic lesions of Watanabe heritable hyperlipidemic rabbits, suggesting a potential role of CD63 in progression of atherosclerosis. CD63 is also a mast cell marker.

CD82 was originally identified as the target of several mAbs inhibitory to syncytium formation induced by human T-cell leukemia virus type I (HTLV-I), the etiological agent of adult T-cell leukemia. Therefore, this gene could be a target for the development of a drug for this leukemia. CD81 is the target of an antiproliferative antibody. A diverse group of human cell lines, including hematolymphoid, neuroectodermal, and mesenchymal cells, express the CD81 protein. Many of the lymphoid cell lines, in particular those derived from large cell lymphomas, were susceptible to the antiproliferative effects of the antibody. CD81 may therefore play an important role in the regulation of lymphoma cell growth. CD9, CD20, CD37, CD63, CD81 and CD82 have been implicated in the regulation of cell growth, adhesion, and signal transduction of B, T lymphocytes and some other non-lymphoid cells. They associate with CD2, CD21, CD4, CD8, MHC Class II molecules, integrins, function as co-receptor for T, B and other lymphoid cells. Some

TM4SF are leukocyte antigens, highly expressed in activated leukocytes, lymphocytes, are highly specific surface marker for lymphoblastic leukemia, lymphoma, melanoma, and neuroblastoma. CD9 has been show to be involved in cell motility and tumor metastasis. These antigen could be a valuable immunogen or
5 target to implement active and passive immunotherapy in patients with cancer. Others have been shown to be involved in inhibition of prostate cancer metastasis. This gene has close homology to C33 antigen (CD82). whic is a member of the transmembrane 4 superfamily (TMSF) and activation antigen of T- cells. C33 Ag (CD82 was originally identified as the target of several mAbs inhibitory to syncytium formation
10 induced by human T-cell leukemia virus type I (HTLV-I), the etiological agent of adult T-cell leukemia. Therefore, this gene could be very important target for developing drug for leukemia. Other members of this family are Sm23, CO-029, R2, TAPA-1, CD9, CD37, CD53, and CD63. CD63 is a 53-Kd lysosomal membrane glycoprotein that has been identified as a platelet activation molecule.

15 There is strong evidence indicating that CD63 and Pltgp40, a platelet membrane glycoprotein are the same molecule and that CD63/Pltgp40 is identical to the well-characterized, stage-specific melanoma-associated antigen ME491. These antigen could be valuable immunogens or target to implement active and passive immunotherapy in patients with cancer.

20 This gene is expressed primarily in fetal tissue (kidney, heart, liver, spleen, brain), macrophages, dendritic cells, retina and to a lesser extent in various other tissues, mostly of lymphoid origin or epithelial cell types. In addition This gene is expressed in cancerous tissues (e.g. breast).

 Therefore, polynucleotides and polypeptides of the invention are useful as
25 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic diseases and/or disorders and cancers in a variety of organs and cell types. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential
30 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at

significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., developmental, proliferating, immune, hematopoietic, integumentary, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid, spinal fluid, or amniotic fluid) or another tissue or cell sample taken
5 from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 133 as residues: Tyr-123 to Tyr-131, Cys-134 to Ser-
10 145, Tyr-234 to Tyr-244. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution fetal cells and tissues and homology to tumor antigens indicates that polynucleotides and polypeptides corresponding to this gene are useful for study, treatment and diagnosis of lymphoid and epithelial disorders and
15 neoplasms. Additionally, tissue distribution in immune cells and other tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting hematopoiesis, including cancers. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere
20 herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

25 Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities,
30 such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity

disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits
5 hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types.

Moreover, the expression within embryonic tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of
10 cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern
15 formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of
20 potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types
25 of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new
30 insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue

markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:41 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is
10 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1918 of SEQ ID NO:41, b is an integer of 15 to 1932, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:41, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 32

The translation product of this gene shares limited sequence homology with VEGF which is thought to be important in regulation of endothelial cell growth. Therefore, it is likely that the protein encoded by this gene would share some similar
20 biological functions.

When tested against U937 Myeloid cell lines, supernatants removed from cells containing this gene activated the GAS assay. Thus, it is likely that this gene activates myeloid cells, and to a lesser extent, other immune and hematopoietic cells and tissue cell types, through the Jak-STAT signal transduction pathway. The gamma activating
25 sequence (GAS) is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and
30 differentiation of cells.

This gene is expressed in brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, nervous system disease and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 134 as residues: Thr-25 to Pro-46. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in brain indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to
5 identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
10 related to SEQ ID NO:42 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
15 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1150 of SEQ ID NO:42, b is an integer of 15 to 1164, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:42, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 33

20 The translation product of this gene shares sequence homology with human p150 which is thought to be important in signal transduction in neuronal cells. Therefore, it is likely that the protein encoded by this polynucleotide would share some similar biological functions with p150.

This gene is expressed primarily in whole embryo and cerebellum.

25 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological and growth defects/disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological
30 probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the CNS, expression of this gene

at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene
5 expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for study and treatment of central nervous system, neurodevelopmental, cognitive, and memory disorders. The tissue distribution
10 also indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not
15 limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive
20 disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation,
25 neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show
30 utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Moreover, the expression within embryonic tissue and other cellular sources marked

by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:43 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1091 of SEQ ID NO:43, b is an integer of 15 to 1105, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:43, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 34

This gene is expressed primarily in PMA stimulated HL-60 cells and to a lesser extent in 6 week embryo.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders affecting cell differentiation, particularly hematopoietic disorders and/or defects. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the metabolic system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 136 as residues: Pro-61 to Asp-68. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study of cellular differentiation and for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia. The tissue distribution also

indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the

5 expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses). Moreover, the protein may

10 represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Additionally, the expression within embryonic tissue and other

15 cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the

"Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere

20 herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain

25 neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating,

30 detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue

differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:44 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1248 of SEQ ID NO:44, b is an integer of 15 to 1262, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:44, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 35

This gene is expressed primarily in colon.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders and/or defects of the digestive tract including but not limited to cancers of the gastrointestinal tract. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the digestive system, expression of this gene at

significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., gastrointestinal, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of disorders of the digestive system particularly disorders involving the colon. Further, expression of this gene product in colon tissue indicates involvement in digestion, processing, and elimination of food, as well as a potential role for this gene as a diagnostic marker or causative agent in the development of colon cancer, and cancer in general. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the colon and/or other gastrointestinal tissue including, but not limited to, stomach, small intestine, large intestine, and rectum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:45 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 503 of SEQ ID NO:45, b is an integer of 15 to 517, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:45, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 36

This gene is expressed primarily in blood cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, immune and hematopoietic diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and hematopoietic system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 138 as residues: Pro-19 to Cys-29, Thr-35 to Glu-44, Val-72 to Lys-78. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis and/or treatment of disorders of the immune and hematopoietic system. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:46 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 844 of SEQ ID NO:46, b is an

integer of 15 to 858, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:46, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 37

5 This gene is expressed in multiple tissue systems such as brain, immune cells, prostate, uterus, testes, placenta, and fetal heart as well as in cancerous tissues such as ovarian tumors. .

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
10 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the immune, reproductive, urogenital, and central nervous system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells,
15 particularly of the central nervous system and immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, reproductive, urogenital, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative
20 to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 139 as residues: Tyr-33 to Lys-38. Polynucleotides encoding said polypeptides are also provided.

25 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of disorders of the immune, urogenital, reproductive, and central nervous systems. The tissue distribution in central nervous system tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis
30 of diseases of the central nervous system, as well as cancers of tissues where expression of this gene has been observed, such as in ovarian tumors. The tissue

distribution in central nervous system tissues also indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Moreover, the expression within embryonic tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in

proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution in uterus indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating female infertility. The protein product is likely involved in preparation of the endometrium of implantation and could be administered either topically or orally. Alternatively, this gene could be transfected in gene-replacement treatments into the cells of the endometrium and the protein products could be produced. Similarly, these treatments could be performed during artificial insemination for the purpose of increasing the likelihood of implantation and development of a healthy embryo. In both cases this gene or its gene product could be administered at later stages of pregnancy to promote healthy development of the endometrium. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution in testes indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of conditions concerning proper testicular function (e.g. endocrine function, sperm maturation), as well as cancer. Therefore, this gene product is useful in the treatment of male infertility and/or impotence. This gene product is also useful in assays designed to identify binding agents, as such agents (antagonists) are useful as male contraceptive agents. Similarly, the protein is believed to be useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene expression of transcripts that is expressed, particularly at low levels, in other tissues of the body. Therefore, this gene product is expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications. Protein, as well as, antibodies directed

against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:47 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 6093 of SEQ ID NO:47, b is an integer of 15 to 6107, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:47, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 38

This gene is expressed in a wide range of tissue systems such as brain, immune cells, fetal liver, kidney, testes, breast, and pancreas as well as cancerous tissue such as ovarian tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the central nervous system, immune system, urogenital, and reproductive system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and central nervous system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, CNS, urogenital, reproductive, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 140 as residues: Met-1 to Ser-7, Asp-32 to Pro-43, Ser-96 to Arg-102. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides
5 corresponding to this gene are useful for treatment and diagnosis of disorders of the immune, reproductive, urogenital and central nervous systems. The tissue distribution in central nervous system tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of diseases of the central nervous system, as well as cancers of tissues where
10 expression of this gene has been observed, such as in ovarian tumors. The tissue distribution in central nervous system tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome,
15 schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. The tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders.
20 Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation
25 of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such
30 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities,

such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Moreover, the expression within embryonic tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue

differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases.

The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:48 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 689 of SEQ ID NO:48, b is an integer of 15 to 703, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:48, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 39

This gene is expressed primarily in macrophages and fetal cells and to a lesser extent in cancerous ovarian tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune diseases, disorders of developing tissues, and cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in

providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the fetal and immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of developmental abnormalities and disorders of the immune systems. The tissue distribution cancerous ovaries indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and intervention of these tumors. Protein, as well as, antibodies directed against the protein may show utility as a tissue-specific marker and/or immunotherapy target for the above listed tissues. Expression of this gene product in macrophage cells strongly indicates a role for this protein in immune function and immune surveillance. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). This gene product may have clinical utility in the treatment of immune dysfunction; in the correction of autoimmunity; in immune modulation; and in the control of inflammation.

The tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or

other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types.

Moreover, the expression within embryonic tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the

polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of
5 degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue
10 markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution also indicates that polynucleotides and polypeptides corresponding to this gene are useful
15 for the treatment, diagnosis, and/or prevention of various skin disorders such as melanomas.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:49 and may have been publicly available prior to conception of
20 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 625 of SEQ ID NO:49, b is an
25 integer of 15 to 639, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:49, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 40

This gene is expressed primarily in neutrophils, bone marrow, brain, and fetal
30 cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hematopoietic disorders, Limbic system disfunction/defects and disorders of the immune system and developing systems. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune, Limbic system and developing systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 142 as residues: Ala-84 to Gln-93. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of disorders of the immune, Limbic system, CNS and developing systems. Expression of this gene product in bone marrow, eosinophils, and neutrophils strongly indicates a role for this protein in hematopoiesis and immune surveillance. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). This gene product may have clinical utility in the treatment of immune dysfunction; in the correction of autoimmunity; in immune modulation; and in the control of inflammation. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune

Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury.

Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Additionally, the expression within embryonic tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the

"Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal-muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:50 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 853 of SEQ ID NO:50, b is an

integer of 15 to 867, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:50, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 41

5 This gene is expressed primarily in ovary and to a lesser extent in fetal tissue, colon, and immune cells.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
10 not limited to, ovarian cancer, gastrointestinal and immune system disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the female reproductive system, expression of this gene at significantly higher or lower
15 levels is routinely detected in certain tissues or cell types (e.g., reproductive, gastrointestinal, immune, cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an
20 individual not having the disorder.

 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 143 as residues: Ile-23 to Ala-29. Polynucleotides encoding said polypeptides are also provided.

 The tissue distribution indicates that polynucleotides and polypeptides
25 corresponding to this gene are useful for diagnosis and treatment of ovarian cancer and related metastases. The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating female infertility. The tissue distribution in colon tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of disorders
30 involving the gastrointestinal tract. This may include diseases associated with digestion and food absorption, as well as hematopoietic disorders involving the

Peyer's patches of the small intestine, or other hematopoietic cells and tissues within the body. Similarly, expression of this gene product in colon tissue indicates again involvement in digestion, processing, and elimination of food, as well as a potential role for this gene as a diagnostic marker or causative agent in the development of colon cancer, and cancer in general. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Moreover, the expression within embryonic tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their

interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
5 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:51 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
10 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1555 of SEQ ID NO:51, b is an integer of 15 to 1569, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:51, and where b is greater than or equal to a + 14.

15 **FEATURES OF PROTEIN ENCODED BY GENE NO: 42**

The translation product of this gene shares sequence homology with retrovirus-related reverse transcriptase pseudogene. In addition, this gene shares homology with human interferon-beta (Genseq accession number T35524; all references available through this accession are hereby incorporated herein by
20 reference), therefore, it is likely that this gene and the protein encoded by this gene shares some similar biological functions with this protein.

This gene is expressed primarily in frontal cortex.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
25 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system,
30 expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, cancerous and wounded tissues) or bodily

fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 The tissue distribution in frontal cortex and homology to retrovirus-related reverse transcriptase pseudogene and human interferon-beta indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neurodegenerative diseases of the brain, particularly of the frontal cortex. The tissue distribution indicates polynucleotides and polypeptides
10 corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of
15 Alzheimer's Disease, Parkinson's Disease, multiple sclerosis, cystic fibrosis, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder,
20 learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

 Potentially, this gene product is involved in synapse formation,
25 neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show
30 utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:52 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1182 of SEQ ID NO:52, b is an integer of 15 to 1196, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:52, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 43

This gene is expressed primarily in immune cells, brain, fetal tissue, and cancerous tissues (such as testes, stomach, lung, pancreas, ovaries) and to a lesser extent in other numerous tissues including, but not limited to, testes and kidney.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system and immune cells expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 145 as residues: Lys-23 to Lys-35, Met-46 to Tyr-52. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neurodegenerative disorders of the frontal cortex, as well as, cancer or a number of tissues including but not limited to testes, stomach, lung, pancreas, and ovaries. The tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell

lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Moreover, the expression within embryonic tissue and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent

of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:53 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 931 of SEQ ID NO:53, b is an integer of 15 to 945, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:53, and where b is greater than or equal to a + 14.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 44

This gene is expressed primarily in epithelioid sarcoma and to a lesser extent in pancreatic carcinoma, aorta endothelial cells induced with TNF-alpha, and amniotic cells induced with TNF. This gene is also expressed, to a lesser extent, in cancerous lung and ovary tissue and fetal tissue.

5 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, epithelioid sarcoma and related cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological
10 probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., amniotic, lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another
15 tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 146 as residues: Tyr-39 to Arg-51. Polynucleotides
20 encoding said polypeptides are also provided.

 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of certain cancers, including epithelioid sarcoma and pancreatic carcinoma. The tissue distribution in tumors of lung, ovary, and pancreas origins indicates that polynucleotides and
25 polypeptides corresponding to this gene are useful for the diagnosis and intervention of these tumors, in addition to other tumors where expression has been indicated. Protein, as well as, antibodies directed against the protein may show utility as a tissue-specific marker and/or immunotherapy target for the above listed tissues. Moreover, the expression within embryonic tissue and other cellular sources marked
30 by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of

developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The tissue distribution indicates polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including

blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:54 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 474 of SEQ ID NO:54, b is an

integer of 15 to 488, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:54, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 45

5 In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

PPVPPWISLPLTGSPRPFGFVPVSPFCFSPMTNGHQVLLLLLLTSAVAAGPWPQ

10 VHAGQWGWMLPPGLPSVQARSLGGLPGGPQWVPGGARGY (SEQ ID NO: 234). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in fetal and infant tissue, particularly infant brain and fetal liver/spleen libraries, and to a lesser extent in breast, ovary tumor, pharynx carcinoma, endometrial stromal cells, thymus, islet cell tumors, and adult
15 cerebellum.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cancer and other proliferative disorders. Similarly, polypeptides and
20 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain and breast, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, developmental, hematopoietic, and cancerous and
25 wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, amniotic fluid, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

30 The tissue distribution in developing cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis

and treatment of cancer and other proliferative disorders. The expression within cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:55 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general
 5 formula of a-b, where a is any integer between 1 to 2846 of SEQ ID NO:55, b is an integer of 15 to 2860, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:55, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 46

10 In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

15 IQQWGDSVLGRRCRDLLQLYLQRPRLRVPVPEVLLHSEGAASSSVCKLDGLI
 HRFITLLADTSDSRALENRGADASMCRKLAVAHPLLLLRHLPMAALLHGR
 THLNFEFRQQNHLSCFLHVLGLLELLQPHVFRSEHQGALWDCLLSFIRLLN
 YRKSSRHAAAFINKFVQFIHKYITYNAPAAISFLQKHADPLHDLSDNSDLVM
 LKSLLAGLSLPSRDDRTDRGLDEEGEEESSAGSLPLVSVSLFTPLTAAEMAPY
 MKRLSRGQTVEDLLEVLSDIDEMSRRRPEILSFFSTNLQRLMSSAECCRNLA
 20 FSLALRSMQNSPSIAAFLPTFMVCLGSQDFEVVQTALRNLPEYALLCQEHA
 AVLLHRAFLVGMYGQMDPSAQISEALRILHMEAVM (SEQ ID NO: 235).

Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in breast cancer, and to a lesser extent in a variety of other cancers, including uterine cancer, synovial sarcoma, and pharynx
 25 carcinoma.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, breast cancer; proliferative diseases and/or disorders. Similarly,
 30 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For

a number of disorders of the above tissues or cells, particularly of the breast, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., reproductive, breast, proliferative, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, breast milk, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 148 as residues: Glu-35 to His-41, Ser-62 to Ala-67, Pro-145 to Leu-155, Glu-157 to Ser-163, Arg-190 to Val-197, Asp-208 to Pro-215, Ser-247 to Pro-252. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in breast cancer tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of cancer. Elevated expression of this gene product in cancers, such as breast cancer, suggest that it is involved in the abnormal proliferation of cells, dedifferentiation, angiogenesis, and other processes that accompany the development of cancer. Thus, therapeutics targeted against this gene product is useful therapeutic products in and of themselves. Alternately, expression of this gene product at elevated levels in breast tissue is reflective of expression within breast lymph nodes, and may suggest a hematopoietic role for this protein. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the

polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of

5 degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue

10 markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly

15 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:56 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or

20 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1545 of SEQ ID NO:56, b is an integer of 15 to 1559, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:56, and where b is greater than or equal to a + 14.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 47

The translation product of this gene shares limited sequence homology with cytochrome-c oxidase. An alternative embodiment is the polypeptide comprising the following amino acid sequence:

MLLKHLQRMVSPQVKASALKVVTLTANDKTSVSFSSLPGQGVIYNVIVWD

30 PFLNTSAAYIPAHTYACSFEAGEGSCASLGRVSSKVFFTLFALLGFFICFFGHR

FWKTELFFIGFIIMGFFFYILITRLTPIKYDVNLILTA VTGSGVGGMFLVA VVWR

FGILSICMLCVGLVLGFLISSVTFFTPLGNLKIFHDDGVFWVTFSCIAILIPVVF
MGCLRILNILTTCGVIGSYSVVLAIIDSYWSTLSYITLNVLKRALNKDFHRAFTN
VPFQTNDFIILAVWGMLAVSGITLQIRRERGRPFPPHPYKLWKQERERRVTNI
LDPSYHIPPLRERLYGRLTQIKGLFQKEQPAGERTPLLL (SEQ ID NO: 236).

- 5 In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

WARLRGPGAHARTSPQPWRGSPQAAMGFLQLLVVXVLXSEHRVAGAAE

- 10 VFGNSSEGLIEFSVGKFRYF

ELNRPFPPEEAILHDISSNVTFLIFQIHSQYQNTTVSFSPRRRSPTM (SEQ ID NO: 237). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in keratinocytes, brain, and spinal cord and to a lesser extent in hematopoietic cells and tissues.

- 15 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative disorders; hematopoietic disorders; integumentary disorders; immune dysfunction; learning disabilities. Similarly, polypeptides and
20 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and nervous systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., integumentary, neural, developmental,
25 cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 30 The tissue distribution in brain and spinal cord cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the

diagnosis and treatment of a variety of neurological and hematopoietic disorders. For example, elevated levels of expression of this gene product in brain and spinal cord indicates that it is involved in neurodegenerative disorders. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in

5 Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms,

10 hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

15 Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Alternately, expression of this gene product in hematopoietic cells indicates that it is involved in the proliferation, differentiation, survival, and activation of all hematopoietic lineages, including stem and progenitor cells. Expression of this gene

20 product in keratinocytes indicates that it is involved in normal skin function, and could be involved in skin disorders, dermatitis, and fibrosis. The protein is useful in detecting, treating, and/or preventing congenital disorders (i.e. nevi, moles, freckles, Mongolian spots, hemangiomas, port-wine syndrome), integumentary tumors (i.e. keratoses, Bowen's Disease, basal cell carcinoma, squamous cell carcinoma,

25 malignant melanoma, Paget's Disease, mycosis fungoides, and Kaposi's sarcoma), injuries and inflammation of the skin (i.e. wounds, rashes, prickly heat disorder, psoriasis, dermatitis), atherosclerosis, urticaria, eczema, photosensitivity, autoimmune disorders (i.e. lupus erythematosus, vitiligo, dermatomyositis, morphea, scleroderma, pemphigoid, and pemphigus), keloids, striae, erythema, petechiae, purpura, and

30 xanthelasma. In addition, such disorders may predispose increased susceptibility to viral and bacterial infections of the skin (i.e. cold sores, warts, chickenpox,

molluscum contagiosum, herpes zoster, boils, cellulitis, erysipelas, impetigo, tinea, athlete's foot, and ringworm). Moreover, the protein product of this gene may also be useful for the treatment or diagnosis of various connective tissue disorders (i.e., arthritis, trauma, tendonitis, chondromalacia and inflammation, etc.), autoimmune disorders (i.e., rheumatoid arthritis, lupus, scleroderma, dermatomyositis, etc.), dwarfism, spinal deformation, joint abnormalities, and chondrodysplasias (i.e. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid). Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:57 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2050 of SEQ ID NO:57, b is an integer of 15 to 2064, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:57, and where b is greater than or equal to a + 14.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 48

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

30 PRVRPASPPVRSPARWGSMAGSPLLWGPRAGGVLLVLLLLGLFRPPPALCA
RPVKEPRGLSAASPPLARLALLAASGGQCPEVRRRGRCRPGAGAGASAGAER

QERARAEAQRLRISRRASWRSCCASGAPPATLIRLWAWTTTPTRLQRSSLALC
SAPALTLPP (SEQ ID NO: 238). Polynucleotides encoding these polypeptides are
also provided.

5 This gene is expressed primarily in human pituitary and to a lesser extent in
pineal gland, and other areas of the brain.

Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, pituitary dysfunction; abnormal growth; neurological defects;
10 insufficient milk secretion; abnormal smooth muscle contraction. Similarly,
polypeptides and antibodies directed to these polypeptides are useful in providing
immunological probes for differential identification of the tissue(s) or cell type(s). For
a number of disorders of the above tissues or cells, particularly of the endocrine and
nervous systems, expression of this gene at significantly higher or lower levels is
15 routinely detected in certain tissues or cell types (e.g., endocrine, developmental,
reproductive, and cancerous and wounded tissues) or bodily fluids (e.g., lymph,
serum, plasma, urine, amniotic fluid, breast milk, synovial fluid and spinal fluid) or
another tissue or cell sample taken from an individual having such a disorder, relative
to the standard gene expression level, i.e., the expression level in healthy tissue or
20 bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic
epitopes shown in SEQ ID NO: 150 as residues: Pro-36 to Gly-42, Pro-64 to Ala-76,
Gly-83 to Ala-90, Ser-100 to Cys-108, Thr-126 to Ser-135. Polynucleotides encoding
said polypeptides are also provided.

25 The tissue distribution primarily in pituitary cells and tissues indicates that
polynucleotides and polypeptides corresponding to this gene are useful for the
diagnosis and/or treatment of a variety of disorders. Elevated expression of this gene
product in the pituitary indicates that it is possibly a hormone-like substance that
either controls pituitary development itself, or various processes controlled by the
30 pituitary. These include growth, milk secretion, smooth muscle contraction, diuresis,
blood pressure, and homeostasis. Thus, this gene product may have numerous clinical

applications. Expression of this gene product in other regions of the brain also indicates that it is involved in normal neurological function, and is useful in the treatment of a variety of neurological disorders. Representative uses are described in the "Biological Activity", "Hyperproliferative Disorders", and "Binding Activity" sections below, in Example 11, 17, 18, 19, 20 and 27, and elsewhere herein. Briefly, the protein can be used for the detection, treatment, and/or prevention of Addison's Disease, Cushing's Syndrome, and disorders and/or cancers of the pancreas (e.g. diabetes mellitus), adrenal cortex, ovaries, pituitary (e.g., hyper-, hypopituitarism), thyroid (e.g. hyper-, hypothyroidism), parathyroid (e.g. hyper-, hypoparathyroidism), hypothalamus, and testes. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:58 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1036 of SEQ ID NO:58, b is an integer of 15 to 1050, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:58, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 49

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

PRVRLATPNIWDL SMLFAFISLLV MLPTWWIVSSWL VWGVILFVYLVIRALRL

PZ030PCT

WRTAKLQVTLKKYSVHLEDMATNSRAFTNLVRKALRLIQETEVISRGFTLVS
AACPFNKAGQHPSQHLIGLRKAVYRTLRFQAAARLATLYMLKNYPLNSES
DNVTNYICVVPFKELGLGLSEEQISEEEAHNFTDGFSLPALKVLFQLWVAQSS
EFFRRLALLLSTANSPPGPLLTPALLPHRILSDVTQGLPHAHSACLEELKRSYE
5 FYRYFETQHQSVPOCLSQTQKSRELNNVHTAVRSLQLHLKALLNEVILEDE
LEKLVCTKETQELVSEAYPILEQKLKLIQPHVQASNNCWEEAISQVDKLLRRN
TDDKKGKPEIACENPHCTVSTFEAAYSTHCRQRSNPRGAGIRSLCR (SEQ ID
NO: 239). Polynucleotides encoding these polypeptides are also provided.

The polypeptide of this gene has been determined to have a transmembrane
10 domain at about amino acid position 7 - 23 of the amino acid sequence referenced in
Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 24 to
390 of this protein has also been determined. Based upon these characteristics, it is
believed that the protein product of this gene shares structural features to type Ib
membrane proteins.

15 The gene encoding the disclosed cDNA is believed to reside on chromosome
12. Accordingly, polynucleotides related to this invention are useful as a marker in
linkage analysis for chromosome 12.

This gene is expressed primarily in prostate and placenta and to a lesser extent
in pancreatic tumors and hematopoietic cells.

20 Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, prostate cancer; pancreatic cancer; prostate dysfunction; hematopoietic
disorders; reproductive diseases and/or disorders, and pancreatitis. Similarly,
25 polypeptides and antibodies directed to these polypeptides are useful in providing
immunological probes for differential identification of the tissue(s) or cell type(s). For
a number of disorders of the above tissues or cells, particularly of the endocrine and
immune systems, expression of this gene at significantly higher or lower levels is
routinely detected in certain tissues or cell types (e.g., reproductive, prostate,
30 pancreas, placental, vascular, and cancerous and wounded tissues) or bodily fluids
(e.g., lymph, serum, seminal fluid, plasma, urine, synovial fluid and spinal fluid) or

another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 151 as residues: Pro-85 to Ser-94, Pro-127 to Thr-136, Glu-154 to Glu-160, Phe-240 to Ser-250, Leu-255 to Leu-265, Leu-341 to Lys-351, Thr-372 to Gly-384. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in prostate and placental cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of a variety of reproductive disorders. Elevated expression of this gene product in the prostate indicates that it is involved in normal prostate function, and is a diagnostic marker for prostate cancer. Alternately, expression of this gene product in placenta indicates that it may play a role in normal vascular function, and is involved in such processes as angiogenesis and endothelial cell chemotaxis. Thus, this gene product is useful in the treatment of myocardial infarction, cancer, ischemia, and diabetic retinopathy. Expression of this gene product in placenta may also be indicative of fetal health and development.

Similarly, expression of this gene product in hematopoietic cells indicates that it is involved in the proliferation, differentiation, survival, or activation of all hematopoietic cell lineages. Finally, expression of this gene product in pancreatic cancers indicates that it may play a role in cancer in general, or in pancreatic function. The secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, and as nutritional supplements. It may also have a very wide range of biological activities. Representative uses are described in the "Chemotaxis" and "Binding Activity" sections below, in Examples 11, 12, 13, 14, 15, 16, 18, 19, and 20, and elsewhere herein. Briefly, the protein may possess the following activities: cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g. for treating human immunodeficiency virus infection, cancer, autoimmune diseases

and allergy); regulation of hematopoiesis (e.g. for treating anemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g. for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g. for treating
5 infections, tumors); hemostatic or thrombolytic activity (e.g. for treating hemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g. for treating septic shock, Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behavior. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. Protein, as well as,
10 antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:59 and may have been publicly available prior to conception of
15 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2519 of SEQ ID NO:59, b is an
20 integer of 15 to 2533, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:59, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 50

When tested against Jurkat and K562 cell lines, supernatants removed from
25 cells containing this gene activated the GAS (gamma activating sequence) and ISRE (interferon-sensitive responsive element) promoter elements, respectively. Thus, it is likely that this gene activates myeloid, leukemia, and to a lesser extent, other immune or hematopoietic cells and tissue cell-types, through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes
30 which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells.

Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. ISRE is also a promoter element found upstream in many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

AAPHPLLRLCLWCPLWPAWPLRGRPRSAWKRPPLVGPAGKLGCSMTTR
QPTAVSWPCWLMSSSLSTACLAWTLTGSLAREATRRARSLSPWNCARQV
PPSPPHSGLGRRGWAHCHLT CLLVTQLFRVGRIHPILSLPLVT (SEQ ID NO:
240). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in brain and placenta.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, vascular diseases; aberrant angiogenesis; neurological disorders; learning disorders; placental insufficiency; and fetal distress. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the vascular and neurological systems (CNS/PNS), expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, reproductive, vascular, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e.,

the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 152 as residues: Met-1 to Thr-7, Glu-36 to Ser-43,
5 Pro-46 to Gly-63. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in brain and placental cells and tissues, combined with the detected GAS and ISRE biological activities, indicates that the protein products of this gene are useful for the diagnosis and/or treatment of a variety of neural, reproductive, and vascular diseases and/or disorders. neurodegenerative disease
10 states, behavioral disorders, or inflammatory conditions. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis,
15 encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and
20 perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Expression of this gene product in placenta indicates that it may play a role
25 in blood vessel development or function, as the placenta is a highly vascularized organ. Thus, this gene product is involved in such processes as angiogenesis, endothelial cell chemotaxis, and vascular cord formation. Thus, it is useful in the treatment of such conditions as myocardial infarction; ischemia; and cancer. Alternately, expression of this gene product in the brain indicates that it may play a
30 role in the survival, proliferation, or function of neurons, and thus is useful in the diagnosis and treatment of such neurological disorders as ALS, schizophrenia, and

Alzheimer's Disease. It may likewise be involved in learning disorders as well.

Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement.

- 5 Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:60 and may have been publicly available prior to conception of
10 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 885 of SEQ ID NO:60, b is an
15 integer of 15 to 899, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:60, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 51

In another embodiment, polypeptides comprising the amino acid sequence of
20 the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

LQLASQSAGIKGMSHCARPTFTLLLASCFWAAAIPNRNVILSVSFRPLHMQ
FTLSILVFILRILILRSFL (SEQ ID NO: 241). Polynucleotides encoding these

- 25 polypeptides are also provided.

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 40 - 56 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 57 to 60 of this protein has also been determined. Based upon these characteristics, it is
30 believed that the protein product of this gene shares structural features to type Ia membrane proteins.

This gene is expressed primarily in spleen derived from patients with chronic lymphocytic leukemia.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, chronic lymphocytic leukemia; hematopoietic disorders; impaired immune function; cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in spleen cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of a variety of hematopoietic disorders. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the uses include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Elevated expression of this protein in the spleens of patients with CLL indicates that it is a useful marker for this Disease. Alternately, it is associated with the development and/or progression of the disease, and is a useful target for therapeutic intervention. Additionally, this gene

product may play more general roles in hematopoiesis, and may serve to control cellular decisions regarding proliferation, survival, activation, and/or differentiation of all hematopoietic cell lineages. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate
5 ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
10 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:61 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
15 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1065 of SEQ ID NO:61, b is an integer of 15 to 1079, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:61, and where b is greater than or equal to a + 14.

20 FEATURES OF PROTEIN ENCODED BY GENE NO: 52

The translation product of this gene shares sequence homology with a putative tyrosine protein kinase from the Chilo iridescent virus. See, for example, Genbank accession no. gi|2738451 (AF003534). Based on the sequence similarity, the translation product of this gene is expected to share at least some biological activities
25 with tyrosine kinase and signaling proteins. Such activities are known in the art, some of which are described elsewhere herein.

This gene is expressed in a variety of tissues, including microvascular endothelial cells, dendritic cells, and fetal tissues. as well as several tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as
30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, vascular, immune, and developmental diseases and/or disorders, particularly cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above
5 tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., vascular, immune, developmental, proliferative, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, amniotic fluid, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having
10 such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 154 as residues: Ala-21 to Lys-31, Arg-41 to Cys-56, Thr-92 to Cys-102, Arg-132 to Val-137, Lys-152 to Ile-159, Pro-199 to Ser-205, Arg-
15 210 to Asp-219, Ser-225 to Lys-230, Tyr-236 to Ala-241, Lys-243 to Leu-249, Thr-375 to Asp-381. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution and homology to a tyrosine kinase indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of cancer. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27,
20 and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes
25 suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such
30 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities,

such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Alternatively, the protein is useful in the detection, treatment, and/or prevention of vascular conditions, which include, but are not limited to, microvascular disease, vascular leak syndrome, aneurysm, stroke, atherosclerosis, arteriosclerosis, or embolism. For example, this gene product may represent a soluble factor produced by smooth muscle that regulates the innervation of organs or regulates the survival of neighboring neurons. Likewise, it is involved in controlling the digestive process, and such actions as peristalsis. Similarly, it is involved in controlling the vasculature in areas where smooth muscle surrounds the endothelium of blood vessels. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:62 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1914 of SEQ ID NO:62, b is an

integer of 15 to 1928, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:62, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 53

5 The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 2 - 18 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ib membrane proteins.

 This gene is expressed primarily in neutrophils.

10 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic diseases and/or disorders, particularly cancer and immune suppression. Similarly, polypeptides and antibodies directed to
15 these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids
20 (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 Preferred polypeptides of the present invention comprise immunogenic
25 epitopes shown in SEQ ID NO: 155 as residues: Gly-63 to Ser-72. Polynucleotides encoding said polypeptides are also provided.

 The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful as a marker for neutrophil monitoring in cancer and/or immune suppressed patients and/or during chemotherapy
30 or radiation therapy. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and

elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes
5 suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such
10 as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic
15 lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages,
20 and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as,
25 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:63 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
30 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or

more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 767 of SEQ ID NO:63, b is an integer of 15 to 781, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:63, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 54

This gene is expressed primarily in IL-1 and LPS induced neutrophils, and to a lesser extent, in fetal brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, hematopoietic, and neural diseases and/or disorders, particularly cancer and immune suppression. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, neural, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, amniotic fluid, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 156 as residues: Ile-28 to Trp-37, Ser-68 to Lys-81. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful as a marker in neutrophils to monitor patients who are immune suppressed or cancer patients during chemotherapy or radiation therapy. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in

regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune
5 responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory
10 bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's
15 Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Alternatively,
20 polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in
25 addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
30 related to SEQ ID NO:64 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1180 of SEQ ID NO:64, b is an integer of 15 to 1194, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:64, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 55

This gene is expressed primarily in prostate.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, urogenital diseases and/or disorders, particularly prostate cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the urogenital system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., urogenital, prostate, renal, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 157 as residues: Arg-30 to Gln-36. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in prostate cancer cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for study, treatment and diagnosis of prostate cancer and other urogenital disorders. Moreover, the expression within cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis,

treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation
5 and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of
10 potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types
15 of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new
20 insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or
25 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:65 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
30 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or

more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1663 of SEQ ID NO:65, b is an integer of 15 to 1677, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:65, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 56

A preferred polypeptide of the invention comprises the following amino acid sequence:

MVLVLRHPLCARERAFREPGRGLLTRTGQHDGAPAVTAVPGPLGAVAAAEG
10 RRSAGWAGGSSPPRKVLWGDMRGRRAGVDVLGPALSSEAAGAEARGWGM
PGMGVGVGASETREGALFLGREGVHGPCPMDGLGPWPWGPW (SEQ ID NO:
242). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in rejected kidney.

Therefore, polynucleotides and polypeptides of the invention are useful as
15 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, diseases and/or disorders affecting the kidney. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of
20 disorders of the above tissues or cells, particularly of the urinary tract, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., urogenital, renal, kidney, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative
25 to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 158 as residues: Ala-30 to Gly-36, Asp-45 to Trp-50, Lys-65 to Cys-71, Pro-80 to Cys-87. Polynucleotides encoding said polypeptides are
30 also provided.

The tissue distribution in kidney indicates the protein product of this gene could be used in the treatment and/or detection of kidney diseases including renal failure, nephritis, renal tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney stones, in addition to Wilm's Tumor Disease, and congenital kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. The protein is useful for modulating the immune response to aberrant proteins, as may exist in proliferating cells and tissues. Such modulation of the immune response would also show utility in inhibiting the rejection of transplanted tissues, particularly of the renal system. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:66 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1223 of SEQ ID NO:66, b is an integer of 15 to 1237, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:66, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 57

The translation product of this gene shares sequence homology with both human and mouse Fibulin-2 which is an extracellular matrix protein found in heart tissue (See Genbank Accession Nos. emb|CAA57876.1 and emb|CAA53040.1, respectively; all references available through these accessions are hereby incorporated

herein by reference; for example, J. Cell Biol. 123 (5), 1269-1277 (1993)). Preferred polypeptides encoded by this gene comprise the following amino acid sequence:

MGPAVKMWTNAWKGLDDCHYNQLCENTPGGHRCSGPCRGYRMQGPSLPCL
DVNECLQLPKACAYQCHNLQGSYRCLCPPGQTLLRDGKACTSLERNQGNVT
5 TVSHRGPLLWLRPWASIPGTSYHAWVSLRPGPMALSSVGRAWCPPGFIRQN
GVCTDLDECRVRNLCQHACRNTEGSYQCLCPAGYRLLPSGKNCQDINECEEE
SIECGPGQMCFNTRGSYQCVDTPCPATYRQGSPSGTCFRRCSDCGTGGPSTL
QYRLLPLPLGVRAHHDVARLTAFASEGV PANRTELSMLEPDRSPFALRPLRA
GLGAVYTRRALTRAGLYRLTVRAAAPHQS VFVLLIAVSPYPY (SEQ ID NO:

10 243). Polynucleotides encoding these polypeptides are also provided.

A preferred polypeptide fragment of the invention comprises the following amino acid sequence:

MRVLVVTIPIYWALARESGEALNGHSLTGGKFRQSHTWSLLQGAHDDPV
ARGLDPDGLLLLDVVVNGVVPRAWLTQIFKCRTLKKHYVQTRAWPAVRG

15 LHTALLPGRPPLVPTLQPQHPVQRGPGPPAPAGAAPAGLSYQLGL (SEQ ID
NO: 244). Polynucleotides encoding these polypeptides are also provided.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the

20 following amino acid sequence:

HASGAFLVVRGEPQGSWGSMTGVINGRKFGVATLNTSVMQEAHSGVSSIHSS
IRHVPANVGPLMRVLVVTIPIYWALARESGEALNGHSLTGGKFRQESHVEF
ATGELLTMTQWPGVWIPMASCSSTWWSMALSPDSLADADLQVQDFEEHYV
QTGPGQLFVGSTQRFFQGGLPSFLRCNHSIQYNAARGPQQLVQHLRASAISS
25 AFDPEAEALRFQLATALQAEENEVGCPEGFELDSQGAFCVDVDECAWDAHL
CREGQRCVNLLGSYRCLPDCGPGFRVADGAGCEDVDECLEGLDDCHYNQLC
ENTPGGHRCSGPCRGYRMQGPSLPCLDVNECLQLPKACAYQCHNLQGSYRCL
CPPGQTLLRDGKACTSLERNQGNVTTVSHRGPLLWLRPWASIPGTSYHAWV
SLRPGPMALSSVGRAWCPPGFIRQNGVCTDLDECRVRNLCQHACRNTEGSY
30 QCLCPAGYRLLPSGKNCQDINECEEEESIECGPGQMCFNTRGSYQCVDTPCPAT
YRQGSPSGTCFRRCSDCGTGGPSTLQYRLLPLPLGVRAHHDVARLTAFASEV

GVPANRTELSMLEPDPRSPFALRPLRAGLGAVYTRRALTRAGLYRLTVRAAA
PRHQSVMFVLLIAVSPYPY (SEQ ID NO: 245). Polynucleotides encoding these
polypeptides are also provided.

When tested against U937 and Jurkat cell lines, supernatants removed from
5 cells containing this gene repeatedly activated the GAS (gamma activating sequence)
promoter element. Thus, it is likely that this gene activates myeloid, T-cells, and to a
lesser extent, other immune and hematopoietic cells and tissue cell types, through the
JAK-STAT signal transduction pathway. GAS is a promoter element found upstream
of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway
10 is a large, signal transduction pathway involved in the differentiation and proliferation
of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of
the GAS element, can be used to indicate proteins involved in the proliferation and
differentiation of cells.

This gene is expressed primarily in kidney.

15 Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, diseases and/or disorders affecting the kidney and renal system.
Similarly, polypeptides and antibodies directed to these polypeptides are useful in
20 providing immunological probes for differential identification of the tissue(s) or cell
type(s). For a number of disorders of the above tissues or cells, particularly of the
urinary tract, expression of this gene at significantly higher or lower levels is
routinely detected in certain tissues or cell types (e.g., renal, urogenital, kidney, and
cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine,
25 synovial fluid and spinal fluid) or another tissue or cell sample taken from an
individual having such a disorder, relative to the standard gene expression level, i.e.,
the expression level in healthy tissue or bodily fluid from an individual not having the
disorder.

Preferred polypeptides of the present invention comprise immunogenic
30 epitopes shown in SEQ ID NO: 159 as residues: Lys-32 to Ser-37, His-89 to Gly-94,
Asn-124 to Gln-130, Ala-163 to Val-168, Cys-196 to Arg-201, Gln-244 to Gln-264,

His-288 to Tyr-294, Leu-314 to Gln-319, Ala-392 to Ser-399, Pro-412 to Asp-419, Ala-452 to Pro-460, Arg-466 to Thr-473. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in rejected kidney, the homology to the conserved
5 Fibulin-2 protein, in addition to the detected GAS biological activity, indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting kidneys, particularly proliferative disorders. Representative uses are described here and elsewhere herein. The protein product of this gene could be used in the treatment and/or detection of kidney diseases
10 including renal failure, nephritis, renal tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney stones, in addition to Wilm's Tumor Disease, and congenital kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. Furthermore, the protein may also be used
15 to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

20 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:67 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is
25 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1920 of SEQ ID NO:67, b is an integer of 15 to 1934, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:67, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 58

Preferred polypeptides of the invention comprise the following amino acid sequence:

MGEKFLLLAMKENHPECFCCKILKILHCMDPGEWLPQTEHCVHLPKEFLIWT
MDIASNERSEIQSVALRLASKVISHHMQTCVENRELIAAELKQWVQLVILSC
5 DHLPTESRLAVVEVLTSTTPLFLTNPHPILELQDTLALWKCVLTLQSEEQAV
RDAATETVTTAMSQENTCQSTEFACQVDASIALALALAVLCDLLQQWDQL
APGLPILLGWLLGESDDLACVESMHQVEEDYLFEKAEVNFWAETLIFVKYL
CKHLFCLLSKSGWRPPSPEMLCHLQRMVSEQCHLLSQFFRELPPAAEFVKTV
EFTRLRIQEERTLACLRLAFLEGKEGEDTLVLSVWDSYAESRQLTLPRTEAA
10 C (SEQ ID NO: 246). Polynucleotides encoding such polypeptides are also provided.

A preferred polypeptide fragment of the invention comprises the following amino acid sequence: MGEPNRHPSM

FLLLLVLERLYASPM DGTSSALSMGPFVFPFIMRCGHSPVYHSREMAARALVP
FVMIDHIPNTIRTLLSTL
15 PSCTDQCFRAKPHSWGHSRFFHLLQAYSDSKTRNEFRLPARAD (SEQ ID NO:
247). Polynucleotides encoding these polypeptides are also provided.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the
20 following amino acid sequence:

MTGREFFSRFPELYPFLLKQLETVANTVDSDMGEPNRHPSMFLLLLVLERLY
ASPM DGTSSALSMGPFVFPFIMRCGHSPVYHSREMAARALVPFVMIDHIPNTIR
TLLSTLPSCTDQCFRQNHIGTLLQVFHLLQAYSDSKHGTNSDFQHELT DITV
CTKAKLWLAKRQNPCLVTRAVYIDILFLLTCCLNRS AKDNQPVLES LGFWEE
25 VRGIISGSELITGFPWAFKVPGLPQYLQSLTRLAIAAVWAAA AKSGERETNPVI
SFSQLLESAPPEVRS LTLEALLEKFLAAASGLGEKGVPLL CNMGEKFLLAM
KENHPECFCCKILKILHCMDPGEWLPQTEHCVHLPKEFLIWTMDIASNERSEIQ
SVALRLASKVISHHMQTCVENRELIAAELKQWVQLVILSCEDHLPTESRLAVV
EVL TSTTPLFLTNPHPILELQDTLALWKCVLTLQSEEQAVRDAATETVTTAM
30 SQENTCQSTEFACQVDASIALALALAVLCDLLQQWDQLAPGLPILLGWLLG
ESDDLACVESMHQVEEDYLFEKAEVNFWAETLIFVKYLCKHLFCLLSKSG

WRPPSPEMLCHLQRMVSEQCHLLSQFFRELPPAAEFVKTVEFTRLRIQEERTL
 ACLRLLAFLEGKEGEDTLVLSVWDSYAESRQLTLPRTEAAC (SEQ ID NO:
 248). Polynucleotides encoding these polypeptides are also provided.

The polypeptide of this gene has been determined to have two transmembrane
 5 domains at about amino acid position 144 - 160, and 462 - 478 of the amino acid
 sequence referenced in Table 1 for this gene. Based upon these characteristics, it is
 believed that the protein product of this gene shares structural features to type IIIa
 membrane proteins. Included in this invention as a preferred domain is the formate
 and nitrite transporters domain, which was identified using the ProSite analysis tool
 10 (Swiss Institute of Bioinformatics). A number of bacterial and archaeobacterial
 proteins involved in transporting formate or nitrite have been shown [1] to be related:
 - focA and focB, from *Escherichia coli*, transporters involved in the bidirectional
 transport of formate. - fdhC, from *Methanobacterium formicicum* and
thermoformicicum, a probable formate transporter. - nirC, from *Escherichia coli* and
 15 *Salmonella typhimurium*, a probable nitrite transporter. - *Bacillus subtilis*
 hypothetical protein yrhG. - *Bacillus subtilis* hypothetical protein ywcJ (ipa-48R).
 These transporters are proteins of about 280 residues and seem to contain six
 transmembrane regions. As signature patterns, we selected two conserved regions.
 The first one is located in what seems to be a cytoplasmic loop between the second
 20 and third transmembrane domains; the second is part of the fourth transmembrane
 region. The 70 Kd yeast hypothetical protein YHL008c is highly similar, in its N-
 terminal section, to the prokaryotic members of this family. The consensus pattern is
 as follows: [LIVMA]-[LIVMY]-x-G-[GSTA]-[DES]-L-[FI]-[TN]-[GS].

Preferred polypeptides of the invention comprise the following amino acid
 25 sequence: IISGSELITG (SEQ ID NO: 249). Polynucleotides encoding these
 polypeptides are also provided. Further preferred are polypeptides comprising the
 formate and nitrite transporter domain of the sequence referenced in Table for this
 gene, and at least 5, 10, 15, 20, 25, 30, 50, or 75 additional contiguous amino acid
 residues of this referenced sequence. The additional contiguous amino acid residues is
 30 N-terminal or C- terminal to the formate and nitrite transporter domain. Alternatively,
 the additional contiguous amino acid residues is both N-terminal and C-terminal to

the formate and nitrite transporter domain, wherein the total N- and C-terminal contiguous amino acid residues equal the specified number. The above preferred polypeptide domain is characteristic of a signature specific to formate and nitrite transporter proteins. Based on the sequence similarity, the translation product of this gene is expected to share at least some biological activities with formate and nitrite transporter proteins. Such activities are known in the art, some of which are described elsewhere herein. It is believed that this gene maps to chromosome 2. Accordingly, polynucleotides derived from this gene are useful in linkage analysis as markers for chromosome 2.

10 This gene is expressed primarily in cells of the immune system, primarily T-cells and to a lesser extent in spleen, liver, thymus, tonsils, and testis.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic diseases and/or disorders, particularly disorders affecting hematopoiesis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of hematopoietic cells, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 160 as residues: Gly-2 to Pro-8, Ser-82 to His-92, Tyr-107 to Asp-117, Arg-162 to Pro-169, Ser-224 to Thr-229, Leu-310 to His-315, Ser-333 to Glu-338, Glu-381 to Ser-388, Gln-428 to Ala-433, Met-446 to Thr-455, Ser-548 to Ser-554, Gly-613 to Asp-618, Ser-627 to Gln-633. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in immune cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting hematopoiesis, including cancers. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are

related to SEQ ID NO:68 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
 5 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3286 of SEQ ID NO:68, b is an integer of 15 to 3300, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:68, and where b is greater than or equal to a + 14.

10 **FEATURES OF PROTEIN ENCODED BY GENE NO: 59**

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

15 VDGIDKLDIEFLQQFLETHSRGPR LHSPGHASQEATPGANMSSGTELLWPGAA
 LLVLLGVAASLCVRCSRPGAKRSEKIYQQRSLREDQQSFTGSRTYSLVGQAW
 PGPLADMAPTRKDKLLQFYPSLEDPASSRYQNFSKGSRHGSEEA YIDPIAMEY
 YNWGRFSKPPEDDDANSYENVLICKQKTTTETGAQQEGIGGLCRGDL SLSLAL
 20 KTGPTSGLCPSASPEEDEGI (SEQ ID NO: 250). Polynucleotides encoding these polypeptides are also provided.

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 10 - 26 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ib membrane proteins.

25 The gene encoding the disclosed cDNA is believed to reside on chromosome 7. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 7.

This gene is expressed primarily in bone marrow, CD34 positive cells, and immune cells, including, neutrophils, T-cells, B-cells, macrophages, monocytes, and
 30 dendritic cells and to a lesser extent in brain and tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders affecting the immune and hematopoietic systems, particularly hematopoiesis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the the immune system and hematopoietic system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 161 as residues: Ser-29 to Thr-57, Pro-74 to Lys-79, Pro-85 to Glu-107, Tyr-118 to Tyr-136, Gln-144 to Gln-152, Ala-182 to Glu-188. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in immune and hematopoietic cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting the immune system and hematopoiesis. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for

immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, and as nutritional supplements. It may also have a very wide range of biological activities.

Representative uses are described in the "Chemotaxis" and "Binding Activity" sections below, in Examples 11, 12, 13, 14, 15, 16, 18, 19, and 20, and elsewhere herein. Briefly, the protein may possess the following activities: cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g. for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g. for treating anemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g. for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g. for treating infections, tumors);

hemostatic or thrombolytic activity (e.g. for treating hemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g. for treating septic shock, Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behavior. Also contemplated is the use of the
5 corresponding nucleic acid in gene therapy procedures.

Based upon the the proteins immune cell specific message distribution, it may be involved in many aspects of the immune response, especially its initial stages, inflammation, allograft rejection, infectious disease response etc. The expression of this clone is
10 frequently found in the hematopoietic cell cDNA libraries. Thus, this factor could be involved in the control of hematopoietic cell proliferation, differentiation, and function. Based on this one can postulate its use in the management of anemias, leukemias, neutropenia, thrombocytopenia, autoimmune diseases, blood tissue engraftment, and poikilothromerythromatosis. Furthermore, the
15 protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

20 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:69 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is
25 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1783 of SEQ ID NO:69, b is an integer of 15 to 1797, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:69, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 60

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

5 VLWREASALVLSNRLSSGLLDLLLQPAIHSRLFPRRSRGLSEGE GSSVSLQRS
RVLSAMKHVLNLYLLGVVLTLLSIFVRVMESLEGLLESPPGTSWTTSQLAN
TEPTKGLPDHPSRSM (SEQ ID NO: 251). Polynucleotides encoding these
polypeptides are also provided.

This gene is expressed primarily in immune cells including activated T cells,
10 macrophages, jurkat cells, bone marrow cells, and osteoblasts and to a lesser extent in
kidney cortex, brain, placenta and lung.

Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
15 not limited to, immune and hematopoietic diseases and/or disorders, particularly
inflammation and diseases related to inflammatory activity. Similarly, polypeptides
and antibodies directed to these polypeptides are useful in providing immunological
probes for differential identification of the tissue(s) or cell type(s). For a number of
disorders of the above tissues or cells, particularly of the immune system, expression
20 of this gene at significantly higher or lower levels is routinely detected in certain
tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded
tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal
fluid) or another tissue or cell sample taken from an individual having such a
disorder, relative to the standard gene expression level, i.e., the expression level in
25 healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic
epitopes shown in SEQ ID NO: 162 as residues: Pro-34 to Met-63. Polynucleotides
encoding said polypeptides are also provided.

The tissue distribution in immune cells and tissues indicates that
30 polynucleotides and polypeptides corresponding to this gene are useful for treating or
diagnosing disease related to the normal or abnormal activation of T cells.

Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell
5 lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for
10 immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity
15 disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful
20 in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as,
25 antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:70 and may have been publicly available prior to conception of
30 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1359 of SEQ ID NO:70, b is an integer of 15 to 1373, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:70, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 61

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

YTFHTQIFLDFPMIFLTVLPLAFLFLHSGFYHYISFSLFSLALFFFLDVATFR
RPGQLFCERSVLFDMFHFGFVSLFLHEWQAKHFWAGLF
IVLPSDVFFSVHHLEAPDGSPNIAKLSLIILLR (SEQ ID NO: 252).

Polynucleotides encoding these polypeptides are also provided.

The polypeptide of this gene has been determined to have two transmembrane domains at about amino acid position 2 - 18 and 22 - 38 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type IIIa membrane proteins.

This gene is expressed in many tissues including brain, liver, prostate, testes, cartilage, gall bladder. Expression is also seen in a number of tumors including colon carcinoma, pancreas tumor, osteoclastoma, ovarian cancer, B cell lymphoma and acute lymphocytic leukemias.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, tumors of various organs including the pancreas, colon, and bone. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the

major organs, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, hepatic, metabolic, reproductive, testicular, skeletal, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or
5 another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in tumors and proliferative tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating or
10 diagnosing tumors of several major organs including the pancreas and large intestine. This protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below
15 and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain
20 neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating,
25 detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in
30 proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the

protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or
5 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:71 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
10 excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1565 of SEQ ID NO:71, b is an integer of 15 to 1579, where both a and b correspond to the positions of nucleotide
15 residues shown in SEQ ID NO:71, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 62

This gene is expressed primarily in dendritic cells and fetal liver/spleen and to a lesser extent in many tissues including tonsils, fetal lung, stromal cell lines, bone
20 marrow cell lines, placenta and tumors including hepatocellular carcinoma, pancreas tumor and osteosarcoma.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
25 not limited to, diseases and/or disorders of the immune and hematopoietic system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels is
30 routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine,

synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 The tissue distribution in dendritic cells and fetal liver/spleen indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosing and treating disorders of the immune system particularly related to the control and generation of precursor cells. polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic
10 related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the uses include bone marrow cell ex-vivo culture, bone marrow
15 transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

 The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of
20 stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies
25 directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:72 and may have been publicly available prior to conception of
30 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1014 of SEQ ID NO:72, b is an integer of 15 to 1028, where both a and b correspond to the positions of nucleotide
5 residues shown in SEQ ID NO:72, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 63

This gene is expressed primarily in adrenal gland tumor and endothelial cells.

Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, endocrine and vascular diseases and/or disorders, particularly diseases associated with the vascular endothelium. Similarly, polypeptides and antibodies
15 directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the vascular system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell
20 types (e.g., endocrine, vascular, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in endothelial cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosing and treating
25 disorders that involve the vascular system including diseases such as atherosclerosis, neoangiogenesis associated with tumor growth and conditions associated with inflammation. Moreover, the protein is useful in the detection, treatment, and/or prevention of a variety of vascular disorders and conditions, which include, but are not limited to microvascular disease, vascular leak syndrome, aneurysm, stroke,
30 embolism, thrombosis, coronary artery disease, arteriosclerosis, and/or atherosclerosis. Alternatively, the protein is useful in the treatment, detection, and/or

prevention of metabolic disorders, particularly lethargy and depression. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:73 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3660 of SEQ ID NO:73, b is an integer of 15 to 3674, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:73, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 64

The translation product of this gene is related to bovine PAM precursor. See Genbank record gi|163482 incorporated herein by reference. Moreover, see following patent publications are also incorporated herein by reference: J04311386 and WO8902460. Many bioactive peptides terminate with an amino acid alpha-amide at their COOH terminus. The enzyme responsible for this essential posttranslational modification is known as peptidyl-glycine alpha-amidating monooxygenase or PAM. An NH₂-terminal signal sequence and short propeptide precede the NH₂ terminus of purified PAM. The sequences of several PAM cyanogen bromide peptides were localized in the NH₂-terminal half of the predicted protein. The forms of PAM purified from bovine neurointermediate pituitary is generated by endoproteolytic cleavage at a subset of the 10 pairs of basic amino acids in the precursor. High levels of PAM mRNA have been found in bovine pituitary and cerebral cortex. In

corticotropic tumor cells, levels of PAM mRNA and pro-ACTH/endorphin mRNA are known to be regulated in parallel by glucocorticoids and CRF.

This gene is expressed primarily in endometrial tumors, dendritic cells, a multiple sclerosis library, kidney, hematopoietic cells, melanocytes, osteoblasts, the spleen, colon, ovary, stromal cells, fetal and adult brain, heart, and in tissues undergoing wound repair.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, endometriosis, endometrial cancer, multiple sclerosis, hematopoietic diseases, bone disease, and wound healing. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly the hematopoietic system and female reproduction, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., reproductive, immune, hematopoietic integumentary, skeletal, gastrointestinal, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in dendritic and hematopoietic cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful as a therapeutic or diagnostic agent in Diseases of hematopoietic origin as well as the female reproductive track due to the gene's primary pattern of expression. polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the uses

include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency
5 etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. The protein may also have a very wide range of biological activities. Representative uses are described in the "Chemotaxis" and "Binding Activity" sections below, in Examples 11, 12, 13, 14, 15,
10 16, 18, 19, and 20, and elsewhere herein. Briefly, the protein may possess the following activities: cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g. for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g. for treating anemia or as adjunct to
15 chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g. for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g. for treating infections, tumors); hemostatic or thrombolytic activity (e.g. for treating hemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g. for treating septic shock,
20 Crohn's Disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behavior. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

25 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:74 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is
30 cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general

formula of a-b, where a is any integer between 1 to 2783 of SEQ ID NO:74, b is an integer of 15 to 2797, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:74, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 65

The translation product of this gene shares sequence similarity with several G-protein coupled receptors (See Genbank Accession No. gb|AAC77910.1| (AF061443); all references available through this accession are hereby incorporated herein by reference; for example, Mol. Endocrinol. 12, 1830-1845 (1998)). G-protein coupled receptors are well known in the art and affect a variety of functions. In particular, the translation product of this gene shares similarity with Follicular Stimulating Hormone Receptor.

Preferred polypeptides encoded by this gene comprise the following amino acid sequence:

15 GTRFPTGETPSLGFTVTLVLLNSLAFLLMAYIYTKLYCNLEKEDLSENSQSSMI
KHVAWLIFTNCIFFCPVAFFSFAPLITAISISPEIMKSVTLIFFP (SEQ ID NO:
253). Polynucleotides encoding such polypeptides are also provided.

A preferred polypeptide fragment of the invention comprises the following amino acid sequence: MIKHVAWLIFTNCIFFCP


20 VAFFSFAPLITAISISPEIMKSVTLIFFPCLLA (SEQ ID NO: 254). Polynucleotides encoding these polypeptides are also provided.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the

25 following amino acid sequence:

GTRFPTGETPSLGFTVTLVLLNSLAFLLMAYIYTKLYCNLEKEDLSENSQSSMI
KHVAWLIFTNCIFFCPVAFFSFAPLITAISISPEIMKSVTLIFFPLPACLNPVLYVF
FNPKFKEWDKLLKRRVTKKSGSVSVSISSQGGCLEQDFYYDCGMYSHLQGN
LTVCDCCESFLLTKPVSCKHLIKSHSCPALAVASCQRPEGYWSDCGTQSAHS
30 DYADEEDSFVSDSSDQVQACGRACFYQSRGFPLVRYAYNLPRVKD (SEQ ID
NO: 255). Polynucleotides encoding these polypeptides are also provided.

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 43 - 59 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 60 to 207 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia membrane proteins. Included in this invention as preferred domains are Zinc finger, C2H2 type domains, which were identified using the ProSite analysis tool (Swiss Institute of Bioinformatics). 'Zinc finger' domains [1-5] are nucleic acid-binding protein structures first identified in the *Xenopus* transcription factor TFIIA. These domains have since been found in numerous nucleic acid-binding proteins. A zinc finger domain is composed of 25 to 30 amino-acid residues. There are two cysteine or histidine residues at both extremities of the domain, which are involved in the tetrahedral coordination of a zinc atom. It has been proposed that such a domain interacts with about five nucleotides. A schematic representation of a zinc finger domain is shown below:



Many classes of zinc fingers are characterized according to the number and positions of the histidine and cysteine residues involved in the zinc atom

coordination. In the first class to be characterized, called C2H2, the first pair of zinc coordinating residues are cysteines, while the second pair are histidines. A number of experimental reports have demonstrated the zinc- dependent DNA or RNA binding property of some members of this class. Some of the proteins known to include

5 C2H2-type zinc fingers are listed below. We have indicated, between brackets, the number of zinc finger regions found in each of these proteins; a '+' symbol indicates that only partial sequence data is available and that additional finger domains is present. In addition to the conserved zinc ligand residues it has been shown that a number of other positions are also important for the structural integrity of the C2H2

10 zinc fingers. The best conserved position is found four residues after the second cysteine; it is generally an aromatic or aliphatic residue. The consensus pattern is as follows: C-x(2,4)-C-x(3)-[LIVMFYWC]-x(8)-H-x(3,5)-H.

Preferred polypeptides of the invention comprise the following amino acid sequence: CDCCESFLLTKPVSCCKHLIKSH (SEQ ID NO: 256). Polynucleotides

15 encoding these polypeptides are also provided. Further preferred are polypeptides comprising the Zinc finger, C2H2 type domain of the sequence referenced in Table for this gene, and at least 5, 10, 15, 20, 25, 30, 50, or 75 additional contiguous amino acid residues of this referenced sequence. The additional contiguous amino acid residues is N-terminal or C- terminal to the Zinc finger, C2H2 type domain.

20 Alternatively, the additional contiguous amino acid residues is both N-terminal and C-terminal to the Zinc finger, C2H2 type domain, wherein the total N- and C-terminal contiguous amino acid residues equal the specified number. The above preferred polypeptide domain is characteristic of a signature specific to zinc finger proteins. Based on the sequence similarity, the translation product of this gene is expected to

25 share at least some biological activities with G-coupled proteins, their receptors, and zinc finger proteins. Such activities are known in the art, some of which are described elsewhere herein.

This gene is expressed primarily in adult and fetal liver, human placenta, colon carcinoma cell lines and fibroblasts and to a lesser extent in the fetal and adult

30 brain, the developing nervous system, lung, pancreas, salivary gland, breast tissue, and dendritic cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, diseases of the liver, developmental abnormalities, neurologic diseases, lung cancer, pancreatic cancer, and colon cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neurological and hepatic origin, as well as the proliferation and/or differentiation of numerous types of tissues, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., hepatic, immune, hematopoietic, neural, gastrointestinal, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 167 as residues: Pro-62 to Asp-67, Arg-74 to Gly-80, Gln-146 to Glu-168. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in fetal liver indicates that polynucleotides and polypeptides corresponding to this gene are useful for a diagnostic marker or therapeutic in a wide variety of disease states. polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the uses include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the

5 differentiation and/or proliferation of various cell types. Alternatively, the protein expression in placental and brain tissue indicates the protein is useful in the detection, treatment, and/or prevention of vascular conditions, which include, but are not limited to, microvascular disease, vascular leak syndrome, aneurysm, stroke, atherosclerosis, arteriosclerosis, or embolism. For example, this gene product may represent a soluble

10 factor produced by smooth muscle that regulates the innervation of organs or regulates the survival of neighboring neurons. Likewise, it is involved in controlling the digestive process, and such actions as peristalsis. Similarly, it is involved in controlling the vasculature in areas where smooth muscle surrounds the endothelium of blood vessels. The protein is useful in the treatment, detection, and/or prevention of

15 bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's Disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic

20 depression, delirium, severe mental retardation and dyskinesias, such as Huntington's Disease or Gilles de la Tourette's syndrome. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed

25 against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:75 and may have been publicly available prior to conception of

30 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2689 of SEQ ID NO:75, b is an integer of 15 to 2703, where both a and b correspond to the positions of nucleotide
5 residues shown in SEQ ID NO:75, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 66

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by
10 the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:
ALENSGSPGLQDSARAHFNXSLRSFSFLRNQMYIFELSLYLEGTSFVVLLFLL
ISVSLDSPPTTKGWDSVLHIWVPLIVQ (SEQ ID NO: 257). Polynucleotides
encoding these polypeptides are also provided.

15 This gene is expressed primarily in placenta and in hematopoietic cells, especially those of T-cell and monocyte origin and to a lesser extent in the brain, endothelial cells, and the lungs.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
20 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hematopoietic, vascular, and developmental diseases and/or disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the
25 immune system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., vascular, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene
30 expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 168 as residues: Ser-30 to Trp-37. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in hematopoietic cells indicates that polynucleotides
5 and polypeptides corresponding to this gene are useful for therapeutic and/or
diagnostic intervention in hematopoietic and developmental disorders. Representative
uses are described in the "Immune Activity" and "infectious disease" sections below,
in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the uses
include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow
10 reconstitution, radiotherapy or chemotherapy of neoplasia.

The gene product may also be involved in lymphopoiesis, therefore, it can be
used in immune disorders such as infection, inflammation, allergy, immunodeficiency
etc. In addition, this gene product may have commercial utility in the expansion of
stem cells and committed progenitors of various blood lineages, and in the
15 differentiation and/or proliferation of various cell types. Alternatively, the protein is
useful in the detection, treatment, and/or prevention of vascular conditions, which
include, but are not limited to, microvascular disease, vascular leak syndrome,
aneurysm, stroke, atherosclerosis, arteriosclerosis, or embolism. For example, this
gene product may represent a soluble factor produced by smooth muscle that
20 regulates the innervation of organs or regulates the survival of neighboring neurons.
Likewise, it is involved in controlling the digestive process, and such actions as
peristalsis. Similarly, it is involved in controlling the vasculature in areas where
smooth muscle surrounds the endothelium of blood vessels. Furthermore, the protein
may also be used to determine biological activity, to raise antibodies, as tissue
25 markers, to isolate cognate ligands or receptors, to identify agents that modulate their
interactions, in addition to its use as a nutritional supplement. Protein, as well as,
antibodies directed against the protein may show utility as a tumor marker and/or
immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
30 available and accessible through sequence databases. Some of these sequences are
related to SEQ ID NO:76 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general
5 formula of a-b, where a is any integer between 1 to 728 of SEQ ID NO:76, b is an integer of 15 to 742, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:76, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 67

10 This gene is expressed primarily in the prostate and to a lesser extent in in human B-cell lymphomas.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
15 not limited to, prostate cancer and diseases of hematopoietic origin, particularly of B-cells. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the prostate and immune systems, expression of this gene at significantly higher or lower
20 levels is routinely detected in certain tissues or cell types (e.g., prostate, reproductive, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, seminal fluid, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the
25 fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 169 as residues: Asp-33 to Lys-42. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in prostate tissue indicates that polynucleotides and
30 polypeptides corresponding to this gene are useful as a therapeutic or diagnostic marker for prostate cancer and disorders involving hematopoietic cells, especially

those of B-cell origin. Moreover, the expression within cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment, and/or prevention of developmental diseases and disorders, including cancer, and other proliferative
5 conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation.

Dysregulation of apoptosis can result in inappropriate suppression of cell
10 death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may
15 also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and is useful in the detection, treatment, and/or prevention of
20 degenerative or proliferative conditions and diseases. The protein is useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. The protein is useful in modulating the immune response to aberrant proteins and polypeptides, as may exist
25 in rapidly proliferating cells and tissues. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the
30 above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:77 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1811 of SEQ ID NO:77, b is an integer of 15 to 1825, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:77, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 68

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:
GHESICGSCRSWIYFSIRCRRRMRPWWSLLLEACATCAQTGPTRSTSCTQEVS
HSSSTAYPAPMRRRCCL PSPRSCT (SEQ ID NO: 258). Polynucleotides encoding these polypeptides are also provided.

The gene encoding the disclosed cDNA is believed to reside on chromosome 17. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 17.

This gene is expressed primarily in the brain and the developing embryo and to a lesser extent in the heart, colon, adipose tissue, kidney, mammary tissue, activated T-cells and dendritic cells.

Therefore, polynucleotides and polypeptides of the invention are useful as
5 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological diseases, developmental conditions, colon cancer, and hematopoietic diseases, especially of T-cell origin. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological
10 probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., neural, developmental, cardiovascular, adipose, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g.,
15 lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic
20 epitopes shown in SEQ ID NO: 170 as residues: Thr-18 to Cys-26, Glu-29 to Thr-36, Ser-50 to Thr-55. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution in brain, combined with the detected GAS biological activity, indicates that polynucleotides and polypeptides corresponding to this gene are useful for therapeutic and/or diagnostic agents in neurological diseases,
25 developmental abnormalities, colon cancer, and hematopoietic diseases, especially those of T-cell origin. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease,
30 Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal

cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated
5 expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Furthermore, the protein may also be used to determine biological activity,
10 to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
15 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:78 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or
20 more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1660 of SEQ ID NO:78, b is an integer of 15 to 1674, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:78, and where b is greater than or equal to a + 14.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 69

The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 2 - 18 of the amino acid sequence referenced in Table 1 for this gene. Based upon these characteristics, it is believed that the protein
30 product of this gene shares structural features to type II membrane proteins.

In another embodiment, polypeptides comprising the amino acid sequence of the open reading frame upstream of the predicted signal peptide are contemplated by the present invention. Specifically, polypeptides of the invention comprise the following amino acid sequence:

- 5 K R A G V E V G G L V M A L A G S V F V L G G V L V L C V E R N G E G E M G W P Q H L P K S Q P L S
P P V A V R R C S F E R S W I D L L V E T S S S M V T C R Q Q V G T P N G M E G R G G G P K T T F P I R L
Q L S G A C A V R P E I Q W E V (SEQ ID NO: 259). Polynucleotides encoding these polypeptides are also provided.

- 10 This gene is expressed primarily in activated monocytes, dendritic cells, and in the tonsils.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic diseases and/or disorders, particularly
- 15 leukemia, lymphomas, tumors of hematopoietic origin. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hematopoietic system, expression of this gene at significantly higher or lower levels is routinely detected in
- 20 certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 25 Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 171 as residues: Gln-30 to Leu-38, Asn-75 to Thr-86. Polynucleotides encoding said polypeptides are also provided.

- The tissue distribution in activated monocytes, dendritic cells, and tonsils indicates that polynucleotides and polypeptides corresponding to this gene are useful
- 30 as a therapeutic and/or diagnostic agent for leukemias, lymphomas, and other diseases associated with cells of hematopoietic origin. Representative uses are described in the

"Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:79 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is

cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2177 of SEQ ID NO:79, b is an integer of 15 to 2191, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:79, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 70

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells, and to a lesser extent, other immune cells and tissue cell types, through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

The gene encoding the disclosed cDNA is believed to reside on chromosome 12. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 12.

This gene is expressed primarily in the placenta, brain, and liver and to a lesser extent in most other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hematopoietic, neurological, vascular, and developmental diseases and/or disorders, particularly cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and nervous systems, expression of this gene at significantly higher or lower levels is routinely detected in certain tissues or

cell types (e.g., hematopoietic, neurological, vascular, developmental, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, amniotic fluid, bile, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful therapeutic and/or diagnostic agent in a multitude of disease states, particularly those involving the immune and neurologic systems. Representative uses are described in the "Regeneration" and "Hyperproliferative Disorders" sections below, in Example 11, 15, and 18, and elsewhere herein. Briefly, the uses include, but are not limited to the detection, treatment, and/or prevention of Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates it plays a role in normal neural function.

Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the protein is useful in the detection, treatment, and/or prevention of a variety of vascular disorders and conditions, which include, but are not limited to microvascular disease, vascular leak syndrome, aneurysm, stroke, embolism, thrombosis, coronary artery disease, arteriosclerosis, and/or atherosclerosis. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement.

Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:80 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1321 of SEQ ID NO:80, b is an integer of 15 to 1335, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:80, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 71

The translation product of this gene shares sequence homology with the murine Fig1 (interleukin-four induced gene 1) which shares homology to the monoamine oxidases, particularly in domains responsible for FAD binding. Preferred polypeptides encoded by this gene comprise the following amino acid sequence:

QDWKAERSQDPFEKCMQDPDYEQLLKVTILEADNRIGGRIFTYRDQXTGWIG
ELGAMRMPSSHRILHKLCQGLGLNLTKFTQYDKNTWTEVHEXKL RNYVVEK
VPEKLG YALRPQEKGHSPEDIYQMALNQALKDLKALGCRKAMKKFERHTLL
EYLLGEGNLSRPAVQLLGDVMSDGGFFYLSFAEALRAXSCLSDRLQYSRIVG
GWDLLPRALLSSLSGLVLLNAPVVAMTQGP HDVHVQIETSPARNLKV LKAD
VVLLTASGPAVKRITFS (SEQ ID NO: 260), and/or

LPRHMQEALRRLHYVPATKVFLSFRPFWREEHIEGGHSNTDRPSRMIFYP PP
REGALLASYTWSDA AAFA GLSREEALRLALDDVAALHGPVVRQLWDGT
GVVKRWAEDQHSQGGFVVQXPALWQTEKDDWTVPYGRIYFAGEHTAYPHG
WVETAVKSALRAAIKINSRKGPASDTASPEGHASDMEGQGHVHGVASSPSH
DLAKEEGS (SEQ ID NO: 261). Polynucleotides encoding such polypeptides are also provided.

A preferred polypeptide fragment of the invention comprises the following amino acid sequence:

MAPLALHLLVLVPILLSLVASQDWKAERSQDPFEKCMQDPDYEQLLKVTIL
EADNRIGGRIFTYRDQXTGWIGELGAMRMPSSHRLHKLCQGLGLNLTKFTQ
5 YDKNTWTEVHEXKLRNYVVEKVPEKLG YALRPQEKGHSPEDIYQMALNQA
LKDLKALGCRKAMKKFERHTLLEYLLGEGNLSRPAVQLLGDVMSEDGFFYL
SFAEALRAXSCLSDRLQYSRIVGGWDL LPRALLSSLSGLVLLNAPVVAMTQG
PHDVH

VQIETSPARNLKVLKADVLLTASGPAVKRITFSPRCPATCRRRCGGCTTCR

10 PPRCS (SEQ ID NO: 262). Based on the sequence similarity, the translation product of this gene is expected to share at least some biological activities with monoamine oxidases, disintegrins, metalloproteinases, and apoptosis modulating proteins. Such activities are known in the art, some of which are described elsewhere herein.

Polynucleotides encoding these polypeptides are also provided.

15 The polypeptide of this gene has been determined to have a transmembrane domain at about amino acid position 235 - 251 of the amino acid sequence referenced in Table 1 for this gene. Moreover, a cytoplasmic tail encompassing amino acids 252 to 319 of this protein has also been determined. Based upon these characteristics, it is believed that the protein product of this gene shares structural features to type Ia

20 membrane proteins.

This gene is expressed primarily in hematopoietic cells, particularly in dendritic cells, and activated monocytes and to a lesser extent in T-cells, endothelial cells, and cells associated with ulcerative colitis.

Therefore, polynucleotides and polypeptides of the invention are useful as
25 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, leukemias, lymphomas, and diseases associated with antigen presenting cells, in addition to apoptosis dependant events. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological
30 probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression

of this gene at significantly higher or lower levels is routinely detected in certain tissues or cell types (e.g., immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g., lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred polypeptides of the present invention comprise immunogenic epitopes shown in SEQ ID NO: 173 as residues: Gln-22 to Gln-44, Ala-90 to Gly-95, Lys-137 to Trp-146, Arg-171 to Asp-181, Glu-370 to Ser-380, Asp-447 to Gly-452, Gln-463 to Trp-469, Asn-504 to Ala-510, Asp-512 to His-519, Ala-541 to Val-550, Asn-558 to His-566. Polynucleotides encoding said polypeptides are also provided.

The tissue distribution immune and hematopoietic cells and tissues, combined with the homology to the murine Fig 1 gene indicates that polynucleotides and polypeptides corresponding to this gene are useful as a therapeutic and/or diagnostic agent for hematopoietic diseases, especially those associated with antigen presenting cells. Representative uses are described in the "Immune Activity" and "infectious disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product is involved in the regulation of cytokine production, antigen presentation, or other processes suggesting a usefulness in the treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product is involved in immune functions. Therefore it is also useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous Disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's

Disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:81 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1853 of SEQ ID NO:81, b is an integer of 15 to 1867, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:81, and where b is greater than or equal to a + 14.

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
1	HISCN02	209878 05/18/98	pSport1	11	1113	1	1113	232	232	103	1	26	27	106
2	HHGDM70	209878 05/18/98	Lambda ZAP II	12	983	102	983	69	69	104	1	57	58	86
3	HHPGO40	209878 05/18/98	Uni-ZAP XR	13	973	1	973	68	68	105	1	37	38	302
3	HHPGO40	209878 05/18/98	Uni-ZAP XR	82	984	1	984	74	74	174	1	37	38	224
4	HAMGG68	209878 05/18/98	pCMVSPORT 3.0	14	1458	1	1458	312	312	106	1	20	21	55
5	HAPOM49	209878 05/18/98	Uni-ZAP XR	15	2005	1	2005	251	251	107	1	22	23	189
5	HAPOM49	209878 05/18/98	Uni-ZAP XR	83	2664	1	2664	448	448	175	1	1	2	123
6	HBGBA69	209878 05/18/98	Uni-ZAP XR	16	943	1	933	62	62	108	1	38	39	60
7	HBJFJ26	209878 05/18/98	Uni-ZAP XR	17	1503	588	1480	290	290	109	1	26	27	128

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
7	HBIFJ26	209878 05/18/98	Uni-ZAP XR	84	1328	413	1305	591	591	176	1	20	21	59
8	HCEDH38	209878 05/18/98	Uni-ZAP XR	18	1512	1	1438	222	222	110	1	26	27	68
9	HDPOJ08	209878 05/18/98	pCMVSPORT 3.0	19	1655	1	1655	159	159	111	1	18	19	122
10	HDPRX82	209878 05/18/98	pCMVSPORT 3.0	20	2525	1	2525	128	128	112	1	32	33	82
11	HELK31	209878 05/18/98	Uni-ZAP XR	21	1396	25	1334	209	209	113	1	29	30	344
11	HCNUA40	97898 02/26/97 209044 05/15/97	pBluescript	85	1342	949	1237	960	960	177	1	33	34	105
12	HFPCX64	209878 05/18/98	Uni-ZAP XR	22	1069	1	1069	181	181	114	1	28	29	181
12	HFPCX64	209878 05/18/98	Uni-ZAP XR	86	1154	84	1154	257	257	178	1	28	29	87
12	HCEBW71	209225 08/28/97	Uni-ZAP XR	87	1197	141	1197	257	257	179	1	28	29	87

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
13	HFXDO60	209878 05/18/98	Lambda ZAP II	23	1658	1	1658	131	131	115	1	46	47	115
14	HHEPG41	209878 05/18/98	pCMVSPORT 3.0	24	1077	385	1043	514	514	116	1	35	36	70
14	HAUAI83	209626 02/12/98	Uni-ZAP XR	88	910	1	886	253	253	180	1	37	38	49
14	HJPAZ83	209626 02/12/98	Uni-ZAP XR	89	1076	398	1076			181	1	11	12	23
15	HKGHA42	209878 05/18/98	pSPORT1	25	1205	1	1205	143	143	117	1	21	22	63
16	HMIAP86	209878 05/18/98	Uni-ZAP XR	26	1674	13	1674	182	182	118	1	19	20	334
17	HMUAP70	209878 05/18/98	pCMVSPORT 3.0	27	1965	531	1914	183	183	119	1	16	17	221
17	HMUAP70	209878 05/18/98	pCMVSPORT 3.0	90	1842	407	1783	413	413	182	1	25	26	103
17	HAGFY16	97923 03/07/97 209071 05/22/97	Uni-ZAP XR	91	1963	209	1922	251	251	183	1	28	29	198

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
17	HBMC37	209683 03/20/98	pBluescript	92	1487	79	1487	170	170	184	1	44	45	70
17	HFLQB16	209641 02/25/98	Uni-ZAP XR	93	1653	394	1637	413	413	185	1	25	26	82
17	HAGFY16	97923 03/07/97	Uni-ZAP XR	94	1830	87	1786	128	128	186	1	26	27	45
18	HRACJ35	209878 05/18/98	pCMVSPORT 3.0	28	1863	8	1863	99	99	120	1	24	25	472
18	HAWAZ34	209141 07/09/97	pBluescript SK-	95	1134	472	1132	687	687	187	1			33
19	HTWDE26	209878 05/18/98	pSport1	29	1626	1	1626	68	68	121	1	30	31	167
19	HMHBN40	97901 02/26/97 209047 05/15/97	Uni-ZAP XR	96	1772	69	1772	129	129	188	1	30	31	231
20	HUSIB13	209878 05/18/98	pSport1	30	605	1	605	172	172	122	1	32	33	46
21	HBAFA02	209877 05/18/98	pSport1	31	931	359	931	46	46	123	1	21	22	108

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
22	H2CBT75	209877 05/18/98	pBluescript SK-	32	1407	1	1407	32	32	124	1	23	24	60
23	HAGDQ42	209877 05/18/98	Uni-ZAP XR	33	1526	1	1526	126	126	125	1	18	19	248
24	HBMCI42	209877 05/18/98	pBluescript	34	1737	41	1580	244	244	126	1	44	45	248
25	HDPBQ71	209877 05/18/98	pCMVSPORT 3.0	35	2242	6	2242	24	24	127	1	33	34	612
26	HCEJG71	209877 05/18/98	Uni-ZAP XR	36	2235	2	2235	28	28	128	1	25	26	447
27	HELHL48	209877 05/18/98	Uni-ZAP XR	37	2971	560	2557	629	629	129	1	16	17	291
27	HSKCT36	209580 01/14/98	Uni-ZAP XR	98	1955	1	1955	31	31	190	1	18	19	184
28	HISAQ04	209877 05/18/98	pSport1	38	1163	1	1163	61	61	130	1	21	22	78
29	HJACB89	209877 05/18/98	pBluescript SK-	39	1932	28	1930	95	95	131	1	23	24	333
30	HTECC05	209877 05/18/98	Uni-ZAP XR	40	881	1	881	27	27	132	1	15	16	164

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
31	HBJLF01	209877 05/18/98	Uni-ZAP XR	41	1932	201	1931	217	217	133	1	46	47	244
32	HBXGP60	209877 05/18/98	ZAP Express	42	1164	1	1164	143	143	134	1	22	23	55
33	HCE5B20	209877 05/18/98	Uni-ZAP XR	43	1105	1	1105	237	237	135	1	25	26	54
34	HCMSQ56	209877 05/18/98	Uni-ZAP XR	44	1262	1	1262	148	148	136	1	19	20	88
35	HCNAH57	209877 05/18/98	Lambda ZAP II	45	517	1	517	35	35	137	1	33	34	61
36	HCUEP91	209877 05/18/98	ZAP Express	46	858	2	858	266	266	138	1	20	21	105
37	HDPCJ91	209877 05/18/98	pCMVSPORT 3.0	47	6107	1	6107	131	131	139	1	28	29	51
38	HDPGK25	209877 05/18/98	pCMVSPORT 3.0	48	703	1	703	345	345	140	1	33	34	119
39	HE2DY70	209877 05/18/98	Uni-ZAP XR	49	639	1	639	137	137	141	1	45	46	58
40	HE2NV57	209877 05/18/98	Uni-ZAP XR	50	867	1	867	99	99	142	1	36	37	99

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	5' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
41	HETBR16	209877 05/18/98	Uni-ZAP XR	51	1569	1	1569	161	161	143	1	21	22	64
42	HFXDGI3	209877 05/18/98	Lambda ZAP II	52	1196	1	1196	43	43	144	1	37	38	66
43	HFXY27	209877 05/18/98	Lambda ZAP II	53	945	1	945	44	44	145	1	19	20	58
44	HHPEC09	209877 05/18/98	Uni-ZAP XR	54	488	1	488	71	71	146	1	19	20	55
45	HISAD54	209877 05/18/98	pSport1	55	2860	1	2860	172	172	147	1	19	20	65
46	HJBCY35	209877 05/18/98	pBluescript SK-	56	1559	93	1272	232	232	148	1	23	24	327
47	HKAEA19	209877 05/18/98	pCMVSPORT 2.0	57	2064	1	1909	83	83	149	1	21	22	89
48	HKGDL36	209877 05/18/98	pSport1	58	1050	1	1050	55	55	150	1	33	34	148
49	HLDBS43	209877 05/18/98	pCMVSPORT 3.0	59	2533	1	2533	73	73	151	1	26	27	390
50	HLWAD92	209877 05/18/98	pCMVSPORT 3.0	60	899	1	899	197	197	152	1	34	35	98

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
51	HL YBI15	209877 05/18/98	pSport1	61	1079	1	1079	92	92	153	1	22	23	60
52	HMEJE05	209889 05/22/98	Lambda ZAP II	62	1928	1	1928	25	25	154	1	30	31	392
53	HNGIX55	209889 05/22/98	Uni-ZAP XR	63	781	1	781	121	121	155	1	19	20	74
54	HNHEX30	209889 05/22/98	Uni-ZAP XR	64	1194	1	1194	138	138	156	1	15	16	81
55	HPJBI33	209889 05/22/98	Uni-ZAP XR	65	1677	1	1677	236	236	157	1	31	32	53
56	HRABA80	209889 05/22/98	pCMVSPORT 3.0	66	1237	1	1237	130	130	158	1	28	29	102
57	HRACD80	209889 05/22/98	pCMVSPORT 3.0	67	1934	1	1934	191	191	159	1	16	17	575
57	HRACD80	209889 05/22/98	pCMVSPORT 3.0	99	1958	1	1958	191	191	191	1	16	17	146
58	HSLCX03	209889 05/22/98	Uni-ZAP XR	68	3300	984	2729	677	677	160	1	22	23	643
58	HSLCX03	209889 05/22/98	Uni-ZAP XR	100	2444	1	2444	392	392	192	1	22	23	124

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
59	HT5GJ57	209889 05/22/98	Uni-ZAP XR	69	1797	92	1797	122	122	161	1	25	26	190
60	HTACS42	209889 05/22/98	Uni-ZAP XR	70	1373	1	1373	213	213	162	1	29	30	63
61	HTEKE40	209889 05/22/98	Uni-ZAP XR	71	1579	1	1579	173	173	163	1	47	48	117
62	HTOBX69	209889 05/22/98	Uni-ZAP XR	72	1028	1	1028	28	28	164	1	20	21	42
63	HUVEO77	209889 05/22/98	Uni-ZAP XR	73	3674	1	3674	55	55	165	1	27	28	47
64	H2CBG48	209889 05/22/98	pBluescript SK-	74	2797	1	2797	125	125	166	1	25	26	45
65	H2CBU83	209889 05/22/98	pBluescript SK-	75	2703	1	2703	157	157	167	1	30	31	207
65	H2CBU83	209889 05/22/98	pBluescript SK-	101	2709	1	2709	157	157	193	1	30	31	51
66	HAPNY94	209889 05/22/98	Uni-ZAP XR	76	742	1	742	94	94	168	1	29	30	50
67	HBHJZ58	209889 05/22/98	Uni-ZAP XR	77	1825	1	1825	102	102	169	1	29	30	42

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
68	HCE2B33	209889 05/22/98	Uni-ZAP XR	78	1674	1	1668	67	67	170	1	18	19	55
69	HDPBQ02	209889 05/22/98	pCMVSPORT 3.0	79	2191	291	2191	460	460	171	1	24	25	108
70	HFYIY170	209889 05/22/98	pSPORT1	80	1335	1	1335	43	43	172	1	15	16	50
71	HDPOZ56	209889 05/22/98	pCMVSPORT 3.0	81	1867	415	1867	103	103	173	1	21	22	566
71	HDPOZ56	209889 05/22/98	pCMVSPORT 3.0	102	1722	1	1722	59	59	194	1	21	22	319

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X and the translated SEQ ID NO:Y are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further

below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in the deposited clone. These probes will also hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used to generate antibodies which bind specifically to the secreted proteins encoded by the cDNA clones identified in Table 1.

Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods. The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed

sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are species homologs. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for the
5 desired homologue.

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides
10 produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains
15 secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced
20 version of a polypeptide, including the secreted polypeptide, can be substantially purified by the one-step method described in Smith and Johnson, Gene 67:31-40 (1988). Polypeptides of the invention also can be purified from natural or recombinant sources using antibodies of the invention raised against the secreted protein in methods which are well known in the art.

25

Signal Sequences

Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of McGeoch, Virus Res. 3:271-286 (1985), uses the information from a short N-terminal
30 charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, Nucleic Acids Res. 14:4683-4690 (1986) uses the

information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. (von Heinje, supra.) However, the two
5 methods do not always produce the same predicted cleavage point(s) for a given protein.

In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., Protein Engineering 10:1-6 (1997)), which predicts the cellular location of a
10 protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1.

As one of ordinary skill would appreciate, however, cleavage sites sometimes
15 vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., + or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely
20 uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the
25 naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

30 **Polynucleotide and Polypeptide Variants**

"Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

- 5 By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the
- 10 polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire
- 15 sequence shown in Table 1, the ORF (open reading frame), or any fragment specified as described herein.

- As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known
- 20 computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence alignment the query and subject sequences
- 25 are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty
- 30 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95%

"identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences shown in Table 1 or to the amino acid sequence encoded by deposited DNA clone can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the the

query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter

the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred.

Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced

5 for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an
10 organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA
15 technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after
20 deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., *J. Biotechnology* 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological
25 activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem* 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every
30 possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See,

Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as to have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used.

(Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of the present invention having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a polypeptide to have an amino acid sequence which comprises the amino acid sequence of the present invention, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of the present invention or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

15

Polynucleotide and Polypeptide Fragments

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence contained in the deposited clone or shown in SEQ ID NO:X. The short nucleotide fragments are preferably at least about 20 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in the deposited clone or the nucleotide sequence shown in SEQ ID NO:X. These nucleotide fragments are useful as diagnostic probes and 25 primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments having a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401- 30 450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-

1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X or the cDNA contained in the deposited clone. In this context "about" includes the particularly recited ranges, larger or
5 smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein.

In the present invention, a "polypeptide fragment" refers to a short amino acid
10 sequence contained in SEQ ID NO:Y or encoded by the cDNA contained in the deposited clone. Protein fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-
15 40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes.

20 Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the
25 mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotide fragments encoding these polypeptide fragments are also preferred.

30 Also preferred are polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and

alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

- 5 Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are specifically contemplated by the present invention. Moreover, polynucleotide fragments encoding these domains are also contemplated.

- Other preferred fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.
- 10

Epitopes & Antibodies

- 15 In the present invention, "epitopes" refer to polypeptide fragments having antigenic or immunogenic activity in an animal, especially in a human. A preferred embodiment of the present invention relates to a polypeptide fragment comprising an epitope, as well as the polynucleotide encoding this fragment. A region of a protein molecule to which an antibody can bind is defined as an "antigenic epitope." In contrast, an "immunogenic epitope" is defined as a part of a protein that elicits an antibody response. (See, for instance, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998-4002 (1983).)
- 20

- Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, R. A., Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985) further described in U.S. Patent No. 4,631,211.)
- 25

- In the present invention, antigenic epitopes preferably contain a sequence of at least seven, more preferably at least nine, and most preferably between about 15 to about 30 amino acids. Antigenic epitopes are useful to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe, J. G. et al., Science 219:660-666 (1983).)
- 30

Similarly, immunogenic epitopes can be used to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow, M. et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle, F. J. et al., J. Gen. Virol. 66:2347-2354 (1985).) A preferred immunogenic epitope includes
5 the secreted protein. The immunogenic epitopes may be presented together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse) or, if it is long enough (at least about 25 amino acids), without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes
10 in a denatured polypeptide (e.g., in Western blotting.)

As used herein, the term "antibody" (Ab) or "monoclonal antibody" (Mab) is meant to include intact molecules as well as antibody fragments (such as, for example, Fab and F(ab')₂ fragments) which are capable of specifically binding to protein. Fab and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear
15 more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody. (Wahl et al., J. Nucl. Med. 24:316-325 (1983).) Thus, these fragments are preferred, as well as the products of a FAB or other immunoglobulin expression library. Moreover, antibodies of the present invention include chimeric, single chain, and humanized antibodies.

20

Fusion Proteins

Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the
25 polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, the polypeptides of the present invention can be used as targeting molecules once fused to other proteins.

Examples of domains that can be fused to polypeptides of the present
30 invention include not only heterologous signal sequences, but also other heterologous

functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, polypeptides of the present invention, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker et al., *Nature* 331:84-86 (1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., *J. Biochem.* 270:3958-3964 (1995).)

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See,

D. Bennett et al., J. Molecular Recognition 8:52-58 (1995); K. Johanson et al., J. Biol. Chem. 270:9459-9471 (1995).)

Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a peptide which facilitates purification of the fused polypeptide.

- 5 In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein.
- 10 Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

15

Vectors, Host Cells, and Protein Production

- The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral
- 20 vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

- The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the
- 25 vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

- The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac
- 30 promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The

expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., *Basic Methods In Molecular Biology* (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography,

phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

- 5 Polypeptides of the present invention, and preferably the secreted form, can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant,
- 10 insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine
- 15 encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.
- 20 In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide
- 25 sequences) that is operably associated with the polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24,
- 30 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al.,

Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

5

Uses of the Polynucleotides

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

10

The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

15

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids

20

containing the human gene corresponding to the SEQ ID NO:X will yield an amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides

25

can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome specific-cDNA libraries.

30

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however,

polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques." Pergamon Press, New York (1988).

5 For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides correspond to the noncoding regions of the cDNAs because the coding sequences are more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

10 Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins
15 University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined.
20 First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide
25 and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using
30 polynucleotides of the present invention. Any of these alterations (altered expression,

chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Both methods
5 rely on binding of the polynucleotide to DNA or RNA. For these techniques, preferred polynucleotides are usually 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J.
10 Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model systems, and the information disclosed herein can
15 be used to design antisense or triple helix polynucleotides in an effort to treat disease.

Polynucleotides of the present invention are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate
20 manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel.
25 In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA
30 markers for RFLP.

The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using
5 this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

Forensic biology also benefits from using DNA-based identification
10 techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co.
15 (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

20 There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by
25 organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination.

In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences
30 in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA

antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

5 Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods. (Jalkanen, M.,
10 et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell . Biol. 105:3087-3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such
15 as iodine (¹²⁵I, ¹²¹I), carbon (¹⁴C), sulfur (³⁵S), tritium (³H), indium (¹¹²In), and technetium (^{99m}Tc), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo
20 imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for
25 the relevant hybridoma.

A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ¹³¹I, ¹¹²In, ^{99m}Tc), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or
30 intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety

needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein.

- 5 In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which
10 involves (a) assaying the expression of a polypeptide of the present invention in cells or body fluid of an individual; (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a disorder.

- 15 Moreover, polypeptides of the present invention can be used to treat disease. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B), to inhibit the activity of a polypeptide (e.g., an oncogene), to
20 activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth).

Similarly, antibodies directed to a polypeptide of the present invention can
25 also be used to treat disease. For example, administration of an antibody directed to a polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as
30 molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also

be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

5

Biological Activities

The polynucleotides and polypeptides of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides and polypeptides could be used to treat the associated disease.

10

Immune Activity

A polypeptide or polynucleotide of the present invention may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, a polynucleotide or polypeptide of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

15

20

A polynucleotide or polypeptide of the present invention may be useful in treating or detecting deficiencies or disorders of hematopoietic cells. A polypeptide or polynucleotide of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency,

25

30

Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

5 Moreover, a polypeptide or polynucleotide of the present invention could also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, a polynucleotide or polypeptide of the present invention could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet
10 disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, a polynucleotide or polypeptide of the present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment of heart attacks (infarction), strokes, or scarring.

15 A polynucleotide or polypeptide of the present invention may also be useful in treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of a polypeptide or polynucleotide of the
20 present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

 Examples of autoimmune disorders that can be treated or detected by the present invention include, but are not limited to: Addison's Disease, hemolytic
25 anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary
30 Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by a polypeptide or polynucleotide of the present invention. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

A polynucleotide or polypeptide of the present invention may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of a polypeptide or polynucleotide of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, a polypeptide or polynucleotide of the present invention may also be used to modulate inflammation. For example, the polypeptide or polynucleotide may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

Hyperproliferative Disorders

A polypeptide or polynucleotide can be used to treat or detect hyperproliferative disorders, including neoplasms. A polypeptide or polynucleotide of the present invention may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, a polypeptide or polynucleotide of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

Examples of hyperproliferative disorders that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by a polynucleotide or polypeptide of the present invention. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

Infectious Disease

A polypeptide or polynucleotide of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, the polypeptide or polynucleotide of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide of the

present invention. Examples of viruses, include, but are not limited to the following DNA and RNA viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza), Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or diseases.

Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but not limited to, the following Gram-Negative and Gram-positive bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Norcardia), Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia, Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, Enterobacteriaceae (Klebsiella, Salmonella, Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), Pasteurellaceae Infections (e.g., Actinobacillus, Haemophilus, Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, and Staphylococcal. These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis,

opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis, Chlamydia, Syphilis,

5 Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or diseases.

10 Moreover, parasitic agents causing disease or symptoms that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but not limited to, the following families: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and
15 Trichomonas. These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), Malaria, pregnancy complications, and toxoplasmosis. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these
20 symptoms or diseases.

Preferably, treatment using a polypeptide or polynucleotide of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the
25 patient (ex vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

30 A polynucleotide or polypeptide of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See,

Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, 5 reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration 10 occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, a polynucleotide or polypeptide of the present invention may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. A polynucleotide or polypeptide of the present invention could also be used 15 prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using a 20 polynucleotide or polypeptide of the present invention to proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral 25 neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotide or polypeptide of the present invention.

30

Chemotaxis

A polynucleotide or polypeptide of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

A polynucleotide or polypeptide of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

It is also contemplated that a polynucleotide or polypeptide of the present invention may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, a polynucleotide or polypeptide of the present invention could be used as an inhibitor of chemotaxis.

Binding Activity

A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable

of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide, either as a secreted protein or on the cell
5 membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

10 The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

Alternatively, the assay can be carried out using cell-free preparations,
15 polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

20 Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

All of these above assays can be used as diagnostic or prognostic markers.
25 The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptide from suitably manipulated cells or tissues.

30 Therefore, the invention includes a method of identifying compounds which bind to a polypeptide of the invention comprising the steps of: (a) incubating a

candidate binding compound with a polypeptide of the invention; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with a polypeptide of the invention, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Other Activities

A polypeptide or polynucleotide of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

A polypeptide or polynucleotide of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, a polypeptide or polynucleotide of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

A polypeptide or polynucleotide of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, circadian rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

A polypeptide or polynucleotide of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

Other Preferred Embodiments

Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95%

identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of
5 positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of
10 positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Similarly preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the
15 range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide
20 sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

25 A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X
30 in Table 1.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X.

Also preferred is an isolated nucleic acid molecule which hybridizes under
5 stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

Also preferred is a composition of matter comprising a DNA molecule which
10 comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide
15 sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of
20 at least 50 contiguous nucleotides is included in the nucleotide sequence of the complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

25 A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule
30 comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X
5 wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and
10 determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence
15 selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

20 A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X
25 wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for identifying the species, tissue or cell type of a biological
30 sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least

one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of positions beginning with the residue at about the position of the First Amino Acid of the Secreted Portion and ending with the residue at about the Last Amino Acid of the

5 Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid

10 sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y.

15 Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

20 Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

25 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

30 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in

the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence
5 at least 95% identical to the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is an isolated antibody which binds specifically to a
10 polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained
15 in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group
20 consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least
25 one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence
30 selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide

comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino

acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

- 5 In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

- 25 Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

30 Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is

expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a secreted portion of a human secreted protein comprising an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is defined in Table 1; and an amino acid sequence of a secreted portion of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a secreted protein activity, which method comprises administering to such an individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to increase the level of said protein activity in said individual.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

20

Examples

Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector. Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 1 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

	<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited</u>
	<u>Plasmid</u>	
	Lambda Zap	pBluescript (pBS)
	Uni-Zap XR	pBluescript (pBS)
5	Zap Express	pBK
	lafmid BA	plafmid BA
	pSport1	pSport1
	pCMVSPORT 2.0	pCMVSPORT 2.0
	pCMVSPORT 3.0	pCMVSPORT 3.0
10	pCR [®] 2.1	pCR [®] 2.1
	<p>Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the f1 origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the f1 ori generates sense strand DNA and in the other, antisense.</p>	
25	<p>Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lafmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into E. coli strain</p>	
30		

XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., *Nuc. Acids Res.* 16:9677-9686 (1988) and Mead, D. et al., *Bio/Technology* 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ³²P-γ-ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for

bacterial colony screening (e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

Alternatively, two primers of 17-20 nucleotides derived from both ends of the
5 SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the
3' NT of the clone defined in Table 1) are synthesized and used to amplify the desired
cDNA using the deposited cDNA plasmid as a template. The polymerase chain
reaction is carried out under routine conditions, for instance, in 25 µl of reaction
mixture with 0.5 ug of the above cDNA template. A convenient reaction mixture is
10 1.5-5 mM MgCl₂, 0.01% (w/v) gelatin, 20 µM each of dATP, dCTP, dGTP, dTTP, 25
pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR
(denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1
min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The
amplified product is analyzed by agarose gel electrophoresis and the DNA band with
15 expected molecular weight is excised and purified. The PCR product is verified to be
the selected sequence by subcloning and sequencing the DNA product.

Several methods are available for the identification of the 5' or 3' non-coding
portions of a gene which may not be present in the deposited clone. These methods
include but are not limited to, filter probing, clone enrichment using specific probes,
20 and protocols similar or identical to 5' and 3' "RACE" protocols which are well
known in the art. For instance, a method similar to 5' RACE is available for
generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et
al., *Nucleic Acids Res.* 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a
25 population of RNA presumably containing full-length gene RNA transcripts. A
primer set containing a primer specific to the ligated RNA oligonucleotide and a
primer specific to a known sequence of the gene of interest is used to PCR amplify
the 5' portion of the desired full-length gene. This amplified product may then be
sequenced and used to generate the full length gene.

30 This above method starts with total RNA isolated from the desired source,
although poly-A+ RNA can be used. The RNA preparation can then be treated with

phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction
5 leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific
10 to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

15 A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X., according to the method described in Example 1. (See also, Sambrook.)

Example 3: Tissue Distribution of Polypeptide

20 Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by, among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P³² using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After
25 labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for mRNA expression.

Multiple Tissue Northern (MTN) blots containing various human tissues (H)
30 or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to

manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70°C overnight, and the films developed according to standard procedures.

5 **Example 4: Chromosomal Mapping of the Polynucleotides**

 An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95°C; 1 minute, 56°C; 1 minute, 70°C. This cycle is
10 repeated 32 times followed by one 5 minute cycle at 70°C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in
15 the particular somatic cell hybrid.

Example 5: Bacterial Expression of a Polypeptide

 A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA
20 sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc.,
25 Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

 The pQE-9 vector is digested with BamHI and XbaI and the amplified
30 fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain

M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r).

Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and
5 confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG
10 (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic
15 agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4°C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The
20 QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

25 The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4,
30 containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250

mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

In addition to the above expression vector, the present invention further
5 includes an expression vector comprising phage operator and promoter elements
operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC
Accession Number 209645, deposited on February 25, 1998.) This vector contains:
1) a neomycinphosphotransferase gene as a selection marker, 2) an *E. coli* origin of
replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a
10 Shine-Delgarno sequence, and 6) the lactose operon repressor gene (*lacIq*). The
origin of replication (*oriC*) is derived from pUC19 (LTI, Gaithersburg, MD). The
promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and
XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating
15 the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA
insert is generated according to the PCR protocol described in Example 1, using PCR
primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or
Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible
enzymes. The insert and vector are ligated according to standard protocols.

20 The engineered vector could easily be substituted in the above protocol to
express protein in a bacterial system.

Example 6: Purification of a Polypeptide from an Inclusion Body

The following alternative method can be used to purify a polypeptide
25 expressed in *E. coli* when it is present in the form of inclusion bodies. Unless
otherwise specified, all of the following steps are conducted at 4-10°C.

Upon completion of the production phase of the *E. coli* fermentation, the cell
culture is cooled to 4-10°C and the cells harvested by continuous centrifugation at
15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit
30 weight of cell paste and the amount of purified protein required, an appropriate
amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM

Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate
5 is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the
10 pellet is discarded and the polypeptide containing supernatant is incubated at 4°C overnight to allow further GuHCl extraction.

Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with
20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA
15 by vigorous stirring. The refolded diluted protein solution is kept at 4°C without mixing for 12 hours prior to further purification steps.

To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 µm membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The
20 filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

25 Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40
30 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium

acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A_{280} monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 μ g of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

Example 7: Cloning and Expression of a Polypeptide in a Baculovirus Expression System

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence identified in

Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in
5 Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment
10 then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a
15 commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified
20 by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five μ g of a plasmid containing the polynucleotide is co-transfected with 1.0 μ g of a commercially available linearized baculovirus DNA ("BaculoGold™
25 baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One μ g of BaculoGold™ virus DNA and 5 μ g of the plasmid are mixed in a sterile well of a microtiter plate containing 50 μ l of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 μ l Lipofectin plus 90 μ l Grace's medium are
30 added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded

in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27° C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27° C for four days.

- 5 After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell
- 10 culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 µl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded
- 15 in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4° C.

- To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of
- 20 infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 µCi of ³⁵S-methionine and 5 µCi ³⁵S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation.
- 25 The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

- 30 **Example 8: Expression of a Polypeptide in Mammalian Cells**

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human HeLa, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a

chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the vector does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five μ g of the expression plasmid pC6 is cotransfected with 0.5 μ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo

contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

15 Example 9: Protein Fusions

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the half-life time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

30 Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These

primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

```
GGGATCCGGAGCCCAAATCTTCTGACAAACTCACACATGCCCACCGTGC
CCAGCACCTGAATTCGAGGGTGCACCGTCAGTCTTCTCTTCCCCCAAAA
CCCAAGGACACCCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGT
GGTGGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG
ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTA
CAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACT
GGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCA
ACCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAAC
CACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAG
GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCCGT
GGAGTGGGAGAGCAATGGGCAGCCGGAACAACACTACAAGACCACGCCT
CCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTG
GACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCA
TGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG
GTAAATGAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)
```

Example 10: Production of an Antibody from a Polypeptide

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) For example, cells expressing a polypeptide of the present invention is administered to an animal to induce the
5 production of sera containing polyclonal antibodies. In a preferred method, a preparation of the secreted protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

In the most preferred method, the antibodies of the present invention are
10 monoclonal antibodies (or protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology. (Köhler et al., Nature 256:495 (1975); Köhler et al., Eur. J. Immunol. 6:511 (1976); Köhler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures
15 involve immunizing an animal (preferably a mouse) with polypeptide or, more preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about
20 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are
25 selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

Alternatively, additional antibodies capable of binding to the polypeptide can
30 be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is

possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody
5 whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies.

It will be appreciated that Fab and F(ab')₂ and other fragments of the
10 antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). Alternatively, secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic
15 chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in
20 the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

25

Example 11: Production Of Secreted Protein For High-Throughput Screening Assays

The following protocol produces a supernatant containing a polypeptide to be tested. This supernatant can then be used in the Screening Assays described in
30 Examples 13-20.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37°C for 6 hours.

While cells are incubating, prepare appropriate media. either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of CaCl₂ (anhyd); 0.00130 mg/L CuSO₄·5H₂O; 0.050 mg/L of Fe(NO₃)₃·9H₂O; 0.417 mg/L of FeSO₄·7H₂O; 311.80 mg/L of KCl; 28.64 mg/L of MgCl₂; 48.84 mg/L of MgSO₄; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO₃; 62.50 mg/L of NaH₂PO₄·H₂O; 71.02 mg/L of Na₂HPO₄; .4320 mg/L of ZnSO₄·7H₂O; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H₂O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL-H₂O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H₂O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H₂O; 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and 0.680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37°C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

5 On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the
10 polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

15 Example 12: Construction of GAS Reporter Construct

One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The
20 binding of a protein to these elements alter the expression of the associated gene.

GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in
25 many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus
30 upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2,

Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, *Ann. Rev. Biochem.* 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN- α , IFN- γ , and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

	<u>Ligand</u>	<u>tyk2</u>	<u>JAKs</u> <u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>	<u>STATs</u>	<u>GAS(elements) or ISRE</u>
	<u>IFN family</u>						
5	IFN-a/B	+	+	-	-	1,2,3	ISRE
	IFN-g		+	+	-	1	GAS (IRF1>Lys6>IFP)
	Il-10	+	?	?	-	1,3	
	<u>gp130 family</u>						
10	IL-6 (Pleiotrophic)	+	+	+	?	1,3	GAS (IRF1>Lys6>IFP)
	Il-11(Pleiotrophic)	?	+	?	?	1,3	
	OnM(Pleiotrophic)	?	+	+	?	1,3	
	LIF(Pleiotrophic)	?	+	+	?	1,3	
	CNTF(Pleiotrophic)	-/+	+	+	?	1,3	
15	G-CSF(Pleiotrophic)	?	+	?	?	1,3	
	IL-12(Pleiotrophic)	+	-	+	+	1,3	
	<u>g-C family</u>						
	IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
20	IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 = IFP >>Ly6)(IgH)
	IL-7 (lymphocytes)	-	+	-	+	5	GAS
	IL-9 (lymphocytes)	-	+	-	+	5	GAS
	IL-13 (lymphocyte)	-	+	?	?	6	GAS
	IL-15	?	+	?	+	5	GAS
25	<u>gp140 family</u>						
	IL-3 (myeloid)	-	-	+	-	5	GAS (IRF1>IFP>>Ly6)
	IL-5 (myeloid)	-	-	+	-	5	GAS
	GM-CSF (myeloid)	-	-	+	-	5	GAS
30	<u>Growth hormone family</u>						
	GH	?	-	+	-	5	
	PRL	?	+/-	+	-	1,3,5	
	EPO	?	-	+	-	5	GAS(B-CAS>IRF1=IFP>>Ly6)
35	<u>Receptor Tyrosine Kinases</u>						
	EGF	?	+	+	-	1,3	GAS (IRF1)
	PDGF	?	+	+	-	1,3	
	CSF-1	?	+	+	-	1,3	GAS (not IRF1)
40							

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

10 5':GCGCCTCGAGATTTCCCCGAAATCTAGATTTCCCCGAAATGATTTCCCC
GAAATGATTTCCCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

15 PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

20 5':CTCGAGATTTCCCCGAAATCTAGATTTCCCCGAAATGATTTCCCCGAAA
TGATTTCCCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCG
CCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTCCGCCCATTCT
CCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCC
TCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCT
25 AGGCTTTTGCAAAAAGCTT:3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter
30 molecules that can be used instead of SEAP include chloramphenicol

acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

Example 13: High-Throughput Screening Assay for T-cell Activity.

The following protocol is used to assess T-cell activity by identifying factors, such as growth factors and cytokines, that may proliferate or differentiate T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells

(ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1% Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

During the incubation period, count cell concentration, spin down the required number of cells (10^7 per transfection), and resuspend in OPTI-MEM to a final concentration of 10^7 cells/ml. Then add 1ml of 1×10^7 cells in OPTI-MEM to T25 flask and incubate at 37°C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with supernatants containing a polypeptide as produced by the protocol described in Example 11.

On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100, 000 cells per well).

After all the plates have been seeded, 50 ul of the supernatants are transferred
5 directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed
10 in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophene covers) and stored at -20°C until SEAP assays are performed according to Example 17. The
15 plates containing the remaining treated cells are placed at 4°C and serve as a source of material for repeating the assay on a specific well if desired.

As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

The above protocol may be used in the generation of both transient, as well as,
20 stable transfected cells, which would be apparent to those of skill in the art.

Example 14: High-Throughput Screening Assay Identifying Myeloid Activity

The following protocol is used to assess myeloid activity by identifying factors, such as growth factors and cytokines, that may proliferate or differentiate
25 myeloid cells. Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

30 To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell

Growth & Differentiation, 5:259-265) is used. First, harvest 2×10^7 U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

- 5 Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 μ g GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 μ M $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 1 mM MgCl_2 , and 675 μ M CaCl_2 . Incubate at 37°C for 45 min.

- 10 Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37°C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 μ g/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 μ g/ml G418 for couple of passages.

- 15 These cells are tested by harvesting 1×10^8 cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of 5×10^5 cells/ml. Plate 200 μ l cells per well in the 96-well plate (or 1×10^5 cells/well).

- 20 Add 50 μ l of the supernatant prepared by the protocol described in Example 11. Incubate at 37°C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.

- 25 When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat pheochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor).

- 5 The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene
10 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

- Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified
15 product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

- To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30
20 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

- PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-
25 inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.

- Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine
30 protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine

growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as 5×10^5 cells/ml.

Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to 1×10^5 cells/well). Add 50 ul supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 17.

Example 16: High-Throughput Screening Assay for T-cell Activity

NF- κ B (Nuclear Factor κ B) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF- κ B regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- κ B appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

In non-stimulated conditions, NF- κ B is retained in the cytoplasm with I- κ B (Inhibitor κ B). However, upon stimulation, I- κ B is phosphorylated and degraded, causing NF- κ B to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- κ B include IL-2, IL-6, GM-CSF, ICAM-1 and class I MHC.

Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF- κ B promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF- κ B would be useful in treating diseases. For example, inhibitors of NF- κ B could be used to treat those
5 diseases related to the acute or chronic activation of NF- κ B, such as rheumatoid arthritis.

To construct a vector containing the NF- κ B promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF- κ B binding site (GGGGACTTTCCC) (SEQ ID NO:8), 18 bp of sequence complementary
10 to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:
5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:
15 5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene)
Sequencing with the T7 and T3 primers confirms the insert contains the following
20 sequence:

5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCC
ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCC
ATCCCGCCCCCTAACTCCGCCCAGTCCGCCCCATTCTCCGCCCCATGGCTGA
25 CTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTA
TTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAA
GCTT:3' (SEQ ID NO:10)

Next, replace the SV40 minimal promoter element present in the pSEAP2-
30 promoter plasmid (Clontech) with this NF- κ B/SV40 fragment using XhoI and

HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF- κ B/SV40/SEAP cassette is removed from the above NF- κ B/SEAP vector using restriction enzymes
5 SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly, the NF- κ B/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

Once NF- κ B/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly,
10 the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 13. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

Example 17: Assay for SEAP Activity

15 As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 μ l of 2.5x
20 dilution buffer into Optiplates containing 35 μ l of a supernatant. Seal the plates with a plastic sealer and incubate at 65°C for 30 min. Separate the Optiplates to avoid uneven heating.

Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 μ l Assay Buffer and incubate at room
25 temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 μ l Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

30 Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

Reaction Buffer Formulation:

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75
14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6
23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

Example 18: High-Throughput Screening Assay Identifying Changes in Small**5 Molecule Concentration and Membrane Permeability**

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants

which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

5 The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.

10 For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO₂ incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

15 A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37°C in a CO₂ incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

20 For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2-5x10⁶ cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37°C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1x10⁶ cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final
25 volume.

 For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

30 To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm;

and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular Ca^{++} concentration.

5 **Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity**

 The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase (RPTK) group are receptors for a range of mitogenic and metabolic growth
10 factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

 Activation of RPTK by ligands involves ligand-mediated receptor
15 dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily
20 of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

 Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of activating tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins
25 capable of activating the tyrosine kinase signal transduction pathways.

 Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr
30 with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St.

Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4°C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 11, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na₃VO₄, 2 mM Na₄P₂O₇ and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4°C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4°C at 16,000 x g.

Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and

PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg₂₊ (5mM ATP/50mM MgCl₂), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl₂, 5 mM MnCl₂, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30°C for 2 min. Initiate the reaction by adding 10ul of the control enzyme or the filtered supernatant.

10 The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37°C for 20 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide.

15 Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37°C for one hour. Wash the well as above.

Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

20

25 **Example 20: High-Throughput Screening Assay Identifying Phosphorylation Activity**

As a potential alternative and/or complement to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine

30

phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4°C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place

of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

Example 21: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95°C for 30 seconds; 60-120 seconds at 52-58°C; and 60-120 seconds at 70°C, using buffer solutions described in Sidransky, D., et al., Science 252:706 (1991).

PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies). The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

PCR products is cloned into T-tailed vectors as described in Holton, T.A. and Graham, M.W., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson, Cg. et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson, Cv.

et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

10 A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

15 For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

25 Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

30 Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard

curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

5

Example 23: Formulating a Polypeptide

The secreted polypeptide composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the secreted polypeptide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of secreted polypeptide administered parenterally per dose will be in the range of about 1 $\mu\text{g/kg/day}$ to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day , and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the secreted polypeptide is typically administered at a dose rate of about 1 $\mu\text{g/kg/hour}$ to about 50 $\mu\text{g/kg/hour}$, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing the secreted protein of the invention are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The secreted polypeptide is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 5 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., *Biopolymers* 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (R. Langer et al., *J. Biomed. Mater. Res.* 15:167-277 (1981), and R. Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions 10 also include liposomally entrapped polypeptides. Liposomes containing the secreted polypeptide are prepared by methods known per se: DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci. USA* 82:3688-3692 (1985); Hwang et al., *Proc. Natl. Acad. Sci. USA* 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and 15 EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal secreted polypeptide therapy.

For parenteral administration, in one embodiment, the secreted polypeptide is 20 formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other 25 compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting the polypeptide uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood 30 of the recipient. Examples of such carrier vehicles include water, saline, Ringer's

solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The secreted polypeptide is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

Any polypeptide to be used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical

compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the

5 polypeptides of the present invention may be employed in conjunction with other therapeutic compounds.

Example 24: Method of Treating Decreased Levels of the Polypeptide

It will be appreciated that conditions caused by a decrease in the standard or

10 normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a pharmaceutical composition comprising an amount of the polypeptide to

15 increase the activity level of the polypeptide in such an individual.

For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 23.

20

Example 25: Method of Treating Increased Levels of the Polypeptide

Antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as

25 cancer.

For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense

30 polynucleotide is provided in Example 23.

Example 26: Method of Treatment Using Gene Therapy

One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37°C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector.

The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

Example 27: Method of Treatment Using Gene Therapy - In Vivo

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata H. et al. (1997) Cardiovasc. Res. 35(3):470-479, Chao J et al. (1997) Pharmacol. Res. 35(6):517-522, Wolff J.A. (1997) Neuromuscul. Disord. 7(5):314-318, Schwartz B. et al. (1996) Gene Ther. 3(5):405-411, Tsurumi Y. et al. (1996) Circulation 94(12):3281-3290 (incorporated herein by reference).

The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection

into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 μ m cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be used to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

Example 28: Transgenic Animals.

The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (i.e., polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e.g., Ulmer et al., Science 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-

mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

Any technique known in the art may be used to produce transgenic clones
5 containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

The present invention provides for transgenic animals that carry the transgene
10 in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory
15 sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is
20 to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only
25 that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the
30 recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA

expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also
5 be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more
10 than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for
15 screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which include, but are not
20 limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

25 **Example 29: Knock-Out Animals.**

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (*E.g.*, see Smithies et al., *Nature* 317:230-234 (1985); Thomas & Capecchi, *Cell* 51:503-512 (1987); Thompson et al., *Cell* 5:313-321 (1989); each of which is incorporated by
30 reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding

regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (e.g., see Thomas & Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and
5 Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For
10 example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological
15 function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

It will be clear that the invention may be practiced otherwise than as
20 particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other
25 disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties.

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>178</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>May 18, 1998</u>	Accession Number <u>209878</u>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	
For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application Authorized officer <u>Misty Walker</u> <u>International Division</u> <u>703-606-6032</u> <u>mwalker@uspto.gov</u>	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer _____

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>179</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit February 26, 1997	Accession Number 97898
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer M. J. Baker International Division TCS-000-0002 Washington, D.C. 20540	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>179</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 15, 1997	Accession Number 209044
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	
For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application Authorized officer [Signature] [Stamp: International Bureau] [Stamp: 708-010-100]	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>179</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit August 28, 1997	Accession Number 209225
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer [Signature] 705-615-2092	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>180</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit March 7, 1997	Accession Number 97923
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer Misty Walker International Division 708-595-0000	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>180</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 22, 1997	Accession Number 209071
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer [Signature] [Stamp: International Division, PCT/US-0002, international@uspto.gov]	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>180</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit February 12, 1998	Accession Number 209626
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<div style="text-align: right; font-size: small; margin-bottom: 5px;">For receiving Office use only</div> <div style="display: flex; align-items: center;"><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div style="border-top: 1px solid black; padding-top: 5px;">Authorized officer <i>[Signature]</i> International Division 709-600-6000</div>	<div style="text-align: right; font-size: small; margin-bottom: 5px;">For International Bureau use only</div> <div style="display: flex; align-items: center;"><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div style="border-top: 1px solid black; padding-top: 5px;">Authorized officer</div>

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>181</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit March 20, 1998	Accession Number 209683
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer [Signature] [Stamp: RECEIVED DIVISION]	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>181</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>February 25, 1998</u>	Accession Number <u>209641</u>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	
For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application Authorized officer <u>[Signature]</u> <u>[Stamp]</u>	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer


INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>181</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit July 9, 1997	Accession Number 209141
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application Authorized officer <i>[Signature]</i>	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer


INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>181</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit February 26, 1997	Accession Number 97901
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>181</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 15, 1997	Accession Number 209047
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>181</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 18, 1998	Accession Number 209877
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer Christy Weisler International Division 703-605-6832	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>182</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit January 14, 1998	Accession Number 209580
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	
For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application Authorized officer 	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>185</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 22, 1998	Accession Number 209889
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer [Signature] 7/26/98	Authorized officer

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:
 - 5 (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a
10 polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - 15 (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X,
20 having biological activity;
 - (f) a polynucleotide which is a variant of SEQ ID NO:X;
 - (g) a polynucleotide which is an allelic variant of SEQ ID NO:X;
 - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
 - 25 (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.
- 5 3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 10 4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 15 5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 20 6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 25 7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.
9. A recombinant host cell produced by the method of claim 8.
- 30 10. The recombinant host cell of claim 9 comprising vector sequences.

11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:
- (a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - 5 (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;
 - (c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included
10 in ATCC Deposit No:Z;
 - (e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (f) a full length protein of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - 15 (g) a variant of SEQ ID NO:Y;
 - (h) an allelic variant of SEQ ID NO:Y; or
 - (i) a species homologue of the SEQ ID NO:Y.
12. The isolated polypeptide of claim 11, wherein the secreted form or the full length protein comprises sequential amino acid deletions from either the C-
20 terminus or the N-terminus.
13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.
- 25 14. A recombinant host cell that expresses the isolated polypeptide of claim 11.
15. A method of making an isolated polypeptide comprising:
- (a) culturing the recombinant host cell of claim 14 under conditions such that
30 said polypeptide is expressed; and
 - (b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount
5 of the polypeptide of claim 11 or the polynucleotide of claim 1.

18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:
(a) determining the presence or absence of a mutation in the polynucleotide of
10 claim 1; and
(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

19. A method of diagnosing a pathological condition or a susceptibility to
15 a pathological condition in a subject comprising:
(a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and
(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

20
20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:
(a) contacting the polypeptide of claim 11 with a binding partner; and
(b) determining whether the binding partner effects an activity of the
25 polypeptide.

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.

22. A method of identifying an activity in a biological assay, wherein the
30 method comprises:
(a) expressing SEQ ID NO:X in a cell;
(b) isolating the supernatant;

- (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 20.

Sequence Listing

<110> Human Genome Sciences, Inc., et al.

<120> 71 Human Secreted Proteins

<130> PZ030PCT

<140> Unassigned

<141> 1999-07-14

<150> 60/092,956

<151> 1998-07-15

<150> 60/092,921

<151> 1998-07-15

<150> 60/092,922

<151> 1998-07-15

<160> 262

<170> PatentIn Ver. 2.0

<210> 1

<211> 733

<212> DNA

<213> Homo sapiens

<400> 1

```

gggatccgga gcccaaatct tctgacaaaa ctcacacatg cccaccgtgc ccagcacctg      60
aattcgaggg tgcaccgtca gtcttcctct tccccccaaa acccaaggac accctcatga      120
tctcccggac tcctgaggtc acatgcgtgg tggtaggagc aagccacgaa gaccctgagg      180
tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg      240
aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact      300
ggctgaatgg caaggagtac aagtgcaagg tctccaacaa agccctccca acccccatcg      360
agaaaacccat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac accctgcccc      420
catcccggga tgagctgacc aagaaccagg tcagcctgac ctgcctgggc aaaggcttct      480
atccaaagcga catcgccgtg gagtgggaga gcaatgggca gccggagaac aactacaaga      540
ccacgcctcc cgtgctggac tccgacggct ccttcttctc ctacagcaag ctcaccgtgg      600
acaagagcag gtggcagcag gggaacgtct tctcatgtc cgtgatgcat gaggctctgc      660
acaaccacta cagcgagaag agcctctccc tgtctccggg taaatgagtg cgacggccgc      720
gactctagag gat                                     733

```

<210> 2

<211> 5

<212> PRT

<213> Homo sapiens

<220>

<221> Site

<222> (3)

<223> Xaa equals any of the twenty naturally occurring L-amino acids

<400> 2

Trp Ser Xaa Trp Ser

1

5

<210> 3
 <211> 86
 <212> DNA
 <213> Homo sapiens

 <400> 3
 gcgcctcgag atttccccga aatctagatt tccccgaaat gatttccccg aaatgatttc 60
 cccgaaatat ctgccatctc aattag 86

<210> 4
 <211> 27
 <212> DNA
 <213> Homo sapiens

 <400> 4
 gcggcaagct ttttgcaaag cctagggc 27

<210> 5
 <211> 271
 <212> DNA
 <213> Homo sapiens

 <400> 5
 ctcgagattt ccccgaaatc tagatttccc cgaaatgatt tccccgaaat gatttccccg 60
 aaatatctgc catctcaatt agtcagcaac catagtcccg cccctaactc cgcccatccc 120
 gccctaact ccgcccagtt ccgcccattc tccgcccacat ggetgactaa ttttttttat 180
 ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt gaggaggctt 240
 ttttgagggc ctaggctttt gcaaaaagct t 271

<210> 6
 <211> 32
 <212> DNA
 <213> Homo sapiens

 <400> 6
 gcgcctcgagg gatgacagcg atagaacccc gg 32

<210> 7
 <211> 31
 <212> DNA
 <213> Homo sapiens

 <400> 7
 gcgaagcttc gcgactcccc ggatccgcct c 31

<210> 8
 <211> 12
 <212> DNA
 <213> Homo sapiens

 <400> 8
 ggggactttc cc 12

<210> 9
 <211> 73
 <212> DNA
 <213> Homo sapiens

<400> 9
 gcggcctcga ggggactttc ccggggactt tccggggact ttccgggact ttccatcctg 60
 ccattctcaat tag 73

<210> 10
 <211> 256
 <212> DNA
 <213> Homo sapiens

<400> 10
 ctccgagggga ctttcccgagg gactttccgg ggactttccg ggactttcca tctgccatct 60
 caattagtca gcaaccatag tcccgcacct aactccgccc atcccgcgcc taactccgcc 120
 cagttccgcc cattctccgc cccatggctg actaattttt tttatttatg cagaggccga 180
 ggccgcctcg gcctctgagc tattccagaa gtagtgagga ggcttttttg gaggcctagg 240
 cttttgcaaa aagctt 256

<210> 11
 <211> 1113
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (393)
 <223> n equals a,t,g, or c

<400> 11
 gatgctcctt tagcttggag gagtttgta ttaccacact tctgaagcct acttctgtca 60
 attcatccaa ctcatctca gtccagtttt gtttccttgc tggtagaggag ttgtgatcct 120
 ttggaggaga agaggcattc tgggtttttg aatttttagc cattttgctc tgggtttcttc 180
 ccattctttgt ggatttatct acctttcatc ttcaatgtta gtgacctatg gatggggtct 240
 ctgagtggat gtgctcttcc tttctgtttg twagttttct ttctaacagt tagccctct 300
 gctgtaggtc tgctggaktt tgctggaggt ccactccaga ccctgtttgc ctgggtatca 360
 ccagtggagg ctgcagaaca gcaaagattg ctncctgttc tttcctctgg aagcttcgtc 420
 tcagagggca cctgccagat gccagccaga gctctctgt atgaggtgtc tgttgcccca 480
 tactgggaga ttctctccag tcaggatata aggaggtcag ggacctactt gaggaggcag 540
 tctgacctt agcagagggt gaacactgtg ctaggaggtc ctctgctctt ttcagagctg 600
 tcaggcgggg cgtataagtc tgctgaatct gtgtccgcag ccacctcttc cccaggtgc 660
 tctgtccag ggagatgggg gttttatttt taagtcccca actggggctg ctgccttttt 720
 ttcagagatg ccctgccag agaggagaaa tctggcagtc tggcctcaga ggccttgctg 780
 agctgccgtt ggctccacct agttcaaact tcccaagggg ctttgtttat actgtaaggg 840
 gaaaaccgcc tactcgagcc tcatcaatgg cagacacccc tcccgcgcc aagcttaagt 900
 gtcccaggtt gatctcagac tgctgctgtg ctggcagtgga gaatttcaag ccagtggatc 960
 ttagtttctg gtgctctgtg ggggtgggac ccattgaacc agactactcg gctccctggc 1020
 ttcagccccc tttccaggag agtgaagggt tctgtctcat tggcattcca ggagtcactg 1080
 gtgtatggaa aaaaaaaaaa aaaggcgccg cgc 1113

<210> 12

<211> 983
 <212> DNA
 <213> Homo sapiens

<400> 12
 ggcacgaggg cagctgcaga gctccaggtt tctctgcccc caagggcagg ggctgcccc 60
 cgcccaggat gactctgcct tccagagcct tggcctccct gggggtggga gtgtggggga 120
 tgctaagggtt aaatcagggt acagtaagtt gtgggggcag caggtggagc agcagagtgg 180
 cactggggagc tttctcttgg gtgtgcggtg tggccttggt tctgcagcca tcaggtgggg 240
 gcttggggact gacttctcct tctgaaggat gctgggaagg tgagctggct ttggcagtgc 300
 ttagagctcc ggggggttcc cctcctaga acatgcaagc tctcacaccg gtgcgtcatc 360
 atcacaccca tcatcaagcc cacagtggta tactgaacac ctgccccaca aagacggtgg 420
 actgctctca gaggagcccc atgaaccacc gatggttaca actatccaat gcctgatggc 480
 agacagccag gccaacctcg gcttccactc tctcttctc accctacaat cagccaaagt 540
 gacctgagtc atgtagtgtg aagttgcttt ctgctttctc ttgtttgtgc ttttgcgtgt 600
 tcttctgccc catactttgt taactccatg agttaaagtc taccattttt cccagacaag 660
 tgctgcttct gcaaggaaac ccttccctgat cccccaccta tctgaaaagt acctctccag 720
 cttgcttctt caggggtgctg agcgttccct cccagcctgt catcaccttc ctccatacgc 780
 tatggtgtgt tctgtcttct tctagtcttg tctctttttt tctgttagat tgtagctcct 840
 tgctgacagg aaccacgcct gctccagctt catacctccc actgctacag cacagaacct 900
 gcttctcaga cttacagcaa atgtttgttt gctgaatgaa ttaattaaag ataaagcaaa 960
 aaaaaaaaaa aaaaaaactc gag 983

<210> 13
 <211> 973
 <212> DNA
 <213> Homo sapiens

<400> 13
 ggcacgagcc cagcggaagc caagccacca ggccccccag cgtccacgcg gagcatgaac 60
 attgaggatg gcgcgtgccc gcggctcccc gtgccccccg ctgcccgcgc gtaggatgtc 120
 ctggccccac ggggcattgc tcttctctct gctcttctcc ccacctctgg gggccggtgg 180
 aggtggagtg gccgtgacgt ctgcccgcgc agggggctcc ccgcgggcca cctcctgccc 240
 cgtggcctgc tctgcagca accaggccag ccgggtgatc tgcacacgga gagacctggc 300
 cgagggtccc gccagcatcc cgggtcaacac gcggtacctg aacctgcaag agaacggcat 360
 ccagggtgatc cggacggaca cgttcaagca cctgcccgcac ctggagattc tgcagctgag 420
 caagaacctg gtgcgcaaga tgcaggtggg cgcttcaac gggctgcccc gcctcaacac 480
 gctggagctt tttgacaacc ggctgaccac ggtgcccacg caggccttcg agtacctgtc 540
 caagctgcgg gagctctggc tgcggaacaa ccccatcgag agcatcccct cctacgcctt 600
 caaccgcgtg ccttcgctgc ggccgctgga cctgggcgag ctcaagcggc tgggaatacat 660
 ctccggaggcg gccttcgagg ggtggtcaa cctgcgctac ctcaacctgg gcatgtgcaa 720
 cctcaaggac atcccccaacc tgacggccct ggtgcgcctg gaggagctgg agctgtcggg 780
 caaccggctg gacctgatcc gcccggtc ctccagggt ctcaccagcc tgcgcaagct 840
 gtggctcatg cagcccagg tagccaccat cgagcgcaac gccttcgacg acctcaagtc 900
 gctggaggag ctcaacctgt cccacaacaa cctgatgtcg ctgccccacg acctcttccac 960
 gccctgcac cgc 973

<210> 14
 <211> 1458
 <212> DNA
 <213> Homo sapiens

<400> 14
 ccacgcgtcc gggaattttc aaaagatcca aacagagact tcttgcattc tctgcctttc 60
 caacagaagc ggtgatcgtc taagtatgag cctgtggctt cctttgtgca tttgagcatg 120
 ctgtaattaa gatgatgata gtttcttaga aaaagctttc ctgaatccct ctgacgttgc 180

ctgggatctt	tctgttgatt	cgtcttttct	ggagattggg	acagagcatc	tgtggtccag	240
ggaagttagt	cctctggcct	caattctgtt	gtggatgtgc	agtataaagc	gggcattgag	300
tgcctcgggg	gatgcctagt	tcgtggcttc	ctggctgttt	tgtccttctg	tgtctttagt	360
ctgtagggtg	ccagctcagg	gagtggggtg	ttggcggcgt	ttccgcgggt	ggcctccttg	420
ctttgccgca	cctccagggt	ctgggcatga	gaggccgtgg	cctcatttct	ggtggataac	480
cttttttagt	taatagcata	tttaattaga	tcacagcatt	gaattcaaaa	tttcttctgc	540
aaagaaagtt	gtggggcata	agacaccggg	aatgagggag	gaggaagaca	gttgtgtttt	600
ctctttaaac	cttgagctct	agccgatgca	tttgtcagga	aatacagcac	tttgtcttaa	660
gaaaacaagg	aaggaggcgg	ggcgagtggt	ctcacgcctg	taateccagc	actttgggag	720
gccgaggcgg	gcggatcacc	tgaggtgggg	agtatgagac	caccctgact	aacatggaga	780
gacctgtctt	ctactaaaag	tacagaatta	gccgggctgt	gttgcgcatg	cccataatcc	840
cagctactga	ggagacttga	ggtaggagaa	tcacttgaac	ctcagcggcg	gaggttgcag	900
tgagtcgaga	tcgcgccagt	gcactccagc	ctgggcaaga	agagcgaaac	tgggtctcaa	960
gttaaaaaaa	gaaagcaagg	aaagagtaat	ttacaacgaa	ggaaaaaaac	ccacagcaca	1020
cccttcgcgg	ctgtcagcgc	tctcctgatg	tcacagtggc	tgctgtctct	tgggtgtggg	1080
gaggtgtggg	gagcccagcc	cctggccctg	cctcccgccg	cccgtctccc	ttctctctct	1140
tactcgggtt	agccatagcg	aggcctcccg	tcgtttcaga	tatgaatttg	ttttatagat	1200
tataaatatg	catataatag	gtatgtataa	agcagaatgc	ctgcctttcc	tggttatatt	1260
ttgtaccata	ttgtaaatta	tattatttat	tctttaccaa	ttttgggaat	aaaaggtggt	1320
ttggttatatt	aataataata	gagctgttaa	acttctgttt	aaatttccag	ttcaacttgt	1380
aaatgttttt	attgtgcata	aatacatact	aatgttgatg	taaaaaaaaa	aaaaaaaaaa	1440
aaaaaaaaaa	aaaaaaaaaa					1458

<210> 15
 <211> 2005
 <212> DNA
 <213> Homo sapiens.

<400> 15						
ggttgctggc	ccaggtagag	ggcgcgctg	gtccaggtga	gcgggcgctg	ccccgcgacg	60
gcgctgcctg	cccaggcggg	ttcacgtaaa	gacagcgaga	tcctgagggc	cagccgggaa	120
ggaggcgtgg	atatggagct	ggctgctgcc	aagtccgggg	cccgcgcgcg	tgcttagcgc	180
gtcctggggg	ctctgtgggg	acgcgcggcg	cgcgcgggct	cggggaccgc	tagagcccg	240
cgctgcgcgc	atggccctgc	tctcgcgcgc	cgcgctcacc	ctcctgctcc	tcctcatggc	300
cgctgttgct	aggtgccagg	agcaggccca	gaccaccgac	tggagagcca	ccctgaagac	360
catccgggaa	ggcgttcata	agatagacac	gtacctgaac	gccgccttgg	acctcctggg	420
aggcgaggac	ggtctctgcc	agtataaatg	cagtgcagga	tctaagcctt	tcccaggtta	480
tggttataaa	ccctccccac	cgaatggatg	tggtcttcca	ctgtttggtg	ktcatcttaa	540
cattgggtatc	ccttccctga	caaagtgttg	caaccaaacac	gacaggtgct	atgaracctg	600
tggcaaaaag	agaatgactg	gtgatgaaga	attccagtat	tgctctcca	agatctgccg	660
agatgtacag	aaaacactag	gactaactca	gcatgttcag	gcatgtgaaa	caacagtgga	720
gctcttggtt	gacagtgtta	tacatttagg	ttgtaaacca	tatctggaca	gccaacgagc	780
cgcatgcagg	tgtcattatg	aagaaaaaac	tgatctttaa	aggagatgcc	gacagctagt	840
gacagatgaa	gatggaagaa	cataaccttt	gacaaaatac	taatgttttt	acaacataaa	900
actgtcttat	ttttgtgaaa	ggattatatt	gagaccttaa	aataatttat	atctttagat	960
taaaacctca	aagcaaaaaa	agtgagggag	atagtggggg	gagggcacgc	ttgtcttctc	1020
aggtatcttc	cccagcattg	ctcccttact	tagtatgcca	aatgtcttga	ccaatatcaa	1080
aaacaagtgc	ttgttttagc	gagaattttg	aaaagaggaa	tatataactc	aattttcaca	1140
accacattta	ccaaaaaaag	agatcaaata	taaaattcat	cataatgtct	gttcaacatt	1200
atcttatatt	gaaaatgggg	aaattatcac	ttacaagtat	ttgtttacta	tgaattttta	1260
aatacacatt	tatgcctaga	aggaacggac	tttttttttc	tatttttaatt	acacataata	1320
tgtaatataa	gtmcaacata	atatgttggt	tctctgtagc	ccgttgagca	tatgagtaag	1380
tcacatttct	attaggacta	cttmcaagga	caaggtttcc	atttttccag	ttgtaaaatt	1440
ggaaccatca	gctgataacc	tcgtaggggg	caaccccgag	atagctaaag	gttatgtaat	1500
atgcctagaa	ggtgatgtga	atgcgattca	gaagcatagc	cactcccat	ttatgagcta	1560
ctcacatgac	aaatgtcatc	ttttgtcata	acctttgcca	agttagagaa	aagatggatt	1620
taatgagata	aatgaaaaga	tatttamcct	aatatatcaa	ggcactat	gtctgttatgc	1680

tttgttattt	atttcccagc	acttggttcct	tattgtagat	tttttaaaga	ctgtaacctt	1740
ttactaactg	tgggtcttact	aaaatttgtg	cttgatactg	cttttcaaaa	agcctttaat	1800
tagagccaaa	aggatggaaa	aggcaagata	taaatgcctt	ttatagatct	cttattttaca	1860
ttgaaaatta	ttaccatatg	tttagagcaa	atccaagaaa	acttcaacag	cttctgaaga	1920
tgtctatgaa	tgttgaaaaac	ttttcaatst	cttggratgc	tcakttaatt	cgcagaccgg	1980
cttaacggat	taaacgcccc	cccc				2005

<210> 16
 <211> 943
 <212> DNA
 <213> Homo sapiens

<400> 16						
ggcacgagct	ccgccccggc	ccgagggggt	ctccccggag	gctcagcccc	ctctgctccc	60
catgggcaac	tgccaggcag	ggcacaacct	gcacctgtgt	ctggccccacc	acccacctct	120
ggctctgtgcc	actttgatcc	tgtctgtcct	tggcctctct	ggcctggggc	ttggcagctt	180
cctcctcacc	cacaggactg	gcctgcgcac	cctgacatcc	cccaggactg	ggctctcttt	240
ttgagatctt	ttggccagct	gacctgtgt	cccaggaatg	ggacagtcac	agggaagtgg	300
cgaggggtctc	acgtcgtggg	cttgctgacc	accttgaact	tcggagacgg	tccagacagg	360
aacaagaccc	ggacattcca	ggccacagtc	ctgggaagtc	agatgggatt	gaaaggatct	420
tctgcaggac	aactggctct	tatcacagcc	agggtgacca	cagaaaggac	tgcaggaaacc	480
tgcttatatt	ttagtgtgt	tccaggaatc	ctaccctcca	gccagccacc	catatcctgc	540
tcagaggagg	gggctggaaa	tgccaccctg	agccctagaa	tgggtgagga	atgtgttagt	600
gtctggagcc	atgaaggcct	tgtgctgacc	aaactgtctca	cctcggagga	gctggctctg	660
tgtggctcca	ggctgtgggt	cttgggctcc	tctctgtctc	tcttctgtgg	ccttctctgc	720
tgtgtcaactg	ctatgtgctt	ccacccgcgc	cgggagtcct	actggtctag	aacccggctc	780
tgagggcact	ggcctagtct	ccgactgtgt	tctcaggtgt	gaatcaactt	cttgggcctt	840
ggctctgagt	tggaaaagggt	tttagaaaaa	gtgaagagct	ggaatgtggg	ggaaaaataaa	900
aagctttttt	gccccaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaa		943

<210> 17
 <211> 1503
 <212> DNA
 <213> Homo sapiens

<400> 17						
caggttcctc	tcagtamarc	ctcarsccga	ggttcccttc	ctcttgcatc	catgtgtgtg	60
tttcaraggc	ggccatcctt	ccctacttcc	agatccttgt	agggcagttg	gtggaggggtg	120
ggaggcacc	cggtgttgcc	tccatgaagc	cctgtgcccag	tactggggct	gcaaggctga	180
ggaaattgtg	tccgtgtcag	aaagctcctc	agctcagagg	tgctgggtacc	tcctgcgtgg	240
taggaaggca	gggggaagag	gccctgtctc	tctgtttctc	tttgccctta	tgagacttga	300
gagtctgtgt	catctgtgcc	ttgcatgtct	ttttttcaga	ctccctgcga	caaggactgt	360
gtactgcatg	aatgaggtcg	agatagttag	tgttgctctg	ggaatcctga	ttgagagccg	420
yaaacaggam	aaggcctgcg	agcagccggc	cctggcgggg	gctgataacc	cagagcactc	480
ccctccctgc	tccgtgtcgc	ctcacacaag	ttctgggagc	agcagtggag	aagaggacag	540
tgggaaacag	gcaactgrtc	caggcctcag	cccttcccag	aggccggggg	gttccagctc	600
tgctgttagc	aggagccctg	aggaggagga	ggaagaggat	gtgctgaaat	acgtccggga	660
gatctttttc	agctagggca	taaactgtgc	actgaactgt	ctgccgagag	cagctggagg	720
acagctgagc	ttccactggg	gctgctgggc	cgcccgccctg	tgggaatggg	gctctctgtg	780
ctcctacctt	tgtgccttct	tgggcctggc	agattcacct	caggccagaa	gccccctggac	840
actccgggcc	ttggggctgc	cgttctgagt	gtgcggaagg	caggactcaa	aatgagatcc	900
catttgactc	cctctgtatg	tactgtgccc	tctcctggct	cttgaggctc	tggagtccca	960
attgtctgtg	ttagttagtg	accaggttcc	agggaaaatg	atgtcatgtg	gtgggtccaac	1020
ttactggaac	caaagagaca	gtactttgca	aagaaaagga	tactgcccag	gtgacttgga	1080
attgtacag	tttagtccgc	atgatctctc	ctgaaggagg	aagcctgttt	caaaaatagt	1140
ttccatcatg	agtctatcaa	tgagctccca	cctctccagc	cagcctagaa	agcaaacagag	1200

7

ctgccccacag	ttctctgccc	tgtctgggag	gttgaggcca	cagtgtatag	actggtaagc	1260
cagacaggcc	tcctcccga	agctgctacc	ttgctttcac	ctgtaccttg	gtccccgggc	1320
agctagctat	aaagcaagag	ggacaggagc	ccagaagaga	cactgaggac	aagagatcac	1380
accagagtac	atgtctctgc	ctctgttttc	agtgtggctt	tggaacaggaa	tatatgaata	1440
aatcactgcc	atacagggtt	tccaatacac	aagtgtctaga	aaatacacac	aattccccaa	1500
tga						1503

<210> 18

<211> 1512

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (207)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (209)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (521)

<223> n equals a,t,g, or c

<400> 18

gcagagcccc	tgggtgtgag	aagctcgtct	cccgtggggt	gcattgggtc	tgccctatct	60
ctgctccag	cacccagggc	ggccgcagat	ggcagtgctc	ctggggacag	cagctgcgaa	120
tgagtccacg	ggccaacgct	gagctgctca	ggctgaggcg	gtgtgctcag	cacagagccc	180
ccggaactgg	catctgcagg	gcgtganena	aggccgcgcg	gatgccgcac	ttcctggact	240
ggttcgtgcc	ggctctacttg	gtcatctcgg	tcctcattct	gggtgggcttc	ggcgccctga	300
tctactactt	cgagccgggc	ctgcaggagg	cgcacaagtg	gcgcatgcag	cgccccctgg	360
tggaccgcsa	cctccgcaag	acgctaattg	tgccgcgaaa	cctggcccttc	ggcgcccgcc	420
aggtctgagc	cgacttgcaa	aggggatagg	csggcgccac	cgggcgccct	ccccagccc	480
gccccgcccc	cccagccccg	agacccccaa	ggcagaggga	ngccggcctg	ttggccctcc	540
acgctatccc	tctgcagcct	gggcccctcc	gacagaggcc	ccaggtgcgc	tgscagtgra	600
gggtggggac	ttaggtgcct	ggctggccca	gggcttgctc	tcctgtgtaa	gccgactcac	660
ccagagccca	ccctcccaag	ctcaggggca	tcctccgctg	ggccccagtg	cctttgcrct	720
gcgcagcact	ctgccctcca	ctggactcag	gcattgtctat	ggctgcctgt	cctgaggctc	780
cggagccctc	atttcttcgt	gaagtcccca	gctcccctgc	ctccactcaa	tgccaccggc	840
cctgcaactt	taggcaggtc	gaagccaacc	caaggaaaaga	aoctaagaac	ctcgtttgga	900
gggatgtcag	cttggggccag	mccagccgca	cccccggggg	ctcaggcttg	gaactgggtga	960
gggtgtgtgg	tgggggtatg	cagagggata	agaccgtggt	agaggagagg	gttgggtgagg	1020
agagagagag	agagagagag	agagagagtc	tggggggagc	gggcaagcat	ggggagatga	1080
gatgtgtata	tgtgagagag	agtgtggggg	ccccaggcag	ggcaggaggt	ggtggaaacg	1140
gggtgaactc	cgtgggctgt	gtgaggactg	tccatagtgg	gtccmaaccc	cctccctctg	1200
ctggagtttc	ctagcccttc	cccctcccya	agactgwggc	agcaggcagg	agcccctgcc	1260
ctccctccct	gtcctgtgcc	acacttcttg	ggccaaaccc	agcccccttg	agccaggccc	1320
tgccagactc	caagcccacc	ctagaacctt	cctcctgtgt	ggagactctg	ttgccccact	1380
ttggacacag	attggcaacc	tgccctcacm	ckccccctw	cgctggggct	tccatcttaa	1440
tttattctca	ataataaaga	cttcatgatg	amaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1500
aaaaaaaaaa	aa					1512

<210> 19

<211> 1655
<212> DNA
<213> Homo sapiens

<400> 19
ccacgcgtcc gggcaaagaa ttaaacctgg tgtttggact tcaacttagc atggctagaa 60
ttggaagtac agtaaacadg aacctcatgg gatggctgta ttctaagatt gaagctttgt 120
taggttctgc tgggtcacaca accctcggga tcacacttat gattgggggt ataactgtga 180
ttctttcact aatctgtgcc ttggctcttg cctacttgga tcagagagca gagagaatcc 240
ttcataaaga acaaggaaaa acaggtgaag ttattaaatt aactgatgta aaggacttct 300
ccttaccctt gtggcttata ttatcatct gtgtctgcta ttatgttgct gtgttccctt 360
ttattggact tgggaaagtt ttctttacag agaaatttgg attttcttcc caggcagcaa 420
gtgcaattaa cagtgttgta tatgtcatat cagctcccat gtccccggtg tttgggctcc 480
tggtggataa aacagggaag aacatcatct gggttctttg cgcatagcag ccactcttgt 540
gtcccacatg atgctggcct ttacgatgtg gaaccttgg attgctatgt gtcttctggg 600
actctcctac tcattgcttg cctgtgcatt gtggccaatg gtggcatttg tagttcctga 660
acatcagctg ggaactgcat atggcttcat gcagtccatt cagaatcttg ggttggccat 720
catttccatc attgctggta tgatactgga ttctcggggg tatttgtttt tggaaagtgt 780
cttcattgcc tgtgtttctt tgtcactttt atctgtggtc ttactctatt ggtgaatcgt 840
gcccaggggtg ggaacctaaa ttattctgca agacaaagga agaaataaaa tttcccatc 900
tgaatgagaa gttaaaatga atgtgtcaga gaatgggctt aacacatcgt tggtttgaaa 960
acttccatct taaaaattta gaggtttagt attagaaaaa ataactggact ggaaagttaa 1020
atattatacc aaatatacct atttcaaagt gtatttgtga ggctgtttt agcctgtgtc 1080
ttttgtattg tgtgttgcta aagaattcta ctttttagtag gctaataaac aatgaaaggg 1140
ttagaaaatt gctgtggaac atccaggtga acttcaggaa agacagtga aaatggaaaa 1200
cggttgagct tctgttgaga taatcttcat taggtatata tcttagggat acagcctttt 1260
ctttatctta tagcaggaaa aaaaaacttt taggggaaat agaagggtg cgttacacaa 1320
aataaacaat ggcattgtca taggccttcc ttttactagt agggcataat gctagggaat 1380
atgtgaagat gtttttttga agtctcttcc tgatcacgaa caatagcttg cgctctactc 1440
tgtagttatg tggattgccg agcaatgacc cttttcaatt tcttatttct gtgttactga 1500
ggaccctaata cacttaggga tgtaatttta tagtataaac tttctgtaca gtttttctta 1560
tagtctaata agtaaaaagt gtocctcaaa ttatgataat tgcctatgta catggataaa 1620
ttaaaacact gcacacggaa aaaaaaaaaa aaaaa 1655

<210> 20
<211> 2525
<212> DNA
<213> Homo sapiens

<220>
<221> SITE
<222> (5)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (10)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (1354)
<223> n equals a,t,g, or c

<400> 20
tgacnctatn gtaaggtacg cctgcaggta ccgggtccgga attcccggtg cgacccacgc 60
gtccgggtctg ccaacaaggt cgttcatgaa agtgtttttc tctttaaggt aattaaaaaa 120

cagtggaatg	gaaaaacagt	gctgtagtca	tcctgtaata	tgctccttgt	caacaatgta	180
tacattcctg	ctaggtgcc	tattcattgc	tttaagctca	agtcgcatct	tactagtga	240
gtattctgcc	aatgaaggta	agttaagact	tggtatatgc	atggagcact	tccatcta	300
cacacatctc	tctcttgcc	ttggttctgt	tatatataac	atggaaataa	taatgccttt	360
tgcttcattg	gagtgataaa	gcatttttaa	atttgattat	ttacacctgc	attcctcaac	420
aagaaaaaat	gtttgataat	ggatgaaatg	tgagtcaatc	agatacaaaa	atcaaacctt	480
ttggtgaaga	accagtcgta	acatttgact	gttaattcaa	tcaacagggtg	tttctggacc	540
tatagcaaaa	tgtgtaattg	cgcccttattt	tgaagtagaa	ggatatattt	gtttgggtcac	600
ttggcatttg	tgaggctact	actattgtaa	ttattgtatc	aatggtaagg	tgctcagcatt	660
atattgtgcg	gtcatattgt	atcaacagta	taaattataa	gctttgataa	gtatgtattt	720
aagaaatctt	tttttatgta	gggatttaag	caaacacttt	aattccacca	aactgtattg	780
agtaacttct	actagttatt	gagtgaagg	gtgggttgcc	cctccacatc	tggtgggtgtt	840
tcctggttag	tggaacgaga	gacttgga	agaaaggac	atagacaaag	tatagagaaa	900
gaaaaaagg	ggcccagg	accggcgctc	agcacacgga	ggatctctgc	cagcctctga	960
gttccmttag	tatttattga	tcattattgg	gtgtttctcg	gagagtggga	tggtggcagga	1020
tcataggata	gtagtggaga	gaggttcaac	aggtaa	gtgaacaaag	gtctttgcat	1080
cataracaak	gtaaagratt	aagtgtcttg	cttttagata	tgcatacaca	taaacactct	1140
aatgctttac	aaagcagtat	tgctgcccgc	akgtcccacc	tccagcccta	aggcgggttt	1200
ycctatctc	agtagatgga	gcatacaatc	gggttttata	ccgagacatt	ccattgcccc	1260
gggacrggca	ggagacagat	gccttcctct	tgctcaact	gcaagaggcr	ttccttcctc	1320
ttttactaat	cctcctcagc	acagaccctt	tacngggtgt	cgggctgggg	gacggctcagg	1380
tccttccctt	cccacgaggc	catatttcag	actatcacat	ggggagaaac	cttggaacaat	1440
acctggcttt	cctaggcaga	ggtcscctgcg	gcyttccrca	gtgttttgtg	tccttgshta	1500
cttgagatta	gggagtgggtg	atgactctta	asgagcatgc	tgcttcaag	catctgttta	1560
acaaagcaca	tcctgcaccg	cccttaatcc	atttaaccct	gagtkgacac	agcacatgtt	1620
tcagagagca	crgggttggg	ggtaagggtta	cagattgcag	aacaaaatgg	agtctcctat	1680
gtctacttct	ttctamacag	acacagtaac	aatctgatct	ctcttttccc	cacaattgag	1740
gacacatata	atcatgatata	gacctttaat	ggctactac	ttggagagtc	agatgtgtac	1800
ccaagtctct	actgcagttta	acatttacct	gccaggcact	aggctaagta	ttagcagcag	1860
gttcaaagtg	cataagatat	agaccttgct	ctcaagactt	agtttattag	gagagacatg	1920
aatgtaaaca	catcatgaaa	atccattata	ataactgcaa	taattgatat	atcctgaaga	1980
tcagagagatt	gtctagagga	taaagttata	tattctgttt	ggtaggggat	gatgtggagt	2040
tcaaattggat	cagagaacac	ttcgctgatt	agaagtcagt	tgatccacta	gaagtcaagg	2100
tgaacaagg	gattcaaaaac	agaggcaaca	gcctgtaaaa	gggaacagag	gcataaaaaa	2160
gcaggatatg	ttgtgagaac	atgtagtttg	aaattaccaa	gcaaaaagtt	taaggactcg	2220
tagccaggca	cagtggctca	tgctgtaatc	ccagcacttt	aggaggccaa	ggccggcgga	2280
tcacttgagg	tcaggaattt	gagaccaggc	tggccaacat	ggtgaaaccc	atctctacta	2340
gaaatacaaa	aaattagctg	gggttggctg	cgtgtgcctg	taatcccagc	tatgaggagag	2400
actgaagcag	gagaatgaac	ccgggaggca	gagattgcag	tgagccgaga	tcactgccact	2460
gcactccagc	ctgggcaaca	gagcaaaact	gtctcaagaa	aaaaaaaaaa	aaaaagggcg	2520
gccgc						2525

<210> 21
 <211> 1396
 <212> DNA
 <213> Homo sapiens

<400> 21						
aagtctcgtgta	tcgcgcccgg	gaggcgccgg	agcccagcgg	ctggcgccag	atccaggctc	60
ctggaagaac	catgtccggc	agctactggg	catgccaggc	acacactgct	gccaaagagg	120
agctgctgtt	tgaattatct	gtgaatgttg	ggaagaggaa	tgccagagct	gccggctgaa	180
aattaccaca	ccaagagaaa	tctgcaggat	ggacttctg	gtcctcttct	gttctacct	240
ggcttcgggtg	ctgatgggtc	ttgttcttat	ctgcgtctgc	tcgaaaaccc	atagcttgaa	300
aggcctggcc	aggggaggag	cacagatatt	ttcctgtata	attccagaat	gtcttcagag	360
agccrtgcat	ggattgcttc	attacctttt	ccatacgaga	aaccacacct	tcattgtctc	420
gcacctgggtc	ttgcaaggga	tggtttatac	tgagtaacac	tggaagatat	ttggctactg	480
tcaggagctg	gagttgtcct	tgcattacct	tcttctgccc	tatctgctgc	taggtgtaaa	540

10

cctgtttttt	ttcacctga	cttgtggaac	caatcctggc	attataacaa	aagcaaatga	600
attatttttt	cttcatgttt	atgaatttga	tgaagtgatg	tttccaaaga	acgtgaggtg	660
ctctacttgt	gattttaagga	aaccagctcg	atccaagcac	tgcatgtgtg	gtaactgggtg	720
tgtgcaccgt	ttcgaccatc	actgtgtttg	ggtgaacaac	tgcatcgggg	cctggaacat	780
caggtaactc	ctcatctacg	tcttgacctt	gacggcctcg	gctgccacog	tcgccattgt	840
gagcaccact	tttctgggcc	acttggtggt	gatgtcagat	ttataccagg	agacttacat	900
cgatgacctt	ggacacctcc	atgttatgga	cacggctctt	cttattcagt	acctgttcct	960
gactttttcca	cggattgtct	tcattgctggg	ctttgtcgtg	gttctgagct	tcctcctggg	1020
tggctacctg	tgttttgccc	tgtatctggc	ggccaccaac	cagactacta	acgagtggta	1080
cagaggtgac	tgggcctggg	gccagcgttg	tccccttggt	gcctggcctc	cgtcagcaga	1140
gccccaaagtc	caccggaaca	ttcactccca	tgggcttcgg	agcaaccttc	aagagatcct	1200
tctacctgcc	tttccatgtc	atgagaggaa	gaaacaagaa	tgacaagtgt	atgactgcct	1260
ttgagctgta	gttcccgttt	atttacacat	gtggatcctc	gttttccaaa	aaaaaaaaaa	1320
aaaaaaaaaa	aaaaaaaaaa	aaaactcgag	ggggggcccg	gtacccaatt	cgccttgag	1380
ttcaagtaga	catcaa					1396

<210> 22

<211> 1069

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (508)

<223> n equals a,t,g, or c

<400> 22

ggcagcagca	cagcctcagg	cctgccccca	gacctgcaga	atcagaaact	ctggggtgag	60
gcctgggttat	ctgctgtaac	agaccttcca	gtgggttctg	atgccctcta	gagcaggaga	120
accactagct	tagaggttgc	agtatgtttg	gcattcttgc	atttgtgtta	gttcagagga	180
atggctgacc	cccatgtctc	atttctaaagc	ttcaggcagc	ttttctcctg	ggcagctgtc	240
attctgttga	ggggaatcct	ggggactgtg	gtctctcctc	cctgtccgtg	tgtccttgat	300
ctggcagctc	accccttcca	tctcccctgt	gaggctccat	gcctagaggt	ggtcttcaaa	360
cagaagaatg	gcaaagataa	ttgtctcgtg	ttttaccctg	acccatttcc	tttaagaggg	420
tcacttcttg	gcccattcat	taaaaaccaa	tgtcatagtt	ctgtgattcc	actatcagac	480
agtgccacgt	ccaaggcgcg	ggctcttnac	ctcccctgaa	gagagactgt	gctgtctgtg	540
cttctctgtg	tctccagctc	cacgtctcca	cggaccacag	cccttgagga	ctccctcggt	600
gtcccagggc	ttctgggtgt	ttcagagacc	tccacactca	acgaccactg	gtgctgcaga	660
aggggccggtg	cttacattcc	aattaacaga	cgtttttccc	atctaattgcc	tcttgccctc	720
tcctaacacc	acctcgggag	tgtttatgtc	tattctaagt	gaatttcaact	gtgtgaaaaa	780
attcacacct	gttgtcccgag	cgatttggga	ggccggggcg	ggtgtatcat	ttgagcccag	840
gagtttgagg	ctagcctggg	caggatgggt	aaaccccgtc	tctataaaga	aattttaaaa	900
attagctggg	catagtggca	cgtgcctgta	gttccatcta	ctggggaggc	tgggggtggga	960
ggatcgcatg	agcccgggag	tttgaggctg	cagttagctg	tgatgcagc	actgcactcc	1020
agtcgggca	acagagcaag	accctgtctc	ttaaaaaaaa	aaaaaaaaaa		1069

<210> 23

<211> 1658

<212> DNA

<213> Homo sapiens

<400> 23

ggcagcagcc	ggcctgccag	agccatgccc	ctgactcctc	agcttcaaaa	tcagggggtct	60
caggacagag	gatgctgggt	gggctcagag	ctcatcaggg	gggctgtgtg	tgagagggga	120
tgccttcttg	atgccctcat	cctcctcggg	gctgggggtc	ccctcaaggc	caccagctc	180
cttctcttgt	ttgctgtctg	tactcctgcc	gcctgtctgc	ttggccctgc	tgtctctctt	240

11

cttggacttc	ttccctccca	ggcagctgt	gtctcccttc	ttgccggacc	actgctctgc	300
caggcaacct	aggggtgtga	ggagagagac	cctcaacaga	agtgcctcag	ggctggggtg	360
ctgggcaagg	agcactgagc	aggagccgt	gggagtagca	actgggactg	tgcttgacat	420
cagcctccct	gcctcctgcc	tctcgctgtg	gccaccaggc	ccctctgggg	gcattctgact	480
tgtctgcccc	tcatcttgca	cctggtttca	gtgactetta	cttcaccatg	tcttgccaat	540
caagcctttc	aagggcagag	atcctacaat	gccctctggg	gccctctggt	tctcctccta	600
cccacctccc	ccaagggaga	gcaaacaaat	catccagagc	cagcctgccc	tgcttcccc	660
aaactcactg	gtgtcttttc	ccttcagcac	gtggttgccg	cagaggaatt	cagtcagggtc	720
ttcctcctgg	tggatcctgt	accagtcctc	gacacacctc	tcaaactctt	caccagcaca	780
tcacacttgt	taatcataat	acctcatatt	ggcaaagccc	cagcacctga	ctcgctccta	840
gaggagctca	gcctaagcct	cgcaaccac	tgcaaggtag	cagtggcacg	gttcacctaa	900
ggaaactgag	gccagagagg	tgaaatgacc	tgaccaaagc	caccccgccc	tgggtggact	960
tcctcagagc	agacccaatc	cccaccagcc	cttcactggg	cacagcaacc	cttccaaggg	1020
ctgaagggcc	tgtacctgct	tcttgaggtc	agccacttct	gcagaagtct	cgttcaacag	1080
ctcatagggg	atgtccatca	ccaccttgac	ccctttgtgt	accaggttgt	gtaatgtctc	1140
aaaggtctct	gacatgccct	ggaagaagcg	accagatatg	gcaggcggag	ctcccttctc	1200
tcctccccc	cctcgtctcc	cagtgggtgg	taagaaccca	gctataagac	caatgctcaa	1260
cgccctctaa	ggatcctcat	cctttttttt	ttgagaagga	gtctcactct	gtcgccagg	1320
ttggagcgct	tcagctcact	gcaacctctg	cctcccaggt	tcaagcgatt	ttcctgcctc	1380
agcctcccaa	gcagctggga	ctacaaaagg	gtgccaccat	acccggctaa	ttttttaga	1440
gttgggggtt	tgtcatgttg	gtcaggctgg	tctcgaactc	ctagcatcaa	gttttccact	1500
cacctcagcc	tcccaaagtg	ctgagattac	aggcgtgagc	caccgcacct	ggcctcatcc	1560
ttgacctgac	cttcctcttc	cctcttttag	gcctgcttcc	cacaaccctc	gcacatatac	1620
cccctgatct	gcctctgcac	acctcatcgc	ttcaaaaa			1658

<210> 24

<211> 1077

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1036)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1038)

<223> n equals a,t,g, or c

<400> 24

ggcacgaggg	gaaagccatg	ctcccaggac	tccttccttg	cagccttaaa	tcggctctgta	60
cggaaaattc	cgcgcttag	aaacccacgc	ttgggtgtaa	cttattattg	ttcttcctga	120
cctacttcct	gtttatcact	tcggggttca	tcattttggc	atttcgggtga	tcgggttgga	180
actattgaag	cccgccttca	ggttcttttc	ccatttttcc	ctttgaaagg	aagacttctg	240
gcttctccta	aatctccgtt	ctctgggtta	ggggagtcca	agcctctgtc	atgaggaacg	300
gaaatgcgag	ggcctcgggt	gttactctaa	aatccgccct	cagcttgcac	gccggaagct	360
gcgattcctg	cagcggaaga	ggcgtgatct	ggccttcgac	tcgctatgtc	cactaacaat	420
atgtcggacc	cacggaggcc	gaacaaagtg	ctgagggtaca	agcccccgcc	gagcgaatgt	480
aacccggcct	tggacgaccc	gacgccggac	tacatgaacc	tgctgggcat	gatcttcagc	540
atgtgcggcc	tcatgcttaa	gctgaagtgg	tgtgcttggg	tcgctgtcta	ctgctccttc	600
atcagctttg	ccaactctcg	gagctcggag	gacacgaagc	aaatgatgag	tagcttcatg	660
ctgtccatct	ctgccgtggg	gatgtcctat	ctgcagaatc	ctcagcccat	gacgccccca	720
tggtgatacc	agcctagaag	ggtcacattt	tggaccctgt	ctatccacta	ggcctgggct	780
ttggctgcta	aacctgctgc	cttcagctgc	catcctggac	ttccctgaat	gaggccgtct	840
cgggtccccc	agctggatag	agggaaacctg	gccctttcct	agggaaacacc	ctaggcttac	900
ccctcctgcc	tcccttcccc	tgccctgctgc	tggggggagat	gctgtccatg	tttctagggg	960

12

tattcatttg	ctttctcgtt	gaaacctgtt	gttaataaag	tttttctactc	tgaaaaaaaa	1020
aaaaaaaaa	aaaaanCncg	agggggggcc	cggaaaccaa	ttcscgggat	agtgagt	1077

<210> 25
 <211> 1205
 <212> DNA
 <213> Homo sapiens

<400> 25

cccacgcgtc	cgcagaggca	gggcaatagt	ggagtctctg	cttggccaag	cagcctagaa	60
ctcaaagtcc	atggcccctt	ctgggcctgg	agaaattgga	tggttatagc	accaggcagc	120
ccttctgggt	gggggacagc	aaatgaggga	cctctctttt	ctctacactc	tcctttggct	180
cccgagatc	tggcaggccc	tggctggagg	cataagatta	gatgaggttg	agctgttgga	240
gaatgaagct	gtgttgggag	aagaaatgag	gttgaccgg	aagatcaacg	aggttgtgct	300
gtcagggat	gaggtgttac	ttgggggcaa	gtgaggctgc	attattagat	aaatgaggtt	360
gtactgtcag	gggatgaagt	gtactttag	tagagatgac	gtcctgctgg	atcagtcggc	420
ttttgtccca	tcagagaaca	cagccacacc	acaggaggaa	ggagagtgtc	cgactcagag	480
gataaatgag	ggtgtcctgc	tggataaatg	agggggcccg	tcaggtgaat	ggagtgtctg	540
tagcaaatga	ggttgtactt	gctggataaa	tgggactggg	gtgctggata	aatggggttg	600
tgctgtcagg	tgaatgcatt	actgctcgtg	ggtgaagggc	atcctgggaa	taatgaggtt	660
gtcctgctgg	atagatgagc	tgccaccacc	aaatggatca	gacctgttcc	atgaaggagg	720
caccatcagc	aacgacgagg	ttatcctgtt	cccactgggg	ctcctggagc	gtcttctggc	780
ccaggggaaa	ctcgggtgtg	gccaccctgg	gttatccaag	tctctctggg	gagcaggggtg	840
gggggctggg	gagggcaggc	agctgcattg	tgcaccgtgg	gacctctcct	tcaccccca	900
tggatgccct	actcctctcc	ctggcaccct	tcagtgggtc	agactgcttc	ggacattctc	960
acccactgc	ctgcttctca	tcctgcctgt	gtcttcttct	tgcccagttt	ggaaaagccc	1020
ctattatgtg	tcagccactc	tgccagctct	tatttaactc	ccctataaca	cagtattact	1080
cctccttgca	catacacact	ttctcttatt	cattcatcca	ttcattcatt	tgacaaacat	1140
ttaagtgtct	agtatgtacc	aaacacatga	ggtacagttt	taaaaagaat	aaaaaaaaa	1200
aaaaa						1205

<210> 26
 <211> 1674
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1663)
 <223> n equals a,t,g, or c

<400> 26

cccagagcagc	tgagtcocctt	ccctgtcttt	cactcttctg	gcacgcgttg	ttttacttct	60
tcgattgaac	cctgccttct	cgacccccc	gggaggccgc	cttcttcagg	cgcctccctt	120
ctctccacga	gtcgcgtctg	acagctgagg	aactggcaag	atcctgtctac	ccagaggggtg	180
aatgggtatc	tttcccggaa	taatccta	ttttctaagg	gtgaagtgtg	caacggcggc	240
cgtgattgta	agcggagtaa	gcaaacacct	ccattgtatt	agtcaccaga	aaagtaccac	300
tgtaagtcac	gagatgtctg	gtctgaattg	gaaacccttt	gtatatggcg	gccttgcctc	360
tatcgtggct	gagtttggga	ctttccctgt	ggaccttacc	aaaacacgac	ttcaggttca	420
aggccaaagc	attgatgccc	gtttcaaaga	gataaaatat	agagggatgt	tccatgcgct	480
gtttcgcatc	tgtaaagagg	aaggtgtatt	ggctctctat	tcagggaattg	ctcctgcgtt	540
gctaagacaa	gcacatcatg	gcaccattaa	aattgggatt	taccaaagct	tgaagcgctt	600
attcgtagaa	cgtttagaag	atgaaactct	tttaattaat	atgatctgtg	gggtagtgtc	660
aggagtgata	tcttccacta	tagccaatcc	caccgatgtt	ctaaagattc	gaatgcaggc	720
tcaagggaagc	ttgttccaag	ggagcatgat	tgggaagctt	atcgatatat	accaacaaga	780
aggcaccagg	ggtctgtgga	gggtgtgtgt	tccaactgct	cagcgtgctg	ccatcggtgt	840

13

aggagtagag	ctaccagtct	atgatattac	taagaagcat	ttaatattgt	caggaatgat	900
ggcgatata	attttaactc	acttcgtttc	cagctttaca	tgtggtttgg	ctggggctct	960
ggcctccaac	ccggttgatg	tggttcgaac	tcgcatgatg	aaccagaggg	caatcgtggg	1020
acatgtggat	ctctataagg	gcactgttga	tggattttta	aagatgtgga	aacatgaggg	1080
cttttttgca	ctctataaag	gattttggcc	aaactggcct	cggcttggac	cctggaacat	1140
catttttttt	attacatacg	agcagctaaa	gaggcttcaa	atctaagaac	tgaattatat	1200
gtgagcccg	ccctgccagc	ctttctactc	ctttgccctt	ttcccggtgt	ctaagtattt	1260
ttgacaatgt	tgtaaagtgt	taccaagccg	ttggtctcct	aagggcctcc	tgatggaaga	1320
acagtggggt	ggttcaaagt	tatttctatg	tttgtgttac	catgttaact	tttccccgag	1380
agaaagtgtt	aacattgaga	ctctggcccc	agattggtat	cttctatgaa	gatggatact	1440
gatgggtgac	attgaaaacg	gcctgctttc	caaattgtgt	taaatgtaat	tggttagccc	1500
cagacttggg	ctagagcaga	aggcataggg	cagggtgggt	attgctatat	gtgttacaga	1560
cctcggttct	cattaaagta	tttattggca	gaatcaaaaa	aaaaaaaaaa	aaaaaaaaaa	1620
aaactcgagg	gggggccccg	taccaaatcc	gccctatggt	gantcgaatg	ggct	1674

<210> 27

<211> 1965

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (333)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1961)

<223> n equals a,t,g, or c

<400> 27

ggatcctcgc	ggcggcggcg	gtgcttacag	cctgagaaga	gcgtctcgcc	cgggagcggc	60
ggcgcccatc	gagacccacc	caaggcgcgt	ccccctcggc	ctcccagcgc	tcccaagccg	120
cagcggccgc	gccccctcag	ctagctcgct	cgctcgctct	gcttccctgc	tgccggctgc	180
gcatggcggt	ggcgttggcg	gcgctggcgg	cggtcgagcc	ggcctgcggc	agccgggtacc	240
agcagtttga	gaatgaagaa	gagttctggg	aaactgaaca	ggctgcagggt	gatgtctcctc	300
caaccttacg	cagcatttct	gcagagagcg	cancatattt	tgactacaag	gatgagtcgt	360
ggttttccaa	gcccccatct	tacaatgtag	ctacaacact	gcccagttat	gatgaagcgg	420
agaggaccaa	ggctgaagct	actatccctt	tggttccctg	gagagatgag	gattttgtgg	480
gtcgggatga	ttttgatgat	gctgaccagc	tgaggatagg	aaatgatggg	attttcoatgt	540
taactttttt	catggcattc	ctctttaact	ggattggggt	tttctgtctt	ttttgcctga	600
ccacttcagc	tgcaggaagg	tatggggcca	tttcaggatt	tggctctctt	ctaattaaat	660
ggatcctgat	tgtcagggtt	tccacctatt	tccctggata	ttttgatggg	cagtactggc	720
tctgggtggg	gttctctgtt	ttaggttttc	tcctgtttct	cagaggattt	atcaattatg	780
caaaagtctg	gaagatgccg	gaaactttct	caaactctcc	caggaccaga	gttctcttta	840
tttattaaag	atgttttctg	gcaaaggcct	tcctgcattt	atgaattctc	tctcaagaag	900
caagagaaca	cctgcaggaa	gtgaatcaag	atgcagaaca	cagaggaata	atcacctgct	960
ttaaaaaaat	aaagtactgt	tgaagagatc	atttctctct	atttgttcct	agggtgtaaaa	1020
ttttaatagt	taatgcagaa	ttctgtaatc	attgaatcat	tagtggttaa	tgtttgaaaa	1080
agctcttgca	atcaagtctg	tgatgtatta	ataatgcctt	atatattggt	tgtagtcatt	1140
ttaagtagca	tgagccatgt	ccctgtagtc	ggtagggggc	agtcttgctt	tattcatcct	1200
ccatctcaaa	atgaacttgg	aattaaatat	tgtaaatat	gtataatgct	ggccatttta	1260
aaggggtttt	ctcaaaagtt	aaacttttgt	tatgactgtg	tttttgaca	taatccatat	1320
ttgctgttca	agttaatcta	gaaattttatt	caattctgta	tgaacacctg	gaagcaaaat	1380
catagtgcga	aaatacattt	aaggtgtggg	caaaaaataag	tctttaattg	gtaaaaaata	1440
agcattaatt	ttttatagcc	tgtattcaca	attctgcggg	accttattgt	acctaaagga	1500
ttctaagggt	gttgtcactg	tataaaaacag	aaagcactag	gatacaaatg	aagcttaatt	1560

14

actaaaatgt	aattcttgac	actctttcta	taattagcgt	tcttcacccc	cacccccacc	1620
cccccccc	ttattttcct	tttgtctcct	ggtgattagg	ccaaagtctg	ggagtaagga	1680
gaggattagg	tacttaggag	caaagaaaga	agtagcttgg	aacttttgag	atgatcccta	1740
acatactgta	ctacttgctt	ttacaatgtg	ttagcagaaa	ccagtgggtt	ataatgtaga	1800
atgatgtgct	ttctgcccc	gtggtaattc	atcttggttt	gctatgttaa	aactgtaaat	1860
acaacagaac	attaataaat	atctcttgtg	tagcaccttt	aaaaaaaaaa	aaaaaaaaaa	1920
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	naaaa		1965

<210> 28
 <211> 1863
 <212> DNA
 <213> Homo sapiens

<400> 28

gactaggccg	cgagcttagt	cctgggagcc	gcctccgtcg	ccgccgtcag	agccgcccta	60
tcagattatc	ttaacaagaa	aaccaactgg	aaaaaaaaat	gaaattcctt	atcttcgcct	120
ttttcgggtg	tggtcacctt	ttatccctgt	gctctgggaa	agctatatgc	aagaatggca	180
tctctaagag	gacttttgaa	gaaataaaag	aagaaatagc	cagctgtgga	gatgttgcta	240
aagcaatcat	caacctagct	gtttatggta	aagcccagaa	cagatccctat	gagcgattgg	300
cactttctgt	tgatactgtt	ggacccagac	tgagtggctc	caagaacctt	gaaaaagcca	360
tccaaattat	gtaccaaaac	ctgcagcaag	atgggctgga	gaaagtccac	ctggagccag	420
tgagaatacc	ccactgggag	aggggagaag	aatcagctgt	gatgctggag	ccaagaattc	480
ataagatagc	catcctgggt	cttggcagca	gcattgggac	tcctccagaa	ggcattacag	540
cagaagttct	ggtggtgacc	tctttcgatg	aactgcagag	aagggcctca	gaagcaagag	600
ggaagattgt	tgttataaac	caaccttaca	tcaactactc	aaggacggtg	caataccgaa	660
cgcagggggc	ggtggaagct	gccaaagttg	gggctttggc	atctctcatt	cgatccgtgg	720
cctccttctc	catctacagt	cctcacacag	gtattcagga	ataccaggat	ggcgtgcccc	780
agattccaac	agcctgtatt	acgggtggaag	atgcagaaat	gatgtcaaga	atggcttctc	840
atgggatcaa	aattgtcatt	cagctaaaga	tgggggcaaa	gacctacca	gatactgatt	900
ccttcaacac	tgtagcagag	atcactggga	gcaaatatcc	agaacagggt	gtactgggtca	960
gtggacatct	ggacagctgg	gatgttgggc	aggggtgccat	ggatgatggc	ggtggagcct	1020
ttatatcatg	ggaagcactc	tcacttatta	aagatccttg	gctgcgtcca	aagaggactc	1080
tgcggctgtg	gctctggact	gcagaagaac	aagggtggagt	tggtgccctc	cagtattatc	1140
agttacacaa	ggtaaatatt	tccaactaca	gtctgggtgat	ggagtctgac	gcaggaaacct	1200
tcttaccac	tgggctgcaa	ttcactggca	gtgaaaaggc	cagggccatc	atggaggagg	1260
ttatgagcct	gctgcagccc	ctcaatatca	ctcaggctct	gagccatgga	gaagggacag	1320
acatcaactt	ttggatccaa	gctggagtgc	ctggagccag	tctacttgat	gactatatac	1380
agtatttctt	cttccatcac	tcccacggag	acaccatgac	tgtcatggat	ccaaagcaga	1440
tgaatgttgc	tgctgctgtt	tgggctgttg	tttcttatgt	tgttgacagc	atggaagaaa	1500
tgtgccttag	gtcctagaaa	cagtaagaaa	gaaacgtttt	catgcttctg	gccagggaatc	1560
ctgggtctgc	aacttttgaa	aactcctctt	cacataacaa	tttcatccaa	ttcatcttca	1620
aagcacaact	ctatttcatg	ctttctgtta	ttatctttct	tgatacttct	caaatctctt	1680
gattctagaa	aaaggaatca	ttctcccctc	cctcccacca	catagaatca	acatatggta	1740
gggattacag	tgggggcatt	tctttatatc	acctcttaaa	aacattgttt	ccactttaaa	1800
agtaaact	taataaattt	ttggaagatc	tctgaaaaaa	aaaaaaaaaa	aaagggcggc	1860
cgc						1863

<210> 29
 <211> 1626
 <212> DNA
 <213> Homo sapiens

<400> 29

cccacgcgtc	cggagccggg	agccggctgc	gggggctccg	ggctgtggga	ccgctggggc	60
ccagcgatg	gcgacctgt	ggggaggcct	tcttcgctt	ggctccttgc	tcagcctgtc	120
gtgcttgccg	ctttccgtgc	tgctgctggc	gcactgtcag	acgccgcca	gtgattgcct	180

15

tcatgtttgtg	gagcccatgc	ctgtgcgggg	gcctgatgta	gaagcatact	gtctacgctg	240
tgaatgcaaa	tatgaagaaa	gaagctctgt	cacaatcaag	gttaccatta	taatttatct	300
ctccattttg	ggccttctac	ttctgtacat	ggtatatctt	actctgggtg	agcccatact	360
gaagaggcgc	ctctttggac	atgcacagtt	gatacagagt	gatgatgata	ttggggatca	420
ccagcctttt	gcaaatgcac	acgatgtgct	agcccgctcc	cgcagtcgag	ccaacgtgct	480
gaacaaggta	gaatatgcac	agcagcgctg	gaagcttcaa	gtccaagagc	agcgaagtc	540
tgtctttgac	cggcatgttg	tcctcagcta	attgggaatt	gaattcaagg	tgactagaaa	600
gaaacaggca	gacaactgga	aagaactgac	tgggttttgc	tgggtttcat	tttaatacct	660
tgttgatttc	accaactggt	gctggaaatt	caaaactgga	agcaaaaact	tgcttgattt	720
ttttttcttg	ttaacgtaat	aatagagaca	tttttaaaag	cacacagctc	aaatcagcca	780
ataatctttt	cctattgtga	cttttactaa	taaaaataaa	tctgcctgta	aattatcttg	840
aagtccttta	cctggaacaa	gcactctctt	tttcaccaca	tagttttaac	ttgactttca	900
agataatttt	cagggttttt	gttgttgttg	ttttttgttt	gtttgttttg	gtgggagagg	960
ggagggatgc	ctgggaagtg	gttaacaact	tttttcaagt	cactttacta	aacaaacttt	1020
tgtaaataga	ccttaccttc	tattttcgag	tttcatttat	attttgcagt	gtagccagcc	1080
tcacaaaga	gctgacttac	tcatttgact	tttgactga	ctgtattatc	tggttatctg	1140
ctgtgtctgc	acttcatggt	aaacgggatc	taaaatgcct	ggtggctttt	cacaaaaagc	1200
agattttctt	catgtactgt	gatgtctgat	gcaatgcac	ctagaacaaa	ctggccattt	1260
gctagtttac	tctaaagact	aaacatagtc	ttgggtgtgt	tggtcttact	catcttctag	1320
tacctttaag	gacaaatcct	aaggacttgg	acacttgcaa	taaagaaatt	ttattttaaa	1380
cccaagcctc	cctggattga	taatatatac	acatttgtca	gcatttccgg	tcgtggtgag	1440
aggcagctgt	ttgagctcca	atgtgtgcag	ctttgaacta	gggctggggt	tggtgggtgcc	1500
tcttctgaaa	ggtctaacca	ttattggata	actggctttt	ttcttctct	ttggaatgta	1560
acaataaaaa	taatttttga	aacatcaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1620
aaaaaa						1626

<210> 30
 <211> 605
 <212> DNA
 <213> Homo sapiens

<400> 30	
ccacgcgtcc	gcccacgcgt
gaggagcggc	ctgctggcca
ccaagcgccc	cogacacctc
atcctgtggc	ggcgtttccg
ctgttccctg	ccatcttcat
cccttcagct	gaggactctc
gactggcctg	cctcctccgc
tcctgccagg	gtgggcagac
ggagctctga	gatcacccca
gatcagctca	gacatatctc
aaaaa	

<210> 31
 <211> 931
 <212> DNA
 <213> Homo sapiens

<400> 31	
gagagtgcct	aagcgggggt
ctgtgtaacc	ttgggtggcc
gccatggcct	ctgcaactgc
agcctagccc	tgggacatca
gtggattctc	cctgggcctc
atcttgccaa	gggtccctgt

16

gtccatattt	aacatgtaaa	aatgtccsc	crckcgtccg	cccaaacatg	ttgtacattt	420
caccatggcc	ccctcatcat	agcaataaca	ttcccactgc	caggggttct	tgagccagcc	480
aggccctgcc	agtggggaag	gaggccaagc	agtgcctgcc	tatgaaattt	caacttttcc	540
tttcatacgt	ctttattacc	caagtcttct	cccgtccatt	ccagtcaaat	ctgggctcac	600
tcaccccagc	gagctctcaa	atccctctcc	aactgcctaa	ggccctttgt	gtaagggtgc	660
ttaatactgt	cctttttttt	tttttaacag	tgtttttag	atttcagatg	actatgcaga	720
ggcctggggg	acccctggct	ctgggcccgg	cctggggctc	cgaaattcca	aggcccagac	780
ttgctggggg	tgggggggta	tccagaattg	gttgtaaata	ctttgcatat	tgtctgatta	840
aacacaaaca	gacctcaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	900
aaaaaaaaaa	aaaaaaaaaa	aaagggcggc	c			931

<210> 32
 <211> 1407
 <212> DNA
 <213> Homo sapiens

<400> 32						
gaattcggca	cgagggcagg	ctcagaagac	gatgcggggc	tgtgtgccgg	ccttcttctg	60
gcatgtactc	agcctcagga	gagcttgctg	caccagggcc	gcccaggtct	tcacagcaca	120
actgcctgga	aggcaggttg	cgagaaggag	aggcggatgg	catgagcagc	aagggggacc	180
gatgtctgtc	agctcacacc	actccagaac	ctgacaaggc	accagcagga	ccccttgcca	240
ggagcatgtc	tgtgcagcag	tgtttttgcc	cctgcacatt	ccagaagccc	tcatgggaag	300
ggatgcagcc	aggcagactc	ctgccagatg	gggcaggtag	tttattcaaa	gagaactctg	360
tatcccatag	gcccaggctc	tcctttcgtc	tggcgtgggc	tttgcctggc	cagtgtgtgc	420
tcctggctca	gcagaaacat	ccatttgagt	tggcatccct	gtagggatcc	cagagcgttg	480
taagccttct	tgtgattggg	agggatggct	gtgggggtgc	ttccaggagg	gggccaccat	540
tgccgcatct	acttctagac	tcccaaagga	gcccaggctc	aggcaggcct	ggcccagagt	600
cacgctggca	accacgagtt	tgggaagcag	tcgtattctc	tctctctctc	tctctctctc	660
agtatccatg	acaggtatga	aacatattgt	ctctttataa	atgtcatttt	acaaattatg	720
tgattatctg	gaagctctaa	gatgagagca	aatgcctgat	cactctggcc	aatgtcaga	780
tactaaagcc	cattcttggc	cgggcatgtt	ggctcccgcc	kgtaatccca	gcactttggg	840
aagcccaagt	gggtgaatca	cctgaggtca	ggagttcaag	accagcctga	ccaacatggg	900
gataccccgt	ctctactaaa	aatacaagcc	gggcgtggtg	gcgcagtcct	gtaatccag	960
ctactcagga	ggctgaggca	ggaaaatcac	ttgaactcgg	gaggcagagg	ttgcagtggg	1020
ctgagatcgc	gccattgcac	tccagcctgg	gtgacagagc	aagactctgt	ctcataaata	1080
aatacaaagc	ccattcttcc	agagtcttgt	gccttaaata	aaacacacct	ctctgctgtg	1140
ggaagactgt	gcaatggcac	agccgcagag	cttggtttgg	gaggttgaag	tgctctgggg	1200
agaattcgta	gatcatcctc	agaaaagcct	tgccttggtg	ttctaccaga	aaaacgtctc	1260
ccaatcacc	aggaaagctg	tccacagtag	tccccctta	tccacgggtg	cactttccat	1320
gggttcagtt	atctgcggtc	aaccacggtc	tgacaatatt	aaatggaaaa	ttcttcaaac	1380
agttaaaaaa	aaaaaaaaaa	aactcga				1407

<210> 33
 <211> 1526
 <212> DNA
 <213> Homo sapiens

<400> 33						
ggcacgagaa	aaaacottca	ggcggcccat	gggtatgccc	aagaggacag	agaacgaatg	60
cacagaaata	ttgtcagcct	tgcacagaat	ctcctgaact	ttatgattgg	ctctatcttg	120
gatttatggc	aatgcttcct	ctggttttac	attggttctt	cattgaaatg	tactcgggga	180
aaaagagttc	cagcgcactt	ttccaacaca	tactgcatt	atttgaatgc	agcatggcag	240
ctattatcac	cttactttgt	agtgatccag	ttgggtttct	ttatattcgt	tcattgtcgag	300
tattgatgct	ttctgactgg	tacacgatgc	tttacaaccc	aagtccagat	tacgttacca	360
cagtacactg	tactcatgaa	gccgtctacc	cactatatac	cattgtattt	atctattacg	420
cattctgctt	ggtattaatg	atgctgctcc	gacctcttct	ggtgaagaag	attgcatgtg	480

17

ggtttagggaa	atctgatcga	tttaaaagta	tttatgctgc	actttacttc	ttcccaattt	540
taaccgtgct	tcaggcagtt	ggtggaggcc	ttttatatta	cgcttccca	tacattatat	600
tagtggtatc	tttggttact	ctggctgtgt	acatgtctgc	ttctgaaata	gagaactgct	660
atgatcttct	ggtcagaaaag	aaaagactta	ttgttctctt	cagccactgg	ttacttcatg	720
cctatggaat	aatctccatt	tccagagtgg	ataaacttga	gcaagatttg	cccccttgg	780
ctttgggtacc	tacaccagcc	cttttttact	tgttcactgc	aaaatttacc	gaaccttcaa	840
ggatactctc	agaaggagcc	aatggacact	gagtgtagac	atgtgaaatg	ccaaaaacct	900
gagaagtgc	cctaataaaa	aagtaaatca	atcttaacag	tgatgagaa	ctattctatc	960
atatatggga	acaagattgt	cagtatatct	taatgtttgg	gtttgtcttt	gttttgttta	1020
tgggttagact	tacagacttg	gaaaatgcaa	aactctgtaa	tactctgtta	cacagggtaa	1080
tattatctgc	tacactggaa	ggccgctagg	aagcccttgc	ttctctcaac	agttcagctg	1140
ttcttttaggg	caaaatcatg	tttctgtgta	cctagcaatg	tggtcccat	ttattaagaa	1200
aagctttaac	acgtgtaatc	tgcagtcctt	aacagtggcg	taattgtacg	tacctgttgt	1260
gtttcagttt	gtttttcacc	tataatgaat	tgtaaaaaa	aacatacttg	tggggtctga	1320
tagcaaacat	agaaatgatg	tatattgttt	tttgttatct	atttattttc	atcaatacag	1380
tattttgatg	tattgcaaaa	atagataata	atttatataa	cagggtttct	gtttatagat	1440
tgggtcaaga	tttgtttgga	ttattgttcc	tgtaaagaaa	acaataataa	aaagcttacc	1500
tacataaaaa	aaaaaaaaaa	aaaaaa				1526

<210> 34

<211> 1737

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1674)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1731)

<223> n equals a,t,g, or c

<400> 34

gtcgaccac	gcgtccgcc	acgcgtccgc	ccacgcgtcc	ggtttataaa	cagaagttaa	60
aaacttgtaag	cttaagcttc	cggtttataaa	cagaagttta	aaattatag	tcctgtttaa	120
cattcagctc	tgtttaactca	ctcatctttt	tgtgttttta	cactttgtca	agatttcttt	180
acatattcat	caatgtctga	agaagttact	tatgcagatc	ttcaattcca	gaactccagt	240
gagatggaaa	aaatcccaga	aattggcaaa	tttggggaaa	aagcacctcc	agctccctct	300
catgtatggc	gtccagcagc	cttgtttctg	actcttctgt	gccttctgtt	gctcattgga	360
ttgggagtct	tggcaagcat	gtttcatgta	actttgaaga	tagaaatgaa	aaaaatgaac	420
aaactacaaa	acatcagtga	agagctccag	agaaatattt	ctctacaact	gatgagtaac	480
atgaatatct	ccaacaagat	caggaacctc	tccaccacac	tgcaacaat	agccaccaa	540
ttatgtcgtg	agctatatag	caaagaacaa	gagcacaaat	gtaagccttg	tccaaggaga	600
tggatttggc	ataaggacag	ctgttatttc	ctaagtgatg	atgtccaaac	atggcaggag	660
agtaaaaatgg	cctgtgctgc	tcagaatgcc	agcctgttga	agataaacaa	caaaaatgca	720
ttggaattta	taaaatccca	gagtagatca	tatgactatt	ggctgggatt	atctcctgaa	780
gaagattcca	ctcgtggtat	gagagtggat	aatataatca	actcctctgc	ctgggttata	840
agaaacgcac	ctgacttaaa	taacatgtat	tgtggatata	taaatagact	atatgttcaa	900
tattatcact	gcacttataa	acaaagaatg	atatgtgaga	agatggccaa	tccagtgcag	960
cttggttcta	catatttttag	ggaggcatga	ggcatcaatc	aaatacattg	aaggagtgtg	1020
kggggtgggg	gttctaggct	ataggtaaat	ttaaatat	tctggttgac	aattagttga	1080
gtttgtctga	agacctggga	ttttatcatg	cagatgaac	atccaggtag	caagcttcag	1140
agagaataga	ctgtgaatgt	taatgccaga	gaggtataat	gaagcatgtc	ccacctccca	1200
ctttccatca	tggcctgaac	cctggaggaa	gaggaagtcc	attcagatag	tgtggggggc	1260
cttcgaattt	tcattttcat	ttacgttctt	ccccttctgg	ccaagatttg	ccagaggcaa	1320

catcaaaaac	cagcaaattt	taattttgtc	ccacagcgtt	gctaggggtg	catgggtccc	1380
catctcgggt	ccatcctata	cttccatggg	actccctatg	gctgaaggcc	ttatgagtca	1440
aaggacttat	agccaattga	ttgttctagg	ccaggtaaga	atggatatgg	acatgcattt	1500
attacctctt	aaaattatta	ttttaagtaa	aagccaataa	acaaaaacga	aaaggcaagt	1560
tacgagactg	acttattttt	aacttctgtg	tgttgagcta	ctgtaagctt	ggcttttgtt	1620
aaagacatac	agcaattagc	tatgcaaaaca	taagcattgt	tctgaaaaaa	aatntataga	1680
tagatatgtt	tatctcccat	aactcataac	tggggagtat	tataccccgg	nggctttt	1737

<210> 35
<211> 2242
<212> DNA
<213> Homo sapiens

<400> 35

tcgacccacg	cgctcgggct	gccatggcgg	cggcgggccc	gctcccgagc	tcctggggccc	60
tcttctcgcc	gctcctcgca	gggcttgcac	tactgggagt	cgggccgggtc	ccagcggggg	120
cgctgcacaa	cgctcacggcc	gagctctttg	gggcccaggc	ctggggcacc	cttgcgggctt	180
tcggggacct	caactccgac	aagcagacgg	atctcttcgt	gctgcgggaa	agaaatgact	240
taatcgtctt	tttggcagac	cagaatgcac	cctattttaa	acccaaaagta	aaggatatctt	300
tcaagaatca	cagtgcattg	ataacaagtg	tagtccctgg	ggattatgat	ggagattctc	360
aaatggatgt	ccttctgaca	tatcttccca	aaaattatgc	caagagtga	ttaggagctg	420
ttatcttctg	gggacaaaat	caaacattag	atcctaacaa	tatgaccata	ctcaatagga	480
cttttcaaga	tgagccacta	attatggatt	tcaatggtga	tctaattcct	gataattttg	540
gtatcacaaa	tgaatccaac	cagccacaga	tactattagg	agggaattta	tcatggcatc	600
cagcattgac	cactacaagt	aaaatgcaaa	ttccacattc	tcattgcattt	attgatctga	660
ctgaagattt	tacagcagat	ttattcctga	cgacattgaa	tgccaccact	agtaccttcc	720
agtttgaaat	atgggaaaat	ttggatggaa	acttytstgw	magtacymta	ttggaaaaac	780
ctcaaaatat	gatgggtggt	ggacagtcag	catttgcaga	ctttgatgga	gatggacaca	840
tggatcatatt	actgccaggc	tgtgaagata	aaaattgcca	aaagagtacc	atctacttag	900
tgagatctgg	gatgaagcag	tgggttccag	tcctacaaga	tttcagcaat	aagggcacac	960
tctggggctt	tgtgccattt	gtggatgaac	agcaaccaac	tgaataacca	attccaatta	1020
cccttcatat	tgagagactac	aatatggatg	gctatccaga	cgctctggtc	atactaaaga	1080
acacatctgg	aagcaaccag	caggccctttt	tactggagaa	cgctccctgt	aataatgcaa	1140
gctgtgaaga	ggcgcgtcga	atgtttaaag	tctactggga	gctgacagac	ctaaatcaaa	1200
ttaaggatgc	catggttgcc	accttctttg	acatttacga	agatggaatc	ttggacattg	1260
tagtgctaag	taaaggatat	acaaagaatg	attttgccat	tcatacacta	aaaaataact	1320
ttgaagcaga	tgcttatttt	gttaaagtta	ttgttcttag	tggtctgtgt	tctaagtact	1380
gtcctcgtaa	gataacaccc	tttggagtga	atcaacctgg	accttatatc	atgtatacaa	1440
ctgtagatgc	aaatgggtat	ctgaaaaatg	gatcagctgg	ccaactcagc	caatccgcac	1500
atttagctct	ccaactacca	tacaacgtgc	ttggtttagg	tcggagcgca	aattttcttg	1560
accatctcta	cgttggtatt	ccccgtccat	ctggagaaaa	atctatacga	aaacaagagt	1620
ggactgcaat	cattccaaat	tcccagctaa	ttgtcattcc	ataccctcac	aatgtccctc	1680
gaagtggag	tgccaaactg	tatcttacac	caagtaatat	tgttctgctt	actgctatag	1740
ctctcatcgg	tgtctgtggt	ttcatcttgg	caataattgg	cattttacat	tggcaggaaa	1800
agaaagcaga	tgatagagaa	aaacgacaag	aagcccaccg	gtttcatttt	gatgctatgt	1860
gacttgccct	taataattaca	taatggaatg	gctgttcact	tgattagtgt	aaacacaaat	1920
tctggcttga	aaaaataggg	gagattaaat	attattttata	aatgatgtat	cccatggtaa	1980
ttattggaaa	gtattcaaat	aaatatgggt	tgaatatgtc	acaaggctct	tttttttaaa	2040
gcactttgta	tataaaaatt	tgggttctct	attctgtagt	gctgtacatt	tttgttcctt	2100
tgtggaatgt	gttgcatgta	ctccagtggt	tgtgtattta	taatcttatt	tgcatcatga	2160
tgatggaaaa	agttgtgtaa	ataaaaaata	ttaaatgagc	aggaaaaaaa	aaaaaaaaaa	2220
aaaaaaaaaa	aagggcggcc	gc				2242

<210> 36
<211> 2235
<212> DNA

<213> Homo sapiens

<400> 36

gtaattcggc	acgaggggttc	caccaacatg	gagctctcgc	agatgtcgsa	gctcatgggg	60
ctgtcgggtg	tgcttgggct	gctggccctg	atggcgacgg	cggcggtasc	gcggggggtgg	120
ctgcgcgcgg	gggaggagag	gagcgccgg	ccgcctgcc	aaaaagcaa	tggatttcca	180
cctgacaaat	cttcgggatc	caagaagcag	aaacaatatc	agcggattcg	gaaggagaag	240
cctcaacaac	acaacttcac	ccaccgcctc	ctggctgcag	ctctgaagag	ccacagcggg	300
aacatatctt	gcatggactt	tagcagcaat	ggcaaatacc	tggtacctg	tgcatgatg	360
cgcaccatcc	gcattctggg	caccaaggac	ttcctgcagc	gagagcaccg	cagcatgaga	420
gccaacgtgg	agctggacca	cgccaccctg	gtgcgcttca	gccctgactg	cagagccttc	480
atcgtctggc	tggtcaacgg	ggacaccctc	cgtgtcttca	agatgaccaa	gcgggaggat	540
gggggctaca	ccttcacagc	cacccagag	gacttcccta	aaaagcacaa	ggcgctgtc	600
atcgacattg	gcattgctaa	cacagggaag	tttatcatga	ctgcctccag	tgacaccact	660
gtcctcatct	ggagcctgaa	gggtcaagtg	ctgtctacca	tcaacaccaa	ccagatgaac	720
aacacacacg	ctgctgtatc	tcctgtggc	agattttag	cctcgtgtgg	cttcacccca	780
gatgtgaagg	tttgggaagt	ctgctttgga	aagaaggggg	agttccagga	ggtggtgcga	840
gccttcgaac	taaaaggcca	ctccgcggct	gtgcactcgt	ttgctttctc	caacgactca	900
cggaggatgg	cttctgtctc	caaggatggt	acatggaaac	tgtgggacac	aratgtggaa	960
tacaagaaga	agcaggaccc	ctacttgctg	aagacaggcc	gctttgaaga	ggcggcgggt	1020
gccgmccgt	gcccctggc	cctctcccc	aacgccagg	tcttgccctt	ggccagtggc	1080
agtagtattc	atctctacaa	tacccggcgg	ggcgagaagg	aggagtgtct	tgagcgggtc	1140
catggcgagt	gtatcgccaa	cttgtccttt	gacatcactg	gccgctttct	ggcctcctgt	1200
ggggaccggg	cgggtcggct	gtttcacaac	actcctggcc	accgagccat	ggtggaggag	1260
atgcaggggc	acctgaagcg	ggcctccaac	gagagcacc	gccagaggct	gcagcagcag	1320
ctgacccagg	cccaagagac	cctgaagagc	ctgggtgccc	tgaagaagtg	actctgggag	1380
ggcccgggcg	agaggattga	ggaggaggga	tctggcctcc	tcatggcact	gctgccatct	1440
ttcctcccag	gtggaagcct	ttcagaagga	gtctcctggg	tttyttactg	gtggccctgc	1500
ttcttcccat	tgaaactact	cttgtctact	taggtctctc	tcttcttgct	ggctgtgact	1560
cctccctgac	tagtggccaa	ggtgcttttc	ttcctcccag	gcccagtggg	tggaatctgt	1620
cccacactgg	cactgaggag	aatggtagag	aggagaggag	agagagagag	aatgtgattt	1680
ttggccttgt	ggcagcacat	cctcacacc	aaagaagttt	gtaaatgttc	cagaacaacc	1740
tagagaacac	ctgagtacta	agcagcagtt	ttgcaaggat	gggagactgg	gatagcttcc	1800
catcacagaa	ctgtgttcca	tcaaaaagac	actaagggat	ttccttctgg	gcctcagttc	1860
tattttgtaag	atggagaata	atcctctctg	tgaactcctt	gcaaagatga	tatgaggcta	1920
agagaatatc	aagtccocag	gtctggaaga	aaagtagaaa	agagtagtac	tattgtccaa	1980
tgatcatgaa	gtggtaaaag	tgggaaccag	tgtgctttga	aaccaaatta	gaaacacatt	2040
ccttgggaag	gcaaaagttt	ctgggacttg	atcatacatt	ttatatggtt	gggacttctc	2100
tcttcgggag	atgatattct	gtttaaggag	acctcttttc	agttcatcaa	gttcatcaga	2160
tattttgagt	ccactctgt	gcccataata	atatgagctg	gggattaaaa	aaaaaaaaaa	2220
aaaaaaaaaa	ctcga					2235

<210> 37

<211> 2971

<212> DNA

<213> Homo sapiens

<400> 37

gacgtgagga	gcgttccatt	tggccagtgg	tgggcggttg	ccacagctgg	tttagggccc	60
cgaccactgg	ggcccttgt	caggaggaga	cagcctcccg	gccgggggag	gacaagtcgc	120
tgccaccttt	ggctgccgac	gtgatccct	gggacggtcc	gtttcctgcc	gtcagctgcc	180
ggccgagttg	ggtctccgtg	gttcaggcgg	gtccccctt	cctgggtctcc	cttctccgc	240
tgggcccgtt	tatcgggagg	agattgtctt	ccagggctag	caattggact	tttgatgatg	300
tttgaccag	cggcaggaat	agcaggcaac	gtgatttcaa	agctgggctc	agcctctgtt	360
tcttctctcg	tgtaatcgca	aaaccattt	tggagcagga	attccaatca	tgtctgtgat	420
ggtggtgaga	aagaagggtg	cacggaaact	ggagaaactc	ccaggcagga	acaccttttg	480
ctgtgatggc	cgcgtcatga	tggcccggca	aaagggcatt	ttctacctga	cccttttctt	540

catcctgggg	acatgtacac	tcttcttcgc	ctttgagtgc	cgctacctgg	ctgttcagct	600
gtctctgccc	atccctgtat	ttgtgccc	gctcttcctt	ttctccatgg	ctacactgtt	660
gaggaccagc	ttcagtgaac	ctggagtgat	tcctcggg	ctaccagatg	aagcagcttt	720
catagaaatg	gagatagaag	ctaccaatgg	tgcggtgccc	cagggccagc	gaccaccgcc	780
tcgtatcaag	aattttccaga	taaacaacca	gattgtgaaa	ctgaaatact	gttacacatg	840
caagatcttc	cggcctcccc	gggcctccca	ttgcagcatc	tgtgacaact	gtgtggagcg	900
cttcgaccat	caactgcccct	gggtggggaa	ttgtgttgga	aagaggaaact	accgctactt	960
ctacctcttc	atcctttctc	tctccctcct	cacaatctat	gtcttcgcct	tcaacatcgt	1020
ctatgtggcc	ctcaaactctt	tgaatttgg	cttcttgagg	acattgaaag	aaactcctgg	1080
aactgttcta	gaagtcctca	tttgcctctt	tacactctgg	tccgtcgtgg	gactgactgg	1140
atttcatact	ttcctcgtgg	ctctcaacca	gacaaccaat	gaagacatca	aaggatcatg	1200
gacagggag	aatcgctcc	agaatcccta	cagccatggc	aatattgtga	agaactgctg	1260
tgaagtgtg	tgtggccct	tgccccccag	tgtgtgggat	cgaaggggta	ttttgccaact	1320
ggaggaaagt	ggaagtgcac	ctcccagtac	tcaagagacc	agtagcagcc	tcttgccaca	1380
gagcccagcc	cccacagaac	acctgaactc	aaatgagatg	cgggaggaca	gcagcactcc	1440
cgaagagatg	ccacctccag	agccccaga	gccaccacag	gaggcagctg	aagctgagaa	1500
gtagccctatc	tatggaagag	acttttgttt	gtgtttaatt	agggctatga	gagatttcag	1560
gtgagaagtt	aaacctgaga	cagagagcaa	gtaagctgtc	ccttttaact	gtttttcttt	1620
ggctcttagt	cacccagttg	cacactggca	ttttcttgc	gcaagctttt	ttaaatttct	1680
gaactcaagg	cagtggcaga	agatgtcagt	cacctctgat	aactggaaaa	atgggtctct	1740
tgggcccctg	caactggtct	ccatggcctc	agccacaggg	tccccttggg	ccccctctct	1800
tccctccaga	tcccagccct	cctgcttggg	gtcactggtc	tcattctggg	gctaaaagtt	1860
tttgagactg	gctcaaactcc	tcccaagctg	ctgcacgtgc	tgagtccaga	ggcagtcaca	1920
gagacctctg	gccaggggat	cctaactggg	ttcttgggg	cttcaggact	gaagaggagg	1980
gagagtgggg	tcagaagatt	ctcctggcca	ccaagtggca	gcattgcccc	caaactcctt	2040
taggaatggg	acagggtacct	tccacttggt	gtatttatta	gtgtagcttc	tcctttgtct	2100
cccattccact	ctgacacctta	agccccactc	ttttccatt	agatatatgt	aagtagttgt	2160
agtagagata	ataattgaca	tttctcgtag	actaccacaga	aactttttta	atacctgtgc	2220
cattctcaat	aagaatttat	gagatgccag	cggcatagcc	cttcacactc	tctgtctcat	2280
ctctcctcct	ttctcattag	ccccctttta	ttgtttttc	cttttgactc	ctgctcccat	2340
taggagcagg	aatggcagta	ataaaagtct	gcactttgg	catttctttt	cctcagagga	2400
agcctgagtg	ctcacttaaa	cactatcccc	tcagactccc	tgtgtgaggc	ctgcagaggc	2460
cctgaatgca	caaatgggaa	accaaggcac	agagaggctc	tcctctctc	tcctctcccc	2520
cgatgtatcc	tcaaaaaaaa	aaaaaatgct	aaccagttct	tccattaaag	ctcggctgag	2580
tgagggaaag	cccagcactg	ctgccctctc	gggttaactca	ccctaaggcc	tcggcccacc	2640
tctggctatg	gtaaccacac	tgggggcttc	ctccaagccc	cgctcttcca	gcactccac	2700
cggcagagtc	ccagagccac	ttcacctgg	gggtgggctg	tggccccag	tcagctctgc	2760
tcaggacctg	ctctatttca	gggaagaaga	tttatgtatt	atatgtggct	atatctccta	2820
gagcacctgt	gttttctctc	ttctaagcca	gggtcctgtc	tggatgactt	atgcgggtgg	2880
ggagtgtaaa	cgggaacttt	tcattctatt	gaaggcgatt	aaactgtgtc	taatgcaaaa	2940
aaaaaaaaa	aaaaaaaaa	aaaaaaaaa	a			2971

<210> 38

<211> 1163

<212> DNA

<213> Homo sapiens

<400> 38

ccacgcgtcc	gccaaagggtg	ctgattaggg	aatgggttatg	gactaggagt	atcagtaaca	60
atgggttagaa	agtggctaac	atgtgttgag	cacctgctgt	gtgcctggcc	ccggctggga	120
gccttcgtgc	ccagagtga	cccgtctgaa	tgcagttctt	tgctcattc	aaactgggga	180
gtggggaggca	gagctgcaca	actcacagg	gccgagctca	agactcactc	ctgggtctgc	240
ctgggctggg	ctgtgcttgt	tgccctctg	gccaacacac	gcgcaccttt	cacctgaaag	300
ccaggatccg	cagaacgttc	cccagaggag	tcattgtttg	gcactatgat	ttgtctcttc	360
ctaaaaaggt	gatagagtta	cactggagag	agcagcatcc	aggtgcagca	gggatgggccc	420
tggggctcac	gggcaggggc	tctgtgtccg	gctggggcct	gggtcctgc	gctgcacctg	480
tgtgtcagaa	gcactcagta	aatctttgct	gatgaaggat	gacaggatat	aggacatgat	540

21

gcttgctgct	gcattgcctg	caatcctgga	tgaatgccca	ggttggtttt	gtccccgctc	600
gggtggatgt	gacgttagct	gtgatgttag	gtccctggct	ttaaaatacg	acggaactgg	660
gaattgaggg	agcagttggg	gcagaaagga	cagccccgca	gaggcctgga	gctgagcagt	720
gcgggcgacc	caggagcagt	gagtgccttc	gtcacagcct	tcacgcgacc	ctgtggctct	780
cataaagggg	atggaatcta	cgaatttagt	tttcccagcc	tccttaaaaa	ctcattcatg	840
ccaggggag	tggctcacac	ctgaaatccc	accactttgg	gaggctgagg	caggctgatt	900
acttgaggtc	aggagtttga	gaccagccta	gccaaacatg	tgaaaccccg	tgtctactca	960
aaagtacaaaa	aaaaaaatta	gtcagacgtg	gtgtcacgca	cctgtaatcc	cagctctttg	1020
ggaggctgag	gcaggagaat	cacttgaacc	caggaggcag	aggttatagt	gagccagtat	1080
tgcgccactg	acctccatct	gggcaataga	gtgagaccct	gtctcaaaaa	aaaaaaaaaa	1140
aaaaaaaaaa	aaaaaaaaaa	aaa				1163

<210> 39

<211> 1932

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1624)

<223> n equals a,t,g, or c

<400> 39

ggcacgagcc	aggccctg	gccgggagct	gaggcggg	cctctgggca	gggcccgggc	60
ggggctgggt	gggcccggcc	tgctgctgcc	gtccatgctg	atgtttgcgg	tgatcggtggc	120
ctccagcggg	ctgctgctca	tgatcgagcg	gggcatcctg	gccgaratga	agccctgccc	180
cctgcacccg	cccggccggg	arggcacagc	ctggcgcggg	aaagccccc	agcctggggg	240
cctgtccctc	agggctgggg	acgcggactt	gcaagtgcgg	caggacgtcc	ggaacaggac	300
cctgcgggag	gtgtgaggac	agccaggcat	gccccgggac	ccctgggact	tgccgggtggg	360
gcagcggcgc	accctgctgc	gccamatcct	cgtaagtgc	cgttaccgct	tcctctactg	420
ctacgtcccc	aaggtggcct	gctctaactg	gaagcgggtg	atgaagggtg	tggcaggcgt	480
cctggacagc	gtggacgtcc	gcctcaagat	ggaccaccgc	agtgcactgg	tgctcctggc	540
cgacctgcgg	cctgaggaga	ttcgctaccg	cctgcagcac	tactttaagt	tcctgtttgt	600
gcgggagccc	ttggaacgcc	tcctctctgc	ctaccgcaac	aagtttggcg	agatccgaga	660
gtaccagcaa	cgctatgggg	ctgagatagt	gaggcggtac	agggctggag	cggggcccag	720
ccctgcaggc	gacgatgtca	cattccccga	gttcctgaga	tacctggtgg	atgaggacct	780
tgagcgcag	aatgagcatt	ggatgcccg	gtaccacctg	tgccagcctt	gtgccgtgca	840
ctatgacttt	gtgggctcct	atgagaggct	ggaggctgat	gcaaatcagg	tgctggagtg	900
ggtacgggca	ccacctcacg	tccgatttcc	agctcgccag	gcctggtacc	ggccaggccag	960
ccccgaaaag	ctgcattacc	acttgctgag	tgcccccccg	gccctgctgc	aggatgtgct	1020
gcctaagtat	atcctggayt	tytccctctt	tgccctacca	ctgcctaagt	tcaccaagga	1080
ggcgtgtcag	cagtgacct	gggtgtgggg	ccagcagctg	gtggggactg	gtttcaacgc	1140
cagctttctg	tgcttctgcc	tgtcattcgg	agaaactctg	gctctggggc	ttggggcttc	1200
tcaggatcct	ggatggcaga	gactgccctc	agaarttcc	tgtccagggg	gggcacccac	1260
agtgactcag	aggacagggc	taggcaggag	acctgtgct	cctcattggg	gggatctctt	1320
ggggggcaga	caccagtttg	ccaatgaagc	aacacatctg	atctaaagac	tggtccaga	1380
ccccgggctg	ccaggattat	gcagtcact	tggtctacct	taatttaacc	tgtggccaaa	1440
ctcagagatg	gtaccagcca	ggggcaagca	tgaccagagc	cagggaccct	gtggctctga	1500
tccccatttt	atccacccca	tgtgcctcag	gactagagtg	agcaatcata	ccttataaat	1560
gactttttgt	cctttctgct	ccagtctcaa	aatttccctac	acctgccagt	tctttacatt	1620
tttnccaagg	aaaggaaaac	ggaagcaggg	ttcttgcctg	gtagctccag	gaccagctc	1680
tgcaggcacc	caaagaccct	ctgtgcccag	cctcttctct	gagttctcgg	aacctcctcc	1740
ctaattctcc	cttccttccc	cacaaggcmt	ttgaggttgt	gactgtggct	ggtatatctg	1800
gctgccattt	ttctgatgca	tttatttaaa	atgtgtactt	tttgatagaa	cccttgtaag	1860
ggctttgttt	tcctaatagc	tgacttttta	ataaagcagt	tttatataaa	aaaaaaaaaa	1920
aaaaaaaaaa	aa					1932

<210> 40
 <211> 881
 <212> DNA
 <213> Homo sapiens

<400> 40
 gaattcggca cgaggggaacc cagaagatgc tgccctctcct gatcatctgt ctccctgcctg 60
 ccattgaagg gaagaactgc ctccgctgct gcccagaact gtctgccttg atagactatg 120
 acctgcagat cctctgggtg accccagggc caccacaga actttctcaa agtattcact 180
 ccttggtcct agaggataat aattttctca aacctggta ccttgatcgt gaccatttgg 240
 aagaagaaac agccaaattc ttactcaag tacaccaagc cattaacacg ttacgagatg 300
 ataaaacagt acttctggaa gagatctaca cgcacaagaa tctctttact gagaggctga 360
 ataagatata tgatgggctg aaggagaagg gagccccacc cytctccatg aatgccttcc 420
 cggctccatc tcctacttgc accccagaac cccttggctc tgtctgcctc cccagcacct 480
 cagtttctct accttctcac ctccctggca gcctgcaatg agtccctgtc caggaaccgg 540
 cggacctccc tgtgggctgt gagtctcagc agtgcctctac tcctggccat agctggagat 600
 gtttctttta ctggcaaagg aagaaggagg cagtaaagga acagggcagc ccgcatgtct 660
 tccagaagtg aacagaggcc gcagctacca ccgtcacaaa gttcactcat ctctgggtcc 720
 cggtgacccc atcccccat accctccatc ctgggtcctg gggccccaaa gctctgaggc 780
 ctaggagact gcgctgtctc gtggtttgcc tactcttaca cctttgtaaa gagtctcttc 840
 attaaaaccc ctcttcataa aaaaaaaaaa aaaaaactcg a 881

<210> 41
 <211> 1932
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (2)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1022)
 <223> n equals a,t,g, or c

<400> 41
 cncggcgcg ctcgggtcat gccccgggc ggggggcaca caggccggcc ggcagccgct 60
 gggaaatagg cccccgggg cggtgggggc gggggggcca tggcgcgagg accccgggcg 120
 cgggcccgt ccggggagga gttctccttc gtcagccgc tggtgaaata cctgctcttc 180
 ttcttcaaca tgctcttctg ggtgatttcc atgggtatgg tggctgtggg tgtctacgct 240
 cggctaataga agcatgcaga agcagcccta gcctgcctgg cagtggaccc tgccatcctg 300
 ctgatcgtgg tgggtgtcct catgttcctg ctacacctct gtggctgcat tgggtccctc 360
 cgcgagaaca tctgcctcct gcagacgttc tccctctgcc tcaccgctgt gttcctgctg 420
 cagctggccg ctgggatcct gggcttcgtc ttctcagaca aggctcgagg gaaagtgagt 480
 gagatcatca acaatgccat tgtgcactac cgagatgact tggatctgca gaacctcatt 540
 gattttggcc agaaaaagt tagctgctgt ggagggattt cctacaagga ctggtctcag 600
 aacatgtatt tcaactgctc agaagacaac ccagtcgag agcgtgctc tgtgccttac 660
 tctgttggct tgcctactcc tgaccaggca gtgatcaaca ctatgtgtgg ccaaggatg 720
 caggcctttg actacttggga agctagcaaa gtcacttaca ccaatggctg tattgacaag 780
 ttggtcaact ggatacacag caacctattc ttacttgggt gtgtggctct aggcctggcc 840
 atccccagc tgggtgggaat tctgtgttcc cagatcctag tgaatcagat caaagatcag 900
 atcaagctac agctctacaa ccagcagcac cgggtgacc catggtagt agaatccatc 960
 ctgcacctcc tcaccatgga aactggcaag cctcataaac gaacagcagt ggggtgctgaa 1020
 ancagcacca aatggagatt tggattccag cccccagtg acagcccagt ggaagaagc 1080

23

aaactccaga	tgggcagaag	gcaggggtgca	caggtggctc	cagtctcagg	aggatgcgcc	1140
tcctctcccc	catcccagcc	ctcagcattg	tgccagagt	atacccttaa	gtgtttgggt	1200
ttatgttttc	agttttgttt	gggaaacagc	agttgcacag	agagttgggg	gtactgctgc	1260
tgccctttca	ccgaggcact	gccaccacca	gctctascag	ggatgctcct	gagcttggcg	1320
gacatactta	gatcctaacg	tgccagttag	acctggctgt	ggagagtagc	actggcagcc	1380
ctgcctggac	tcacttggc	atgataccag	ctccagaagg	gaagggagt	gagcaggcag	1440
tgaggagaga	gcctgggggt	cggctgggga	cagccgtatg	tgctaggtag	gagtggaggg	1500
agatatgttt	accaaagtc	tgctctgcca	tcctcccagg	tagtcagagt	gagctacatc	1560
ctgccccgcc	ttcattttcca	tggaacatg	gcagctagga	cacggggtag	aacagcagcc	1620
aaattcttcc	ccacctccct	tacttcgaaa	aaaagtgttg	aacctgggtc	cctatactct	1680
gcagtcagaa	gtgggactga	gccatacatg	cccttgaatt	cctccctgtc	tgccctccc	1740
tctccagcaa	gcaggggttt	ctttaacttg	gcagtgtgca	gaggagaagt	ggtaacaccc	1800
ccacccatt	ccctgcacg	ggagctcagt	attctacag	ggtaagaggt	aggaatcttg	1860
ctgggacgag	gggagccaga	agtggcaata	aaagcgtgtt	gacctggaaa	aaaaaaaaa	1920
aagggcgccc	gc					1932

<210> 42

<211> 1164

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (582)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (592)

<223> n equals a,t,g, or c

<400> 42

ggcacgagct	tgtgtgtcac	cagcctcctg	atctgccagg	gtctgctctg	ggttggcact	60
gaccaggggtg	tcatcgctcct	gctgcccgtg	cctcggtctg	aaggcatccc	caagatcaca	120
gggaaaggcca	tgtgtctact	caatgggcac	tgtgggcctg	tgcccttcct	ggctgtgggt	180
accagcatcc	tgccccctga	catcctgcgg	agtgaccagg	aggaggctga	ggggccccgg	240
gctgaggagg	acaagccaga	cgggcaggca	cacgagccca	tgcccgacag	ccacgtgggc	300
cgagagctga	cccgcgaagaa	gggcatacctc	ttgcagtacc	gcctgcgctc	caccgcacac	360
ctccccgggcc	cgctgctctc	catgcggggag	ccggcgccctg	ctgatggcgc	agctttggag	420
cacagcgagg	aggacggctc	catttacgag	atggccgacg	accccgacgt	ctgggtgcgc	480
agccggccct	gcgcccgcga	cgcccaccgc	aaggagattt	gctctgtggc	catcatctcg	540
gcggggcagg	ctaccgcaac	tttggcagcg	ctctgggcac	antgggaagc	angccccgtg	600
tgggggagacg	gacagcaccc	tcctcatctg	gcagtgcctt	tgatgctata	gcgcctcccc	660
tctccctca	gagggcacag	ctgcaggcct	gaccaaggcc	acgcccggct	ctcgtgctct	720
aggacctgca	cgggacttgt	ggatgggcct	ggactctcca	gaaactactt	gggccagagc	780
aaaggaaaac	ctcttggttt	aaaaaaattt	ttttcagagt	gttttgggga	ggagttttag	840
ggcttgggga	gagggaggac	acatctggag	gaaatggcct	tctttttaaa	agcaaaaaac	900
acaaaaacctc	acaactgcct	ggcaagccca	gtatcacttg	tttgggcccct	agcgggactc	960
caaggcagcc	acacgcccct	cctggaagg	tgtgtgcgtg	tgagtgtgtg	cgagtgtgtg	1020
ggctggtgtg	tgaatatcta	taataagta	tatatggtgt	atattatatg	tgtataaata	1080
aagtctgtac	atattggagc	tctgggagat	gctggaataa	aagacaagag	ttacatctgg	1140
acttgaaaaa	aaaaaaaaaa	aaaa				1164

<210> 43

<211> 1105

<212> DNA

<213> Homo sapiens

<400> 43

gaattcggca	cgagaacaaa	ttgaaacccat	ctgggtcatga	actttttat	gttaagaggt	60
tttctaatat	tgattcaatc	tcttttgctta	ttctaaatgt	gttcagattt	tccacttctt	120
gagtcattt	ggtaatttat	ttgtttctag	gaatttgtcc	atttcattct	gtttacctaa	180
tttttgacat	ataaaattat	atatggaaat	ttctaaaata	tttaaaatt	tctgtaattg	240
caatagtaat	gtccctctct	ttgtttaccaa	ttgtttat	gaatcttctc	cttttttttg	300
tcaatctagc	taaaaatttg	tcaattttgt	tcgtctcttc	aaaaaaatat	acgtttgtct	360
tcatgatttc	tctawtggtt	ttccatcyat	atttcatttg	aatacatttt	taaaacyttay	420
ctttattatt	tcattccttc	tgggagcttt	gggtctcatt	tttttttcct	gataatctag	480
ttgtttattg	tataagatta	agtatttatt	tgaaatctgt	atgttcttta	atgtaggcat	540
tcactactat	aaatttactt	ctcaggagca	tctctgcgc	attccatgtt	ttagtatgtt	600
gtgtttta	ttgtattcat	aactagaggg	aaacagaggt	gacggagaaa	aagacgtaca	660
aatatcatcc	acttgcaaa	tatagatttg	ttgtattgk	ratatgaatr	aaaatattac	720
gagacagata	agaaaaattg	aacactgacc	attgatgcag	ttacagttaa	ttttaaaatc	780
aagggttaata	acatttttagt	tatttttaaag	aatgatagta	atttagagat	gtattctgaa	840
tgttttttaa	tgaaaagata	tgcctgggat	ttcttccaaa	atgaatcttg	taggttggga	900
agaaaaatgag	aacatagtg	aaacaagact	gacaatgagt	tgttgaggtt	gggcaatgcg	960
tacactaaag	cttatttttat	cttatttttac	tgtatatact	gttaaagctt	gcattatttt	1020
cataaatgca	tttgctaagt	gcaactgtta	tcaataaag	tggattgggc	tctaaaaaaa	1080
aaaaaaaaaa	aaactcgagg	ggggg				1105

<210> 44

<211> 1262

<212> DNA

<213> Homo sapiens

<400> 44

cagcatgtac	ccagttgttc	tttctcctga	gaaagcaaaa	tgcttgatat	ttcttataat	60
ccaggctgcc	acgtttacct	tgtaaaaatca	atacttaatt	tttagatttt	tattattatct	120
tttctcgtga	agcaagactt	ctaaattatg	gtataaatat	cttttgaatt	gttgttctta	180
atgaatcttc	caactgtaaa	ctcatctaat	ttcaaactta	tcataacctga	ggatgtaaca	240
ttgtcctttg	tttctcatct	tgatattacc	gtcaatcatt	ttgtatttct	gagtacattt	300
gaacttgctg	gagtaataga	gggaaaacct	ctgcctgatt	ctaaatcaga	tctttgtcct	360
atactcggac	aattatgggt	tcataattta	ttatttttta	ttttctgggt	ttaacaaatg	420
agataacatt	ttagacataa	tatttgtaaa	catcttgact	tatttcagca	ttttcctttt	480
ttgtgtatct	tcagagagtt	tgttgaaagt	agcaatttcc	aagtaatttt	aaattattga	540
agtctactag	cacgaaaggt	caaattctta	ggatatttaa	aaaatgttgt	ttaataatca	600
aactcatctt	aaaaaatggt	catcagactc	tgtctttgat	gcacattttg	ccaaaagaga	660
gccttatttc	tgtgaaagaa	atacagtatg	tactttggga	tttactaaa	taaaaactgtt	720
actttaaggc	acagagcaga	tatagaatcc	ccctctctcc	ccactcctag	tgactgggtat	780
tctacattaa	tatttatctt	ccatgcatag	tgtacttgag	ggaaaaaac	aataactctt	840
aattgtttta	tatcaaacaa	taaaatcctg	tgtatcagtg	actgtcaata	gatggctttc	900
tgtttaaaaa	ctgaagctac	tccagaagta	ggaattaatt	tatttagtaa	acaaagtcag	960
tcaaaccaga	gccatgtcct	ggggaactgt	caaaagaatg	gttcctaagg	gccagaggcc	1020
acatccactg	gtagatgaca	gaacaacccat	acttcagatg	gcaaaaccgg	tcagtttggt	1080
ttgcgttggtg	tgcttatcct	ctttctgtgt	gcttcagctg	aattaagtgc	ttggagagct	1140
caaatagttc	aagatagcca	agatgaccaa	ttctgccagg	tggcaagcct	gatcttgcaa	1200
ttttgattaa	aataaagaac	attccccaag	aacagtttgt	tgcaaaaaaa	aaaaaaaaaa	1260
aa						1262

<210> 45

<211> 517

<212> DNA

<213> Homo sapiens

<400> 45
gaattcggca cgagtgcact tccaccagct atgtatgaga cttcccattg ctccacatct 60
ccagtatttt atgtgggcag tccttttggt tttgggcatt ttgggtggata tgaaatggca 120
tctcagtgtg gcttttcatt atatttcctt gatgactaat ggtattcctt caccctttca 180
gtgcttattg gccattcatg tatctttggt ttttggttag cacttcaggt cttttgcccc 240
tagatttagt ggggtgattg ctctttatta atgatttgta gggatgttat atatatcttg 300
gacacaagat tattgttaga gatacgtact tcagatattt tctcccagtc tgtagcttgc 360
ctaattatta ttattattat tatttgagat gaagtctcac tctgtcgcgc aggcagaggt 420
tgcagtgggc cgagatagca ccactacact caagcctggc tgacagagtg agactctgtc 480
tcaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aactcga 517

<210> 46
<211> 858
<212> DNA
<213> Homo sapiens

<400> 46
agaaaaaatc ctacatggat attggtagga aagagagaaa ggaagtggcc agtgtcccgt 60
ggcctcttcc accttctgga ttgttgaagc tggggcctgg aggggatggt cctgccactc 120
agcagggggc actaatggga ccaagctaac ctgtccagtg agaatcctgc agggagacct 180
gaggggtacca ggaagtgcga ggggaaggcc cgggaaatgg agagagctgg tctggagggg 240
aggagcaagc cgcgtggggc agggcatgtg ccttttgctt gggggagtat tactcatttg 300
gagctgtgcg tctggaacgc ctgcctcaca cacaaggac tggggcagat gtaagtcttc 360
tgcagcaacg aagcgcacag ctgagagtaa cttagaaagc acccagctaa tgctggcatc 420
ccagatcgac cccctcctcg ctgaatgttg gcactctctg gcctcagttt cctcatctgt 480
aaatgggggt gataagaaat gtgtacacac ctcccgggca gtggggagga ttaaaactgtg 540
ctctgacacg atccgggcat gttcagggtg gtatctgcag taaaccgcgc tcggaaaatg 600
gcgggcgcatc agggccagcg gtgggagctc tccgtgcttg gcttgacgcc attgtggagg 660
tgaggagggg gctgcaagac tctgagcagg aagacccgcg aaagcaggaa agcagagcca 720
gagttggggg ccagccgcag aaacgagagc ccccgtagt ttgaggcacc ctttgagag 780
ggcaggaagc aggaagggtg aattttctcc aaaacccaag aggcagagtg accccacatg 840
ataactgagt ttctcgag 858

<210> 47
<211> 6107
<212> DNA
<213> Homo sapiens

<220>
<221> SITE
<222> (5749)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (5892)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (5896)
<223> n equals a,t,g, or c

<220>
<221> SITE

<222> (5906)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (5957)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (5966)
 <223> n equals a,t,g, or c

<400> 47
 gcagtttagtt ccttgatgtc agtagtgggc taaaggcagc ttactgtgtg tttgctggag 60
 ctttcactca gccaaagtgtt agagtcagga aacccattga ggcaatggcg tcaaatgggtg 120
 tttcacaaaga atgagccatt cagtctttgc tcactatata tttaatattt tattattgtt 180
 gttattgtta ttattaattg gctttctgta ttctatgcct tttatttata aagacactaa 240
 gaaaaacccat gtttgtaatt ttaataacat ttttccatc ttgtaatatc cagagctact 300
 ttataaaattc totgaaccaa aagtattttc ctccagtgtat ctcttctccc ccagccccta 360
 ttgggaaaaa ttaccagta tagttcaggt tatgaggagg atcagccaca caatccagtg 420
 cttcagtttg aaaatgtaaa attctaacc taaagtaggg ttggttgaaa tttcagacaa 480
 agcaaaccga gcaggtataa aaagtagtat aaatacaaat ctgtaagtta tttttgaatt 540
 ttctgaactt ttttctaaga gattacatag gagactaaag aaatctatct gttcaagttc 600
 taattaggat gattgttaat actgcactgt ggatgaagtg ggcactggct tgtgtgctga 660
 cttctgtggt ttagcaagag gtttattgtt atcaaatgct aattggcaat gccaaagtcac 720
 tgggaccaat tttctgtttt ataatatcta agtttagaac agaatatata cctgaactgt 780
 agtggtttga tcggatggag acagaaaacc cgatttttct tctcataaat tttgtggtta 840
 tttatacaag ggctgtgcta tgctaccata ttcttggtca ataataatag gtttgttgtt 900
 ttttttaccat tgtaaagtgt tccttaccoc taaagggtcaa tgtaaagtac aacattctga 960
 aaatacaatt tggctacgaa gagtattcat cttctttgaa gctcagtggt tgatatttgt 1020
 gctaataatg caatttctctg attcctgtta caagttatag ctacatatgg gagagactca 1080
 gtgagccagc aaaggccata gaaacaacaa tttattaaat gtatttatgg cagaaggacc 1140
 taaataaaact gtgagccacc ttttcttctt tatattgtta catttaagt tcttgccttt 1200
 cagcaactca cattaatgct tggagcttat ctcttctctc ctctctctct ctctctctct 1260
 ctgtgtgtgt gtgtgtatgt gtgtgtgtgt gtgtgtgtgt ttccctattg tcattccatt 1320
 atatatccac accaactatg gtgacgataa ttcaaagtca tattttgcct ctaagcttga 1380
 tcattgttacc tttatgatta aagtatcatg ttatttagcc aatgcaaact tgttttaaaa 1440
 caaatagttt aaaaaaagaa caagttttta agggctttat tatagaagaa gtattaatga 1500
 aggactttcc ttctccctc cctttccctc cctccctgcc tcccttcttc ccttccatct 1560
 cccctccctc cctgccttct ttgtttctcc ttcccttatt cctccctccc tcccttctcc 1620
 cttcttctct ttcttccatt catccttctc tgccctttat ttttattttt tgtaatatca 1680
 catgtgctgt agtttggaa tttattctag tgcatttctt gctcatcaga acctcagcta 1740
 atctacctag gaaaaatagt atcaaaggaa atgagaaagt tgtatctgag tccctccaga 1800
 actaagataa ttctttttga ccatttaagc ctttataaat gcgttttgac catttaagcc 1860
 tttataaatg cttgttttag gaaagtgaat ctgttagatg catcaacaaa taatgaccag 1920
 gacaaaacga tttataaatt aaagtcctca atcaccatgg ttatacattt tcaccagaaa 1980
 tagtaatctt acaatttttc atttttctga tgaagatttc tgttccaata tctgtttcct 2040
 aatagatttt ttaaattaat tagctttcct ctgctttatg accacaggtt ttatccctaa 2100
 ccgagacagc tgtcttatat ctgcatgcct tagactgtgt ggagggactc catgaagaaa 2160
 gaccataggt tagaaaaata actcatagta tataccctag taagtgggtt agtagaatct 2220
 cataacatgt attaaaaaga ggttttcttc tctgcttgtt tgtgtcacta gagcaaaatt 2280
 gttagagataa tgctcataat gcagtaataa tcagaataat ctacaatatc atttgggat 2340
 ggtcccaggt cccagtgtc tagttacttt acttcttttt ttttttttga gatggagtct 2400
 tgctctgtct ctcaggctag agcagtgtgc gatctcagct cactgcagcc tccacctccc 2460
 aggttcaagc gattctcctg cctcagcctc ccaagtagcc aggattacag gcacctcca 2520
 ctaggccggg ctaatttttt ttgtattttt tttagtaga tgggggtttg ccatgttggc 2580
 caggtcgggt tcgaactcct aacctccagt gatccacctg cctcggcgct ccaaagtgtc 2640

aggattacag	gcatgagcca	ccacatccgg	cctaattact	tctttaatcc	ccatttattt	2700
ttatgccatt	ctagcctcat	ttattaataa	aattatgttt	ttactttctc	tttcaggaaa	2760
tttttttaaat	taatatTTTA	tatctagatc	taatgctatg	gaaaagtgcc	tttttatcat	2820
ttataaTTTC	atTTTTCact	atTTTCCaaa	acacataaac	aaatagTTTC	agtaggtccc	2880
agctTTTact	TTTTccattt	aaaccttctt	ttctccattt	cttccctttg	gcttaagaat	2940
aaaagaaaag	gtacattgct	agaattgttt	ctttgggaga	gggtaaaaga	ttacagaatt	3000
agactgttca	gcctttatat	aaactaaatt	tgtcttcac	tcaaccagct	aatggtaggt	3060
cttatctgaa	tactcatgag	aatttttagca	tctgtgaaac	tccatgcacc	agatgtgtgt	3120
aaatttcagg	aagaaaagtgt	tgaaaagcatt	ttctctgatg	ttaattagat	ggaaataaat	3180
cactaaaaaca	tagtttaggt	aaagcctgat	tatgccactt	TTTTttaact	agacagggca	3240
aagttgttta	tgttagtgt	cttcttgtct	atcctcagtt	aatttaccta	gacaaaaagt	3300
gtcaaaaggaa	atgagaaaaa	ggttatatct	gactccctcc	agacctaa	taattccttt	3360
tgatcagata	cagtcagatg	gagtgccctg	gtttttgtta	atTTTgcctc	tattccagct	3420
ccttaccaca	gcggtgtgtc	ttaaagaaa	gatcatcagc	aacaggtcag	gatagttcta	3480
cctttgggat	agggctgctt	ttcccgtgct	agtattttctg	tgactgttag	tggcactgag	3540
gactgcaaac	TTTTatgcaa	tattcttaaat	accctattga	tattatgcac	TTTaatcatt	3600
ccaaagaagc	caagaatgct	gtatagtgt	gatttcttcc	taatgaattc	atcttaacta	3660
tttagaatgt	tatgtccctt	ttcttttTga	tagccaaact	ggtataaatg	ttatatggat	3720
TTTTctaaaa	tgactatata	ggacttaaga	ctttgaaatg	taatttactt	ataaggggaa	3780
ataatttatgc	tttagcacat	catttttagaa	acgtcacatt	ttagaaacat	tcagcttgct	3840
aacctacatg	tttgTgaatt	cattaaaaac	agttgtctat	atatTTTgtg	ccatgtatat	3900
aagaacatta	caatatatct	TTTTctacat	atgtagtatg	tgcaaccagt	ggttctcaga	3960
gtatggTTct	cagcccacca	gctagtatca	gtatcacctg	ggaactagtt	agaaatgtaa	4020
attctttggc	cccattcccag	acatactgag	tcagaaattc	tggaataggg	cccccgcaat	4080
ctgTTTTcac	aagccctcca	ggtgattctg	atgcacactt	taaagTTtag	gaaccactgg	4140
gctaagactc	tgTTgagata	tagagTTTTt	cttccactca	gactgatata	gTTatacatt	4200
gTTcttcatg	taaattcagc	TTaacctggT	tatctataat	ctTTtatTgg	caaaagTTaa	4260
TTctcagTac	Tgcctataga	gatacagtgt	atTTtatgt	catacacaat	tagtctaatt	4320
cttgataaTT	cagTTaaTTt	agTTTggcat	TTtcttacca	cttactaaaa	ggtttacatt	4380
aaatgactga	TTTaaatata	taggtgcaat	gttctatgtt	tattTTtaatt	gTTatgacat	4440
TTaagtagct	aatataaTTg	accggtgcta	aagtctctcg	TTtatccata	aaatgggtac	4500
atttatgggca	gtgtaataca	agctTTcttt	tcattgccta	gtactTTtacc	agcagaccac	4560
agTTTtgccc	tggttagacc	aaacctcaga	acaaaatcat	cattccttgt	atttatattt	4620
gtatctgaga	tagtaaacaa	gatggctggc	caggtcaaca	tggcacctta	acttatTTtt	4680
TTaataggt	aaacttcttc	aaaagtagct	tgctTTgtat	aagaactaag	ctatcagtat	4740
agatatagct	atccttgag	cttatgtttc	agacaagaat	tatttactaa	aataaataat	4800
aaacaagata	atgcattata	caatttgggc	atTTtctcgt	tctcaagtgt	atgcacatg	4860
gTaaatataa	actaaccaca	agataggtag	attgattcat	ttcattTTta	tctccttgtg	4920
taatttcagta	cctccataat	tgTTtcaatc	ttcttcccac	tgTTtaca	ttaccagTTa	4980
attaactcgt	gaaagaaaaa	ttcacatatc	agaataaaaa	taaatgtata	ctcactTTtat	5040
aaaaatcacc	actgctgtct	ttccttaata	ctagcagTgg	aaatgtaagt	ggcttactct	5100
acaaaTTttg	gtgctggcaa	atacataggc	aaactgtTgg	gagctgctct	agttacattc	5160
ctcccttctt	attccctTTt	tctcttccct	actTTtattg	ataacataat	ctgttaoccca	5220
aagcattctc	ccacagttct	atTTgactcc	cacttgtaat	aactcctTTa	aaaaattcca	5280
tgTTtaacca	tatgaccttg	cttgcttact	catattctcc	ctccctctcc	ccttccctttc	5340
tctctcttcc	agaagtcatt	TgcctggTTt	gaaatatttt	gtagggattg	cttatttatat	5400
tatttttagct	gatgaacctc	aggacaacgt	ctacacacac	acacatacat	acacgcacac	5460
aaaatctcag	ctgttgaaga	gtgggctTgg	aatcagactt	ctgtgtccag	taaaaaactc	5520
ctgcactgaa	gtcattgtga	cttgagtagt	tacagactga	ttccagtga	cttgatctaa	5580
TTtctTTtga	tctaataaat	gtgtctgctt	accttgTTtc	ctTTtaattg	ataagctcca	5640
agtagttgct	aattTTtttga	caactTTtaa	tgagTTtcat	tcacttcttt	tacttaatgt	5700
TTtaagtata	gtaccaataa	TTtcattaac	ctgttctcaa	gtggTTtanc	taccattctg	5760
ccattTTttaa	TTTTtatTTa	atTTtatTTg	cttgagcaca	ctgatcaacc	actgaactgc	5820
cttcttccat	Tgtcctgcaa	Tgatataagg	gttcaatttt	Tgtgtatatg	gctttcatag	5880
TTgggatttc	anagcncTga	taccanatat	TTtcagTTtg	TTctctgggg	gaatttcatt	5940
Tgcatctatg	TTTTtancta	tctgtTataa	cttgTtaa	attaaaaaga	tattttgctt	6000
ctattTgaac	atTTgtatac	Tcgcaactat	atTTctgtaa	acagctgcag	tcaaaaaataa	6060
aacactgaaa	gTTTTcaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaa		6107

<210> 48
 <211> 703
 <212> DNA
 <213> Homo sapiens

<400> 48
 ccacgcgtcc gcaggacatc gttttctaca tgggtggctgt gttcctgacc ttcctcatgc 60
 tcttccgtgg cagggtcacc ctggcatggg ctctgggtta cctgggcttg tatgtgttct 120
 atgtgggtcac tgtgattctc tgcacctgga tctaccaacg gcaacggaga ggatctctgt 180
 tctgccccat gccagttact ccagagatcc tctcagactc cgaggaggac cgggtatctt 240
 ctaataccaa cagctatgac tacgggtgatg agtaccggcc gctgttcttc taccaggaga 300
 ccacgggtca gatcctggtc cgggccctca atccccctga ttacatgaag tggagaagga 360
 aatcagcata ctggaaagcc ctcaagggtgt tcaagctgcc tgtggagtcc ctgctgctcc 420
 tcacagtccc cgtcgtggac cgggacaagg atgaccagaa ctggaaacgg cccctcaact 480
 gtctgcatct gggtatcagc cccctgggtg tggtcctgac cctgcagtcg gggacctatg 540
 gtgtctatga gataggcggc ctcgttcccc tctgggtcgt ggtgggtgac gcaggcacag 600
 ccttggcttc agtgaccttt tttgccacat ctgacagcca gccccccagg cttcactggc 660
 tctttgcttt cctgggcttt ctgaccagcg cctgtggat caa 703

<210> 49
 <211> 639
 <212> DNA
 <213> Homo sapiens

<400> 49
 ggcacgagca ttcacagggt acaaatgctg ctgccaaactg tctgggcaa atgactctgc 60
 atcacaaacc tttccttgca tgtggagggg atggatttac tcagtccaac tttgatggct 120
 gcataccttc tgccctatgt gttctggaag ctttaaagaa ttatatttag tgcctatata 180
 cttattctct acatgtgtat tgggttttta ttttcacaat tttctgttat tgattatttt 240
 gttttctatt ttgctaagaa aaattactgg aaaattgttc ttcacttatt atcatttttc 300
 atgtggagta taaaatcaat tttgtaattt tgatagttac aacctatgct agaatggaaa 360
 ttcctcacac cttgcacctt ccctactttt ctgaattgct atgactactc cttgttggag 420
 gaaaagtggg acttaaaaaa taacaaacga ctctctcaaa aaaattacat taaatcacaa 480
 taacagtttg tatgccaaaa acttgattat ccttatgaaa atttcaattc tgaataaaga 540
 ataatcacat tatcaagacc ccaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 600
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 639

<210> 50
 <211> 867
 <212> DNA
 <213> Homo sapiens

<400> 50
 ggcacgagca ggtactgggt gactgcctgg ctgaggaaaa gttaactaga cacttgggga 60
 aaggagatcc aaggagtaga gaggcaaaat gcctttgcat gcttttcttc ctatctcttt 120
 ttctttctct ccttctcact ctctcccttc ctctctttct tctttctctt ttcttttttt 180
 tttctctttt ccccccactc tctgctgccc tcttctcttc cctcccctcc cctcccctcc 240
 cctctccctc ctccctccct tcttctcttc ctctctcttc tcttctcttc ctctctctcc 300
 tctctctctc ctctctccct gccttctttc ctctctcttg ccaacttgcc agaaggagcc 360
 caagaaaaag caccagatg cttcagtcac cttcttagaa ttcttctttt ttttatgttc 420
 agaaaaagatg gaaattcatt tctgctaaa agaaagaaaa aattggaaga cagggtgaag 480
 gtgaacaggc ccattataag aaagaaacaa aaatctatat tctgtctaca aggaagcgag 540
 agagagaaag agagagaaga aagaagttcc aggattctaa tgtaccaaag ggatctcctt 600
 tttcttgttt tgttctgaaa atttcaccaa aagagcacag gagaacatct tggctaattc 660

29

attggcgatg	atgtaagaaa	actgagagaa	atgaaagaaa	tgaagaatta	ctgctgcaga	720
taatatacag	ccttgaggaa	agaaaggctt	ttaagattat	agatataaag	gctattgctg	780
tattctggga	taaaagaaa	tctgatgtca	gggaaagggg	aagttggaaa	aactggaaaa	840
agaaaaaaga	aaaaaaaaa	aaaaaaa				867

<210> 51
 <211> 1569
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (341)
 <223> n equals a,t,g, or c

<400> 51						
gtattggcca	ggctgggtctc	aaactcctga	cctcgtgatc	cacccacctt	ggcctcccaa	60
agtgcagaga	ttacaggcat	gagccactgc	acctggcctc	aagaaaaatt	atatatcacg	120
tggaaatagg	tagtagtctc	tgcactgatt	ttcgttgata	atggctgttc	ttcttatcac	180
catttttgcta	tttctttgtc	tgggctatta	caggggtatt	acagaaattt	ccagaaagac	240
ccctgcctgt	cgaatgttta	cttcaagctt	gagctcctgg	tatattatga	ggaaattata	300
tgatacceca	ggagaggtct	tcctttccca	tgccattgta	naattcctaa	agtaaaatta	360
atttgccttc	ttgtcaaaga	aggagccaat	gttgttttaa	aatttttagct	tgagagatag	420
gtggggaaga	aattaaatag	acaagtaatc	mctattcaga	agagaaggga	gagtcattgt	480
acgaggccca	agatacttgc	ccaaaaatat	cgcagagaaa	aactagtctt	tggggtccta	540
tttttttagt	ggaacatttg	agttatttaa	aattagaatt	ttattttggt	cagattagaa	600
tttctagggt	atgtcatatg	tgttttttaa	ttgaaagctc	ttaaaactcc	tattgtagtt	660
taatgtcatt	atccattaat	ttacataaat	ctgatttgga	tctctatttt	catcgtagac	720
tgtgtagggg	caatttttcc	taaaggttct	gtgacatagt	gctacctttt	ttttaaaacc	780
tgtcttgccc	aggcattatt	gagtgccccc	tgggtgccagc	atgtgtattt	cacgactgta	840
tcaacaaatc	atgatcatct	tctctggcca	ttgtgccttt	tcagattcca	aactgtgtac	900
ctctcagtc	ttcctacaaa	cttagaaagt	ctaatatctt	aatgtttact	tatgtagcaa	960
cctcccttcc	tcccatccct	aaatcctctt	gtaattaatt	attttccttt	ggaacttttt	1020
aaatctacaa	tttccctata	atatggtaac	caatattaat	tttcttggtc	tgcgccaagt	1080
ttgattttat	acaaattggt	tccagtttgg	gtcatgagca	caaaaccagg	tatttttaaa	1140
aatctatata	acccttcaat	gaggcagtat	taattttatt	aactcattaa	ttcaaccaat	1200
aattcctgat	tgtttactgt	gttagatatt	ggggtatccc	caatacctga	cagctgtgag	1260
caaaacaaat	gccctacaca	catgaggtgt	acagtcaggt	agaaaagata	aacaataagc	1320
aaattaatag	ataatatgat	gtccaataag	gacttcaaag	gaaaataaag	cagagtaaag	1380
agccagagaa	tgacagttag	ctgtttttca	catgagtcac	cagaaaaggc	ctctttaaag	1440
aattgacatt	tgaacagaaa	aacgaatcaa	gggcgtcaac	tgtttattgc	ttttattgct	1500
taccatttga	ccaagcaatt	ctacacatag	gattcacctt	aaaaaaaaa	aaaaaaaaa	1560
aaactcgag						1569

<210> 52
 <211> 1196
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (590)
 <223> n equals a,t,g, or c

<400> 52						
gattgggtct	gtttatgtga	tagattactt	ttattgattt	gtatgttgaa	ccagccttgc	60

atcctagggg	tgaagccgac	ttgggtgtgg	tggataagct	ttttgatgtg	ctgctggggt	120
tggcttgcca	gtgttttatt	agggattttt	gcgtcaatat	tcacagggga	tattggcctg	180
gaattttctt	tttttgttat	gtgtctgcca	ggttttggta	tcaggggtgat	gctggcctca	240
taaaataagt	tagggagggc	tccctctttt	tctttcattt	ggaagaattt	cagaagggaat	300
ggtaccagat	ccyctttgta	cctctggtag	aatttggtctg	tgaatccatc	tggtcckgag	360
cttttttttt	gttggttaggc	tattaattac	tgctcaattt	tcagaacttg	ttattgggtct	420
attcagggat	ttgacttctt	cctggtttag	tcttgggagg	ttgtatgtgt	gcaggaattt	480
attcattttct	tctagatttt	ctcgtttatt	tgtgtagagg	tgtttatagc	atyctctgat	540
ggtagtttgt	attctgtggg	atcagtgggtg	atctcccttt	tatcattttt	atttgtgtcta	600
tttgattctt	ctctcttktc	ttctttatta	ttctygttaa	tggtctatgt	attttgttaa	660
tctyttacaa	aaacaggctt	ctagattcat	ggatgttttg	aaagggttyt	cgtgtctcta	720
tctccttcag	ttcttccctg	atcttagcta	tttcttgtct	tctgctagct	tttgaaattg	780
tttgcttttg	cttctctagt	tcttttaacc	gtgatgtcca	gtgtgtcaat	ttcagatctt	840
tccagccttc	tgatatgggc	atttaatgct	ataaatttcc	ctcttaacac	tgcttttagct	900
gtgtcctaga	gattctggta	cggtgtctct	ttgttctcat	tggtttcaaa	taacttcaat	960
atttctgcct	taattttggt	atttaccag	cagtcattca	agagcagggt	gttcaatttc	1020
catgtagttg	tgtggttttg	agtgaatttc	ttaatcttga	gttctaattt	gattgcactg	1080
tgtctgaga	gacggttaca	atttccattc	ttttgcattt	gctgagaagt	gttttacttc	1140
caattgtgtc	tcgtgccgaa	ttcgatatca	agcttatcga	taccgtcgac	ctcgag	1196

<210> 53

<211> 945

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (295)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (875)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (914)

<223> n equals a,t,g, or c

<400> 53

gaatggtgaa	atattaagtg	ctttctcccc	caggttcagg	attatgacag	ctatgtccat	60
tcacctcttc	tgtacagcat	tgtcctgtgg	aagttctggc	cagtgaata	aggcaattaa	120
aagaaataaa	atatcaaacg	attggaaaga	tgtaaatgtg	tcacattcca	tagaaaacat	180
gattcataga	tatacataca	cgaatgcttt	gaattcataa	gtagattcag	ccagttgctg	240
gatataaagt	caatatacaa	aaactatttt	tatagacatg	aaacacgcaa	tgagnaaaaa	300
aatttaacca	tttttagtag	catcaaaaaa	cccccatacc	taggaatatg	aattttagtg	360
actatttggg	atatgttgat	ggatatattat	catttccagt	ttgggattat	tataaagaaa	420
atagccctga	acatttgtaa	tatatgactt	ttgggtgaatg	tagcattcat	ttctgttgat	480
tacaaactca	ggggtgaaat	tgttgagtc	taagggagct	atagatgtat	tcaacttcag	540
ctgatatggc	taataaaatt	tgcgaaaaag	attgcatcaa	gttatgctcc	catcagcaat	600
atgagagttc	ctgtttttcc	acattgtcag	caacactttg	tactgttact	ccttttaatt	660
ttagccgatt	tggtgaagg	tgtggttaata	tctcattgta	gtggccaggc	gtggtgctca	720
cgctctgaat	cccagcactg	tgggaagcca	aggtgggccc	atcacgaggt	caggagatcc	780
agaccatcct	ggctaacatg	atgaaacctt	gttgccctgta	gtcccaacta	cttgggaggc	840
tgaggcagga	gaatggcatg	aactcgggag	gcgngcttg	cagtgaacct	ccagcctggg	900
caacagagtg	agantctctc	aaaaaaaaaa	aaaaaaaaac	tcgag		945

<210> 54
 <211> 488
 <212> DNA
 <213> Homo sapiens

<400> 54
 ggacacgagga gagtagagggc tattcatgta atgtctataa aaaaataaca ccaaggctgg 60
 gattacaggc atgagccact gcacctggcc agtttgctta ttttgtttg tgcctcctcc 120
 catgggagac ctcaaggagg tatgcctgcc ccacagatgc cctggaagga cagcttgctg 180
 ctctactca gaaccacacc tgcagacaga ggaggacaga cggacactca tttgctgagc 240
 acccatgtaa catgaactaa gagctgggtg gagacaatga acggtggagc catcgttccc 300
 gatgtggagg gagaacagct caagaccacg gaacagcctg ctctcccgt tcttggttc 360
 cgtgcgcttt tgtccaatca ggcttttga ccaatcggcc aggcgcgcta tgtaaatttc 420
 tgacattttc aaagctgtct ttttaataaa cctttcagtg taaaaataaa aaaaaaaaaa 480
 aaaaaaaaaa 488

<210> 55
 <211> 2860
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (753)
 <223> n equals a,t,g, or c

<400> 55
 ggacacacagg gctggcaggc ccgcggtggc tgggtgtgag gcatgaacaa attgtaccgg 60
 gtatccccca cccactctg accaccagtt cctccttggc tatcactccc cctgacaggc 120
 agcccaccca ggccctggatt tgtccctgtc tcccccttt gcttttccc catgactaat 180
 gggcaccagg tcttgctgct cctgcttctc acctctgcag tggcagcagg cccctggccc 240
 caggtgcagc ccggtcagtg ggggttgatg tgccctcctc caggcctgcc ctctgtccaa 300
 gccccggagt ggcttggtgg gctccctggg ggccccagc ggggtgccagg tgggtgcccg 360
 ggttatttag ggggtggtgt atcactgtag ggacaggctt ctgccccag cctggagagc 420
 tgttttcttc aggaaggctc tggagatgga gacttggttg cgaattcacc acaactccag 480
 gctgggaggc tgggtctctg ctctcagagc cgagacacca gggaggatag ccaggctgcc 540
 ctgcctggga attctgctgg gccgtcaaat tcaacccgca ccaacgtggg caggaggcca 600
 cagtgtcctg ccaggagcag agggctgaag gtctgcagga ggaagaccct atcctgggtg 660
 ggggcacctg ctgcccaccc tgcctccagc gtgcctgggg ggagcacacc tgggcatgga 720
 ggagtccagg gtgctggggc acacaagaga ggnnggggag aggcctggac agtaggaaga 780
 tcttgccag ggtcctggat ccgccactct gggggtgacc ttggacaaac ctctgccttg 840
 gccctcagtc tccccatcaa ggtttttcca ttcaggaggg tttgggccat cttcagccac 900
 cctaccagcc ctgaaaagga tgtgactcct gtttctggga agtgtgtggt gtgttaggtg 960
 ggccctacagc cctggttgtg gggagggaag gatggagaga cagcacagt acagagccca 1020
 gactgcaggc tggagttagg gttccacttc ccgctgctg tgtgtcctgg accagtgcct 1080
 ctgaaccttg gcacttgggg cagtggatat taacatcttt ccaagcccaa ttcttggggc 1140
 atcagggcct ccggtcctct gggaggtggc aggtcctcag attggagatg ccattggggg 1200
 gggaggtgcc tctccttttg aggggtatgga agtggagaca ggagtggcct ggcgcagctg 1260
 ccgtgggttct taggggctgg gcccggggag cccatggggc ttgtgcctag aaagcctggg 1320
 ctctcactg gggcttagat gtgcagactt catgtctccc cagctccagc tctgttctct 1380
 ataggtcaag cctccacaat gccagaggcc cagggttagc cccctccagc tccctcctag 1440
 atctacagct gcccttttga tgacagcgcc attgagctcc ctgggctggg ggggtcatgc 1500
 aggggtgagg cagtgcctg ccgcccgtac tcattgcctg gccaggcagg acacaggctg 1560
 gcgggcactg agagtgggccc ccacgaaatc cattgtcagg ttaccaggat gaagaaccca 1620
 ggctggctct ggagtgcagg gcggggcctg ccggaagaat tatgggact gcagcaggag 1680

ggcagcctgg	gccattagct	cctgatgtca	tcgatttggg	tgaggggaca	gggaagtcag	1740
aggaagctgg	ccagtggctc	tcacgcagac	ttacagcagt	ggagtgggtc	ctgattcctg	1800
gtacagctgc	tccactgag	tctccaggga	tctgtgggtc	aggaccccct	gcaaccccct	1860
cccagacccc	tgtactgggtg	ggaggagagg	acctagagga	aagggtgctg	gcagataagc	1920
agctgaggga	ggccctgggt	ttagcttata	agtcttctg	gccctcctgc	cccaggaagg	1980
gcagcggagga	ccatgggtgtt	gcccctgtca	tcgttatcgt	cctggccatg	agcttgcagg	2040
actgggaggg	ccggagtcag	ccaggcagac	ggcagcacag	catttgctg	ttggcagggtg	2100
gccttggtgg	cttcccaaag	gcaatcgctc	cacgcagaac	aaaactcaact	tttttggggg	2160
gtgaagcacc	ttggttcatt	tgttttagttc	gttaattcca	gcagtctgtt	tctaagggaa	2220
acatggctgc	agccggctcct	gcgcctccca	ccctcccacc	aggtgcccag	tgttcccaag	2280
ggccccgaat	cccaacctta	ttcaggcgctc	agcatctctg	caccccaaat	gcctgttagg	2340
gaggatagtg	aaggctgagc	cctcctgggc	ccatcaaaag	ccagcagtga	gagaacaccc	2400
ccatctctct	gaggtgacct	tgtaggcgag	tccgtgctgt	ctggctggcc	tgggtgagggt	2460
gggcaggggac	caaggcctgg	cgccctgggc	tcgctggcct	tgctctgctg	gctgacttca	2520
tcctgatagt	accttgattt	tcctactgtg	acttcccctt	ctgtcgactt	cctcaccaac	2580
tttaaaattc	cgtattgaga	gcagtttctc	aagttacctc	aaatcctatt	cagaagaagg	2640
ttcttctctg	aagttgggag	ggcggaaaac	aagtttagtc	acagaagact	actccatgtt	2700
tgagcttctg	tttcaaggga	agtgaagtaac	tgccggaggga	gccctgcccc	tctgcagtgt	2760
gtggtgtgtg	cctgatactt	ttcagattga	ggtgttactt	acatgtaata	aaatgcacag	2820
acttaagtgt	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa			2860

<210> 56

<211> 1559

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1445)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1551)

<223> n equals a,t,g, or c

<400> 56

atccagcagt	ggggagacag	cgtgctgggc	aggcgctgcc	gagaccttct	cctgcagctc	60
tacctacagc	ggccggagct	gccccgtgcc	gtgcctgagg	tcctactgca	cagcgaagg	120
gctgccagca	gcagcgtctg	caagctggac	ggactcatcc	accgcttcat	cacgctcctt	180
gcggacacca	gcgactcccc	ggcggttgag	aaccgagggg	cggatgccag	catggcctgc	240
cggaagctgg	cggtggcgca	cccgtgctg	ctgctcaggc	acctgcccat	gatcgcggg	300
ctcctgcacg	gcccgaacca	cctcaacttc	caggagttcc	ggcagcagaa	ccacctgagc	360
tgcttctctg	acgtgctggg	cctgctggag	ctgctgcagc	cgcacgtgtt	ccgcagcgag	420
caccaggggg	cgctgtggga	ctgccttctg	tccttcatcc	gcctgctgct	gaattacagg	480
aagtcctccc	gccatctggc	tgcccttcac	aacaagtttg	tgcagttcat	ccataagtac	540
attacctaca	atgccccagc	agccatctcc	ttcctgcaga	agcacgccga	cccgtctccac	600
gacctgtcct	tcgacaacag	tgacctgggtg	atgctgaaat	ccctccttgc	agggctcagc	660
ctgcccagca	gggacgacag	gaccgaccga	ggcctggacg	aagagggcga	ggaggagagc	720
tcagccgggt	ccttgccccct	ggtcagcgctc	tcctgtttca	ccctctgac	cgcggccgag	780
atggccccct	acatgaaacg	gctttccccg	ggccaaacgg	tgaggatctt	gctggagggtt	840
ctgagtgaca	tagacgagat	gtcccgggcg	agaccgaga	tcctgagctt	cttctcgacc	900
aacctgcagc	ggctgatgag	ctcgcccgag	gagtggttgc	gcaacctcgc	cttcagcctg	960
gccctgcgct	ccatgcagaa	cagccccagc	attgcagccg	ctttcctgcc	cacgttcatg	1020
tactgcctgg	gcagccagga	ctttgaggtg	gtgcagacgg	ccctccggaa	cctgcctgag	1080
tacgctctcc	tgtgccaaga	gcacgcgggt	gtgctgctcc	accgggcctt	cctggtgggc	1140
atgtacggcc	agatgggacc	cagcgcgcag	atctccgagg	ccctgaggat	cctgcataatg	1200

33

gaggccgtga	tgtgagcctg	tggcagccga	ccccctcca	agccccggcc	cgccccgtcc	1260
ccggggatcc	tcgaggcaaa	gcccaggaag	cgtgggcgtt	gctgggtctgt	ccgaggaggt	1320
gagggcgccg	agccctgagg	ccaggcaggc	ccaggagcaa	tactccgagc	cctgggggtg	1380
ctccggggccg	gccgctggca	tcagggggccg	tccagcaagc	cctcattcac	cttctggggc	1440
acagnccctgc	gcggagcggc	ggatccccc	gggcatggcc	tgggctgggt	ttgaatgaaa	1500
cgacctgaac	tgtcaaaaaa	aaaaaaaaaa	aaaccgrgg	gggggcccgg	nacccaatt	1559

<210> 57

<211> 2064

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (2001)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2024)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2049)

<223> n equals a,t,g, or c

<400> 57

atggggcgagg	ctgccccggcc	ccggcgccga	cgeccgcacc	tctccccagc	cctggcggtgg	60
gcccagcccg	gcccaggcag	caatgggggtt	cctgcagctg	ctgggtcgta	cggtgctggy	120
atccgaacac	cgggtgggctg	gtgcagccga	ggtcttcggg	aattccagcg	arggtcttat	180
tgaattttct	gtggggaaat	ttagatactt	cgagctcaat	agggcccttc	cagaggaagc	240
tattttgcat	gatatttcaa	gcaatgtgac	ttttcttatt	ttccaaatac	actcacagta	300
tcagaataca	actgtttcct	tttctccgag	gcgtagatcc	cccaccatgt	gacgctggga	360
cagaccagga	ctccagggtg	agggttgcat	atgatgtcta	tcagtatttt	ctgcctgaga	420
atgacctcac	tgaggagatg	ttgctgaagc	atctgcagag	gatgggtcagt	gtgccccagg	480
tgaaggccag	tgctctcaag	gtgggttacc	taacagctaa	tgataagaca	agtgtttcct	540
tctcctccct	ccggggacaa	ggtgtcatat	acaatgtcat	tgtttgggac	ccgtttctaa	600
atacatctgc	tgccctacatt	cctgctcaca	catacgcttg	cagctttgag	gcaggagagg	660
gtagttgtgc	ttccctagga	agagtgtctt	ccaaagtgtt	cttcaactct	tttgccttgc	720
ttggtttctt	catttggttc	tttgacacac	gattctggaa	aacagaatta	ttcttcatag	780
gctttatcat	catgggattc	ttcttttata	tactgattac	aagactgaca	cctatcaagt	840
atgatgtgaa	tctgattctg	acagctgtca	ctggaagcgt	cggtggaatg	ttcttggtag	900
ctgtgtgggt	gcgatttgga	atcctctcga	tctgcagctg	ctgtgttgga	ctagtgtctg	960
ggttctctcat	ctcgtcagtg	actttcttta	ctccactggg	aaacctaaag	atttttcatg	1020
atgatgggtg	attctgggtc	actttctctt	gcatagctat	cctcattcca	gtagttttca	1080
tgggctgcct	aagaatactg	aacatactga	cttgtggagt	cattgggtcc	tattcggtgg	1140
ttttagccat	tgacagttac	tggtccacaa	gcctttccta	catcactttg	aacgtactca	1200
agagagcgct	caacaaggat	ttccacagag	ctttcacaaa	tgtgcctttt	caaactaatg	1260
acttcattat	cctggcagta	tggggcatgc	tggtgtgaag	tgggaattac	ttacagattc	1320
gaagagagag	aggacgaccg	ttcttccctc	cccaccata	caagttatgg	aagcaagaga	1380
gagagcgccg	agtgacaaac	attctggacc	ctagctacca	cattcctcca	ttgagagaga	1440
ggctctatgg	ccgattaacc	cagattaaag	ggctcttcca	gaaggagcag	ccagctggag	1500
agagaacgcc	tttgcttctg	tagatgcccc	ggggcttgg	cagtgtgcct	cagctttgga	1560
gttcatgcct	ggagtgggtc	aacagtctct	ggtgcaagtc	taataagaga	tcaggcatat	1620
atattctgttc	tttgcataat	attatgggtc	ccttattgat	atatggtaag	ggtgtactag	1680
gggattagga	tgattgtgaag	agaatgagaa	agatgaccaa	aagggttggtg	gtagggaggc	1740

34

ttttttcttat	ttccaaatac	ttgagaaatt	acctttttggt	ttacaaatct	atgatcaact	1800
tattccatta	aatagatata	ttaaaaaat	taaaaactga	ttcttctgca	gagcactggt	1860
gtttcttttt	ataacccctt	gaaacaagtc	tctcacctga	gcctgtctaa	actttcggag	1920
ggagtttatt	attgagtctt	tatctgtgac	agtattttgga	gatttaggga	tttgatactt	1980
aggcctttga	attttagaat	ncaaaaagg	gagccagcca	gacngggggg	ccaacccgga	2040
tccccaacng	ggaccagggg	ggtc				2064

<210> 58
 <211> 1050
 <212> DNA
 <213> Homo sapiens

<400> 58						
ccacgcgtcc	ggccagccag	tccgcccgtc	cggagcccg	ctcgctggg	cagcatggcg	60
gggtcgccgc	tgctctggg	gccgcggg	ggggcgctcg	gccttttgg	gctgctgctg	120
ctcgccctgt	ttcggccg	ccccgcgctc	tgccgcg	cggtaaagg	gcccccg	180
ctaagcgag	cgtctccg	cttggtaga	ctggcgctcc	tcgccgcttc	cggcggtcag	240
tgccccgagg	tgaggcg	ggggcggtgc	agacctggcg	cggcgctgg	cgcctctgct	300
ggagccgaac	gtcaggagcg	ggcgcggg	gaggcgaga	ggctgaggat	cagcaggcgc	360
gcgtcctggc	gcagctgctg	cgcgtctggg	gcgcccccg	caactctgat	cgggctctgg	420
gcctggacga	cgaccccgac	gcgcctgcag	cgcagctcgc	tcgcgctctg	ctcccgcccc	480
gccttgaccc	tgccgcccta	gcagcccgag	ttgtccccgc	gcccgtcccc	gcccgggcgc	540
tccgaccccg	gcccccggtc	tacgacgacg	gcccccg	cccggatgct	gaggaggcag	600
gcgacgagac	acccgacgtg	gaccccgagc	tggtgaggtg	cttgctggga	cggattcttg	660
cgggaagcgc	ggactccgag	ggggtggcag	ccccgcg	cctccgccc	gccgcgacc	720
acgatgtggg	ctctgagctg	ccccctgagg	gcgtgctggg	ggcgctgctg	cgtgtgaaac	780
gcctagagac	cccgccgccc	caggtgcctg	cacgcgcct	cttgccacc	tgagcactgc	840
ccggatcccc	tgacccctgg	gacccagaag	tgcccccgcc	atccccccac	caggactgct	900
ccccgccagc	acgtccagag	caacttacc	cggccagcca	gccctctcac	ccgaggatcc	960
ctacccccctg	gccccacaat	aaacatgatc	tgaagcagca	aaaaaaaaa	aaaaaaaaa	1020
aaaaaaaaa	aaaaaaaaa	aaaaaaaaa				1050

<210> 59
 <211> 2533
 <212> DNA
 <213> Homo sapiens

<400> 59						
ccacgcgtcc	gcctggcaac	ccctaataatt	tgggatctct	caatgctatt	tgccctcatt	60
agcttgctcg	ttatgcttcc	cacttggtgg	atttgttctt	cctggctggt	atggggagtg	120
atttctatttg	tgtatctgg	cataagagct	ttgagattat	ggaggacagc	caaaactacaa	180
gtgaccctaa	aaaaatacag	cgttcatttg	gaagatatgg	ccacaaacag	ccgagctttt	240
actaacctcg	tgagaaaagc	tttacgtctc	attcaagaaa	ccgaagtgat	ttccagagga	300
tttacactgg	tcagtgtctg	ttgccattt	aataaagctg	gacagcatcc	aagtcagcat	360
ctcatcggtc	ttcggaagc	tgtctaccga	actctaagag	ccaacttcca	agcagcaagg	420
ctagctaccc	tatatatgct	gaaaaactac	cccctgaact	ctgagagtga	caatgtaacc	480
aactacatct	gtgtggtgcc	ttttaaagag	ctgggccttg	gacttagtga	agagcagatt	540
tcagaagagg	aagcacataa	ctttacagat	ggcttcagcc	tgctgcatt	gaaggttttg	600
ttccaactct	gggtggcaca	gagttcagag	ttcttcagac	ggttagccct	attactttct	660
acagccaatt	cacctctctg	gcccttactt	actccagcac	ttctgcctca	tcgtatctta	720
tctgatgtga	ctcaaggctc	acctcatgct	cattctgcct	gtttggaaga	gcttaagcgc	780
agctatgagt	tctatcggtg	ctttgaaact	cagcaccagt	cagtaccgca	gtgtttatcc	840
aaaactcaac	agaagtcaag	agaactgaat	aatgttcaca	cagcagtgcg	tagcttgag	900
ctccatctga	aagcattact	gaatgaggtg	ataattcttg	aagatgaact	tgaaaagctt	960
gtttgtacta	aagaaacaca	agaactagtg	tcagaggctt	atcccatcct	agaacagaaa	1020
ttaaagttga	ttcagcccca	cgttcaagca	agcaacaatt	gctgggaaga	ggccatttct	1080

35

cagggtcgaca	aactgctacg	aagaaatata	gataaaaaag	gcaagcctga	aatagcatgt	1140
gaaaacccac	attgtacagt	aagtaccttt	gaagcagcct	actctacaca	ttgcagacaa	1200
agatccaatc	ccagaggagc	aggaattaga	agcttatgta	gatgatatag	atattgatag	1260
tgatttcaga	aaggatgatt	tttattactt	gtctcaagaa	gacaaagaga	gacagaagcg	1320
tgagcatgaa	gaatccaaga	gggtgctcca	agaattaaaa	tctgtgctgg	gatttaaagc	1380
ttcagaggca	gaaaggcaga	agtggagca	acttctattt	agtgatcatg	tgtttcttca	1440
tatagcttta	aaattatgct	attgacatta	tgggaaagat	ttatcaatga	gagaaatgtg	1500
tctctttttc	agccgtgttg	aaatccttgt	ctcctgtaga	cccagtgga	cccataagta	1560
attcagaacc	atcaatgaat	tcagatatgg	gaaaagtcag	taaaaatgat	actgaagagg	1620
aaagtaataa	atccgccaca	acagacaatg	aaataagtag	gactgagtat	ttatgtgaaa	1680
actctctaga	aggtaaaaaat	aaagataatt	cttcaaatga	agtcttcccc	caaggagcag	1740
aagaaagaat	gtgttaccac	tgtgagagt	aagatgaacc	acaagcagat	ggaagtggct	1800
tgaccactgc	ccctccaact	cccagggact	cattacagcc	ctccattaa	cagaggctgg	1860
cacggctaca	gctgtcacca	gattttacct	tactgctgg	ccttgctgca	gaagtggctg	1920
ctagatctct	ctcctttacc	accatgcagg	aacagacttt	tggatgatg	gaggaagaac	1980
aaataataga	agaaaaataa	aatgagatag	aagaaaagta	agaaccaaga	ttcatatgaa	2040
gtgatattag	attgttcctt	ttacaaaagt	gttttagctt	aagactggaa	agggaaatag	2100
agtgtaaagt	tactatatat	aaagctaaga	tgtggattta	caggaagaac	cctgggtttga	2160
ataactgatc	tgaattagtg	agttacctgt	aaatggcaga	tcttttagga	aaataagaga	2220
aaggtaaagg	ctcttttgaa	taaactgctg	ttttatttgt	ggcacaactg	atcaatcttg	2280
gaaattcttt	aagtattttt	aataagaaat	gaattatcat	ttcttgccag	aatttgctac	2340
cttaagggtga	ttgggaaaat	tctgttgcaa	gaacattaac	atttagtatg	actccttttt	2400
actgtattct	tgcatgtaat	aactgcagct	attatgttaa	taacaagttg	tttgtatttt	2460
atttttggtt	ataccagctc	taaagatcca	ggttctgaat	aaaaaaatta	attgatacaa	2520
aaaaaaaaaa	aaa					2533

<210> 60
 <211> 899
 <212> DNA
 <213> Homo sapiens

<400> 60						
ggcagatttc	ccggcacctt	cgtgggcacc	acagagcccg	cctccccacc	cctgagcagc	60
acctcaccca	cactgctgc	ggccactatg	cctgtgtgtg	cctctgtggc	cagcctggcc	120
cctccggggg	aggcctcgct	ctgcctggaa	gaggtggccc	cccctgccag	tgggacccgc	180
aaagctcggg	tgctctatga	ctacgaggca	gccgacagca	gtgagctggc	cctgctggct	240
gatgagctca	tcactgtcta	cagcctgcct	ggcatggacc	ctgactggct	cattggccag	300
agaggcaaca	agaagggcaa	ggtccctgtc	acctacttgg	aactgctcag	ctaggcaggt	360
gcccccatcc	cccccgcat	ctggcctagg	caggagagga	tgggcgcact	gccacttaac	420
ttgtttgttg	gtgacacagt	tgttcagagt	ggggagaatt	caccccatc	tgtccctgcc	480
cctagtcaac	tagctgtgag	ggtgcctgag	gctgaatggc	tccaccctcc	cccagccctg	540
cttctgacct	tggtctctgg	agcccctgcc	cctgcctgca	tccccgagca	ccccaccctc	600
caggctccac	taaggaggga	ggggctgtct	gcagcagctg	cactcagcac	ctaggccagg	660
gtggggccgc	cgcatatggg	ctcaggaagc	cccaggtgca	ctcagcgaga	gccttgccct	720
tcagttgcc	aaagctgcat	caggggaatg	cggaaggca	cacagggtc	tggcagcccc	780
tggggactgg	gcgctgcccc	tgggagggga	gagcctggcc	agggctgggt	ttgggcccgg	840
agcagcatct	tccggtgcta	tcctcccctc	ccaccctca	cagctcaagc	caagtccag	899

<210> 61
 <211> 1079
 <212> DNA
 <213> Homo sapiens

<400> 61						
tcgacccacg	cgccgggtt	tcaccacgtt	ggccaggtg	gtctcaaaact	cctgacttca	60
gttggcctcc	caaagtgtg	ggattaaagg	catgagccac	tgcgcccggc	ctacctttct	120

aactctactt	ctagcttctt	gcttctgggc	tgtctgtata	ccaaacagga	atgtaatact	180
ttctgtcagc	ttcaggcctt	tgcacatgca	gttcactttg	tctatcttgg	tttttattct	240
taggatttta	attctcctaa	gaagctttct	ctgaccagcc	taaaacttac	gtaagccctg	300
ggttagggtg	tatgcttatg	tcctcccata	gcatttttgc	tttgcattgt	ttgtaactct	360
taatgtacag	catcatgatt	gcctatttta	actttctctg	ttgttacagt	agactttaat	420
ctcttttaagg	acaggaactg	tgtcttgttt	agaatcccca	gagcttattt	agtacatgg	480
ctatgcttat	aatttaagta	tttattgaac	aaatgaaatt	ttcctaagcc	ctaaaacctt	540
gcaagatggt	ttagtgagg	aaactggcct	cggtaggtt	gaataactag	cacgaggtca	600
ctcacctaaa	aagtggtag	gagggattaa	aatctaaatc	tgttttagctg	taaagattgg	660
gcttttttgc	ttgctgtgct	acatgactgc	ycctctcat	gttgctgtga	cacatccctg	720
tcaagtgttc	aaacagcccg	tgcctaacaa	ccccatccat	agcttctgag	gaaagtgtgtg	780
tcactctttg	acagctctga	gagctgaagc	gagtctttgc	agaataattt	cccatctatt	840
ggctcttaatt	tatgctttgg	agaatataac	ttattttcaa	aaaaaataatg	attcagaatt	900
tgtcatctcc	ttaaggtccg	tttattagtt	tatttcattc	cttcattcac	tgataaccat	960
ttactgagca	ccagcctggg	caacatggtg	agaacccatc	tctaccaatt	taaaaaaaa	1020
aaaaaagggc	ggcgcgtcta	gaggatccaa	gcttacgtac	gcgtgcatgc	gacgtcata	1079

<210> 62
 <211> 1928
 <212> DNA
 <213> Homo sapiens

<400> 62						
ggcacgagag	taggtctgcc	ggcgatggag	tggtgggcta	gctcgccgct	tcggctctgg	60
ctgctgttgt	tcctcctgcc	ctcagcgag	ggccgccaga	aggagtcagg	ttcaaaatgg	120
aaagtattta	ttgaccaa	taacaggtct	ttggagaatt	acgaaccatg	ttcaagtcaa	180
aactgcagct	gctaccatgg	tgtcatagaa	gaggatctaa	ctcctttccg	aggaggcatc	240
cccaggaaga	tgatggcaga	ggtagtcaga	cggagagctag	ggaccacta	tcagatcact	300
aagaacagac	tgtaccggga	aatgactgc	atgttccct	caaggtgtag	tggtgttgag	360
cactttat	tggaagtgat	cggcgctctc	cctgacatgg	agatgggtgat	caatgtacga	420
gattatcctc	aggttcctaa	atggatggag	cctgccatcc	cagtcttctc	cttcagtaag	480
acatcagagt	accatgatat	catgtatcct	gcttgacat	tttgggaagg	gggacctgct	540
gtttggccaa	tttatcctac	aggtcttggg	cgttgggacc	tcttcagaga	agatctgtta	600
aggtcagcag	cacagtggcc	atggaaaaag	aaaaactcta	cagcataatt	ccgaggatca	660
aggacaagtc	cagaacgaga	tcctctcatt	ctctctgtctc	ggaaaaaccc	aaaacttgtt	720
gatgcagaat	acacaaaaa	ccaggcctgg	aaatctatga	aagatacctt	aggaaagcca	780
gctgctaagg	atgtccatct	tgtggatcac	tgcaaataca	agtatctgtt	taattttoga	840
ggcgtagctg	caagtttccg	gtttaaacac	ctcttctctg	gtggctcact	tgttttccat	900
gttggtgatg	agtggctaga	attcttctat	ccacagctga	agccatgggt	tcactatata	960
ccagtcaaaa	cagatctctc	caatgtccaa	gagctgttac	aatttgtaaa	agcaaatgat	1020
gatgtagctc	aagagattgc	tgaaggggga	agccagttta	ttaggaacca	tttgcagatg	1080
gatgacatca	cctgttactg	ggagaacctc	ttgagtgaat	actctaaatt	cctgtcttat	1140
aatgtaacga	gaaggaaagg	ttatgatcaa	attattccca	aaatgttgaa	aactgaacta	1200
tagtagtcat	cataggacca	tagtccctct	tgtggcaaca	gatctcagat	atcctacggt	1260
gagaagctta	ccataagctt	ggcacctata	ccttgaatat	ctgctatcaa	gccaaatacc	1320
tggttttctt	tatcatgctg	cacccagagc	aactcttgag	aaagatttaa	aatgtgtcta	1380
atacactgat	atgaagcagt	tcaacttttt	ggatgaataa	ggaccagaaa	tcgtgagatg	1440
tggattttga	acccaactct	acctttcatt	ttcttaagac	caatcacagc	ttgtgcctca	1500
gatcatccac	ctgtgtgagt	ccatcactgt	gaaattgact	gtgtccatgt	gatgatgccc	1560
tttgtcccat	tatttggagc	agaaaattcg	tcatttggaa	gtagtacaac	tcattgtctg	1620
aattgtgaaa	ttattcaagg	cgtgatctct	gtcactttat	tttaattgtag	gaaaccttat	1680
ggggtttatg	aaaaatactt	ggggatcatt	ctctgaatgg	tctaaggaa	gggtagccat	1740
gccatgcaat	gatgtaggag	ttctcttttg	taaaaccata	aactctgtta	ctcaggaggt	1800
ttctataaat	ccacatagaa	agaggccaat	tgcatgagta	attattgcaa	ttggatttca	1860
ggttcccttt	ttgtgccttc	atgccctact	tcttaatgcc	tctctaaagc	caaaaaaaa	1920
aaaaaaaa						1928

<210> 63
 <211> 781
 <212> DNA
 <213> Homo sapiens

<400> 63
 ggcacgagat tttcagcctt tttggactgg tttctccaca tcttcgtgga tttatctaac 60
 tttggtcttt gatgttggtg accttcagat tgggtctctg agtgaacatc ctttttgttg 120
 atgttgatac tattcctttc tgtttggttg tttgttttcc ttctaacagt cagggccctc 180
 tgctgcaggc ctgctggagt ttgggtgagg tccactccag accctgtttg tctgggtttt 240
 gccagaggag gctgcagaat agcaatgatt gctgcctgtt tttcctctgg aagctttgtc 300
 ccagaggggc acccaccaga tgcagccag agctctcctg tatgagggtg ctgttgccc 360
 atacttgagg gtgccttcca gtcaggatac acagggtgtc ggtaccactc tgaggaggca 420
 ctctgtcccc tatcagagct cgaacactgt gctgggagat ccactgttct cttcagagct 480
 gtcagacagg gacgtttaag tctgctgaag ctatgcccac agctgccctt tccccagat 540
 gctctgtccc agggagaagg gaggttttatc tataagtctc tgactggggc tgctgccttt 600
 tcttcagaga tgccctgccc caagacgggg actctagaga ggcagtctgg ctgcagtggc 660
 cttgctgaac tgtggtgggc ttcacccagt tggaccttcc ctgagccttt ttttttacc 720
 tgtgagggta aaaatgccta atcaagcctc agcaatgggtg gatgcccttc cccccacaa 780
 g 781

<210> 64
 <211> 1194
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1172)
 <223> n equals a,t,g, or c

<400> 64
 ggcacgagaa gacatggagt cttaagtgtg atcagtggga gggggctgga atcatttaga 60
 ggcacatctca ttcacaaaac caggagctga tactggctgt cagccaggac ttcaactgac 120
 ctatgtagaa cctgtccatg tggccccctc ttgcagtctc cccatttggg ctggtttggg 180
 cttcatcaca gtccggcagc ttacttctaa gggcaagcat tccacgacaa cacagcagaa 240
 gggcatggca tttttacagt gaagtttggc aatctcatag cgtcgttctt gtcctacttt 300
 atttattggt cagggcaatc acaaagatgt gcataggctc aaagaaaaga gacataacct 360
 cgaccacgcy atggaagaag tgacacggtc atgttatgag aggagtgtgt gggatgggag 420
 atagggctgt ggccacctgc agaaaacagc atctgctata ggctgtcatg gaagcgcagg 480
 atggggattt agcctacctg aggggtcagt cagcaaaggc ctctgggagg aagttagatc 540
 ttcggctgag gatgtgaagg gctaaaagga gaatgaggaa gagtttcagg gagaggaaac 600
 aatgaaacga gtccagagac gctggtgagt tggatggttt gcttcagtat gatgacaata 660
 cagaggggca aggagactgg tgcaggagaa gagagaaggc gccatgtgct ctgggtcgtg 720
 tcttctatgc cagactccct tagaagagga gcagcctcca gtcagcgggtg tcccaggaac 780
 acggaggcta gacaggacaa tggcagccaa tcctgtctcc caaactgggt acagtgggga 840
 aaagctgcat ggtctagatc caccctgctc cctggcccca gtatagaaga tcaaattcaa 900
 tctgcccatt cttatccaga taaagtaaag gaagactgga aaaaagaact aatccacggc 960
 tccatctgcc catgactttc tctgctgatg ccggaggcag ctatggataa agagacggca 1020
 cacggcatgt cccgacgctg tggaggtggg gagacccgc aagtccacag gaaaagagtt 1080
 aagttgctgc cacctgggca tccgctatcc tctgctcttc tgctcatcc tcaattcaga 1140
 ccatgatgga gctgattgtc tcccatttta tnccttggat tgaatggctc cgag 1194

<210> 65
 <211> 1677

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1012)

<223> n equals a,t,g, or c

<400> 65

ggtgcagtgg	tgccatcaca	gttcaactgca	gccttgacct	ccggggctca	agcaatcctc	60
ccacctcagc	cacttgagta	gctgagacct	cagatatgtg	ccatcacacc	cagctgattt	120
tttaaaatta	atTTTTtGta	gagatagggt	ctcatatgtt	gcccattgtg	gtctcaaaact	180
actgggttca	aatgatcctc	ctgcctcagc	cttccaaagt	actgggatta	caggcatgag	240
ccaccatgcc	gggctgggag	gcggaatttt	gttcagtcta	aagataagct	ttttcatagc	300
tctggctgta	gtgggaggga	gcagaggagt	gaatgattgt	cagttgggag	ggtgcagagt	360
gggctcctgc	cctagggtgr	aggtragggt	ggcttaggtg	asmcamcaca	gaggcctgt	420
tcagcccccac	gtcccctccc	tgtgctccct	cctcctctct	cctcctcctgc	aggcgtggga	480
ggtatcatca	ttcagcagat	ttcaccagag	gcagtggagg	aggcaggtag	ctgagccaga	540
attcagaatg	tcttattctc	cacttgactc	tgccactaac	ttgtttgtgca	actttgggccc	600
tttccccagg	ccttcatttt	cttttctttt	ctttttcttt	yttttttttt	gaggcggagt	660
ctcgctatgt	tgcccaggct	ggagtgcagt	ggcgcagcat	catctcggct	cactgcaagc	720
tocaccttct	gagttcacgc	cattctactg	cctcagcctc	ccgagtagcc	gggactgcag	780
gorccccacca	ccacgcccgg	cttatttttt	gtattttttag	tagagacagg	gtttcaccac	840
gttagccaag	atggtctcga	tctcctgacc	tcgtgatcca	ccgcctggg	cttcccaaaag	900
tgctgggatt	acaggcgtga	gccactgcgc	ccggccattt	tcttaaatat	ctaataaaaa	960
atatatagca	aatgcagttt	ttaaactacg	acaatatgac	cacgcaaaag	antattatct	1020
tccaagactg	ctgggtccaag	gaaaagtcag	taataaagtg	gaagcattgt	agcttatgga	1080
atgactgggt	asatttgga	gaagccttag	caataatcta	gaatctgcat	agataatata	1140
tctgaggatt	gggctttgtg	gtttacaaaag	catttttttt	tcctcttttg	atoccagccg	1200
tttgtctgga	ctgatacaaa	gcatttttat	tagtttgtct	tattcaatcc	tcacaccacc	1260
tcaaatttac	agaggatatg	gatctgggtta	atttgtatga	ctatgtaacc	tcattgtcagt	1320
ccacagcact	gcctggagggt	gggtagagggt	ggctcctgggc	tggaatccca	gccccagtg	1380
gaccttgagc	aagttacttt	agctgtctgc	acctaaattt	cctcactggc	aaaacaggaa	1440
tactgtgggt	tcacacctgc	aattccagca	ctttgggagg	ctgagggtggg	aggattgctt	1500
gagtcagaa	gttcaaaacc	agactgggca	acatagcaag	accatctcta	caaaaattaa	1560
ataaatataa	cattttacaag	ggttgtgtgtg	aagattaaat	gagatcactc	acgaaaaaagc	1620
tcagcagacc	ctgatgtgca	gtaggtgctc	aataaatgtt	agccagcaaa	aaaaaag	1677

<210> 66

<211> 1237

<212> DNA

<213> Homo sapiens

<400> 66

agcaaaccca	ggaagggtgtg	gcgtccccgc	ttcgcgccaa	gatgggtgctg	gtgctgcgcc	60
atccttttgtg	tgcccgggaa	agggcggttc	gggagccggg	tggggggctc	ctgactcgca	120
ctgggcagca	tgacgggtgcg	ccggctgtca	ctgctgtgcc	gggacctctg	ggcgctgtgg	180
ctgctgctga	aggccggcgc	agtgcgtggg	gcgcggggcg	gtcctcgctc	ccccggaagg	240
tgttgtgggg	cgacatgcgg	ggacgcgggg	cggggggtgga	cgttctgggc	ccagccctgt	300
cctcagaagc	tgctggggca	gaagcccggg	gctgggggat	gccgggggatg	ggtgttgggg	360
tgggtgcctc	cgagaccaga	ggagccctgt	tccttgccag	ggaagggtgtg	cacgggcctt	420
gcccgatgga	tggttttaggg	ccatggccct	ggggtccctg	gtgagcagtg	gggcccctc	480
tgcccttgcc	ctgtgaggga	ctgtctgtgc	tggtcccaga	aggctgggat	cacctttcca	540
ctggctcctt	tgttcgaggt	ttttcataga	caggctatgt	ggacaaatga	gggcagcgcc	600
cacgtctggc	tgggtggagg	gctgcggctc	ctccttgagg	gggacgcctg	gccactgctg	660
tcccacaat	ggggccaccc	gtgggtgcaag	gcgtgacaag	ctgcctctc	taggtaagca	720
ggacttggga	ggcccctggc	caagcctgtg	gacccggctg	ggcggcctct	gtgggtctcag	780

39

gtttgggtgt	gtttgggtctg	gtcagggctc	aggggctgct	ggtccacact	ggccccatcc	840
tgacaattgg	agctttgggg	caaggtccct	ggagaagggg	tcacgtcggg	aggaaacagc	900
ctgggttttg	ttgatgcttt	tctaagaatg	gagtactcgt	tttcaagaga	tttgtcctaa	960
ttatatatttc	cagcgggtac	ttatgccaag	tattgatgaa	taattcataa	aataagcatc	1020
tttggtgaatt	ttagtgaatc	agaccttaac	tatcaacggc	aatgaatgaa	catctaaagt	1080
ttccaatttt	aaagtaaaga	actggctggg	tacagcagtt	cacgcctgta	atcccagcac	1140
tttgggaggc	caaggctaga	ggatcgcttg	agcccaggag	tttgagatca	gcctggggcaa	1200
cataccaaga	cctcatctgt	taaaaaaaaa	aaaaaaa			1237

<210> 67

<211> 1934

<212> DNA

<213> Homo sapiens

<400> 67

ccacgcgtcc	ggggcggtcc	tggctgtgag	aggggagccc	caggggagct	ggggcagcat	60
gactgggggtg	ataaatggcc	ggaattttgg	cgtggccaca	ctcaacacca	gcgtgatgca	120
ggaggcacac	tccggggtca	gcagcatcca	cagcagcatc	cgccatgtcc	cagcaaacgt	180
ggggcctctg	atgcgggtgc	tcgtggtcac	catcgccccc	atctactggg	ccctggccag	240
agagagtggtg	gaagccctga	atggccactc	tctgactggg	ggcaagttcc	ggcaggagtc	300
acacgtggag	tttgctacag	gggagctgct	cacgatgacc	cagtggcccg	gggtctggat	360
cccgatggcc	tcctgctcct	cgacgtggtg	gtcaatggcg	ttgtccccgg	acagcctggc	420
tgacgcagat	cttcaagtgc	aggactttga	ggagcactac	gtgcaaacag	ggcctggcca	480
gctgttcgtg	ggctccacac	agcgcttctt	ccagggcggc	ctccccctgt	tcctacgctg	540
caaccacagc	atccagtaca	acgcggcccg	gggccccccg	ccccagctgg	tgcagcacct	600
gcgggcctca	gctatcagct	cggcctttga	tccagaggcc	gaggccctgc	gcttccagct	660
cgctacagcc	ctgcaggcgg	aggagaacga	ggtcggctgc	cccgagggct	ttgagctgga	720
ctcccaggga	gcgtttttgt	tggatgtgga	cgagtgtgcg	tgggatgtc	acctctgccg	780
agagggacag	cgctgtgtga	acctgctcgg	gtcctaccgc	tgctcccccg	actgtggggc	840
tggcttccgg	gtggctgatg	gggcccggctg	tgaagatgtg	gacgaatgcc	tggagggggt	900
ggacgactgt	cactacaacc	agctctgcga	gaacacccca	ggcggtcacc	gctgcagctg	960
ccccaggggt	taccggatgc	agggccctag	cctgccctgc	ctagatgtca	atgagtgcct	1020
gcagctgccc	aaggcctgcg	cctaccagtg	ccacaacctc	cagggcagct	accgctgcct	1080
gtgcccccca	ggccagaccc	tccttcgcga	cggaaggcc	tgcacctcac	tggagcggaa	1140
tggacaaaat	gtgaccaccg	tcagccaccg	aggccctcta	ttgccctggc	tgcggccctg	1200
ggcctcgatc	cccggtacct	cctaccacgc	ctgggtctct	ctccgtccgg	gtcccatggc	1260
cctgagcagt	gtgggcggg	cctgggtgcc	tcctggtttc	atcaggcaga	acggagtctg	1320
cacagacctt	gacgagtgcc	gcgtgaggaa	cctgtgtcag	cacgcctgcc	gcaaacactga	1380
gggcagctac	cagtgcctgt	gccccgcgg	ctaccgtctg	ctccccagcg	ggaagaactg	1440
ccaggacatc	aacgagtgcg	aggaggagag	catcgagtgt	ggacccggcc	agatgtgctt	1500
caacacccgt	ggcagctacc	agtgtgtgga	cacacctgt	cctgccacct	accggcaggg	1560
ccccagccct	gggacgtgct	tccggcgctg	ctcgcaggac	tgcggcacgg	gcggcccttc	1620
tacgtgcag	taccggctgc	tgccgctgcc	cctgggctg	cgcgccacc	acgacgtggc	1680
ccgcctcacc	gccttctccg	aggtcggcgt	ccccgccaac	cgacccgagc	tcagcatgct	1740
ggagcccgac	ccccgcagcc	ccttcgcgct	ggtccgctg	cgcgccggcc	ttggcgcggt	1800
ctacacccgt	cgcgcgctca	ccccgcggcg	cctctaccgg	ctcaccgtgc	gtgctgcggc	1860
accgcgccac	caaagcgtct	tcgtcttgct	catcgccgtg	tccccctacc	cctactaaac	1920
gggagagggc	attg					1934

<210> 68

<211> 3300

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (3)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (15)
 <223> n equals a,t,g, or c

<400> 68
 ncngcagccg gacgnccgag cgcagcgagt cagtgcgcga ggaagcggaa gagcgcccaa 60
 tacgcaaacc gcctctcccc gcgcgttggc cgattcatta atgcagctgg cacgacaggt 120
 ttcccgaactg gaaagcgggc agtgagcgca acgcaattaa tgtgagttag ctcaactcatt 180
 aggcaccccca ggctttacac tttatgcttc cggctcgtat gttgtgtgga attgtgagcg 240
 gataacaatt tcacacagga aacagctatg accatgatta cgccaagctc gaaattaacc 300
 ctcaactaaag ggaacaaaag ctggagctcc accgcggtgg cggccgctct agaactagt 360
 gatcccccg gctgcaggaa ttcggcacga gaacacatct taagggaacc aagtctcaag 420
 agaaaatcaag taattatgaa tgaacagctc taaaaaagag agagagaata ttttcttaaa 480
 tcaacttagt tgctgttatg accaaagaac agatgtttgt gtgttcaccc cagagaagca 540
 agagattttc ccttaaacct cagcttataa tgaatggaag tgaatgacag ggagagaggt 600
 tttctctcgt tccccagaac tctatccttt tcttctcaaa cagttggaaa ctgtagccaa 660
 tacagtagac agtgatatgg gagaaccaa tctgcatcca agcatgtttc tcttactttt 720
 ggtgttggag agactctacg cttccccgat ggatgggtact tcttctgctc tcagcatggg 780
 accttttgtt cccttcatta tgaggtgtgg tcactcacct gtctaccact cccgtgaaat 840
 ggcaagctcgt gccttggtec catttgttat gatagatcac attcctaata ccattcgaa 900
 tctgttgttc acactcccca gctgcactga ccagtgtttc cggcaaaacc acattcatgg 960
 gacacttctc caggtttttc atttgttgca agcctactca gactccaaac acggaacgaa 1020
 ttccagacttc cagcagcagc tgactgacat cactgtttgt accaaagcca aactctgggt 1080
 ggccaagagg caaaatccat gtttgggtgac cagagctgta tatattgata ttctcttct 1140
 attgacttgc tgccctcaaca gatctgcaaa ggacaaccag ccagtctctg agagtcttgg 1200
 cttctgggag gaagtcagag ggattatctc aggatcagag ctgataacgg gattcccttg 1260
 ggccctcaag gtgccaggcc tgccccagta cctccagagc ctaccagac tagccattgc 1320
 tgcagtgtgg gccgcggcag ccaagagtgg agagcgggag acgaatgtcc ccattctctt 1380
 ctctcagctg ttagaatctg ccttccctga agtgcgctca ctaacactgg aagccctctt 1440
 ggaaaagttc tttagcagcag cctctggact tggagagaag ggctgtgccac ccttgcctgt 1500
 caacatggga gagaagttct tattgttggc catgaaggaa aatcaccag aatgtctctg 1560
 caagatactg aaaattctcc actgcatgga ccctgggtgag tggcttcccc agacggagca 1620
 ctgtgtccat ctgaccccaa aggagtctct gatctggagc atggatattg cttccaatga 1680
 aagatctgaa attcagagtg tagctctgag acttgcttcc aaagtcattt ccaccacat 1740
 gcagacatgt gtggagaaca gggaattgat agctgctgag ctgaagcagt gggttcagct 1800
 ggtcatcttg tcatgtgaag accatcttcc tacagagtct aggttggccg tcgttgaagt 1860
 cctcaccagt actacaccac ttttctccac caaccccat cctattcttg agttgcagga 1920
 tacacttgct ctctggaagt gtgtccttac ccttctgcag agtgaggagc aagctgttag 1980
 agatgcagcc acggaaccg tgacaactgc catgtcaca gaaaatacct gccagtcaac 2040
 agagtttgcc ttctgccagg tggatgcctc catcgctctg gccctggccc tggcgtctct 2100
 gtgtgatctg ctccagcagt gggaccagtt ggcccctgga ctgccatcc tgctgggatg 2160
 gctgttggga gagagtgatg acctcgtggc ctgtgtggag agcatgcac aggtggaaga 2220
 agactacctg tttgaaaaag cagaagtcaa cttttgggcc gagaccctga tctttgtgaa 2280
 atacctctgc aagcacctct tctgtctcct ctcaaaagtc ggctggcgct cccaagccc 2340
 tgagatgctc tgtcaccttc aaaggatggt gtcagagcag tgccacctcc tgtctcagtt 2400
 cttcagagag cttccaccag ctgctgagtt tgtgaagaca gtggagttca caagactacg 2460
 cattcaagag gaaaggactt tggcttgctt gaggctgctg gccttttttg aaggaaagga 2520
 aggggaagac accctagttc tcagtgtttg ggactcttat gcagaatcga ggcagttaac 2580
 tcttccaaga acagaagcgg catgttgaag aaaatctggg ggattgggat ggggggtatgt 2640

41

gtggattttt	cctccactaa	atctgcagga	aacatgttga	acataaatc	aaaaatttta	2700
tcccaaaaaa	aaaaaaaaa	aaactcgagg	gggggcccgg	taccaattc	gccctatagt	2760
gagtcgtatt	acaattcact	ggccgtcgtt	ttacaacgtc	gtgactggga	aaaccctggc	2820
gttaccacaac	ttaatcgcc	tgcagcacat	ccccctttcg	ccagctggcg	taatatgcga	2880
gaggcccgca	ccgatcgccc	ttcccaacag	ttgcgagcc	tgaatggcg	atggcaaatt	2940
gtaagcgta	atattttgtt	aaaattcgcg	ttaaattttt	gttaaatcag	ctcatttttt	3000
aaccaatagg	ccgaaatcgg	caaaatccct	tataaatcaa	aagaatagac	cgagataggg	3060
ttgagtgttg	ttccagtttg	gaacaagagt	ccactattaa	agaacgtgga	ctccaacgtc	3120
aaagggcgaa	aaaccgtcta	tcaggggcat	ggcccactac	gtgaaccatc	accctaatac	3180
agtttttttg	ggtcgagggtg	ccgtaaagca	ctaaatcgga	accctaaagg	gagccccga	3240
tttagagctt	gacggggaaa	gccggcgaa	gtggcgagaa	aggaaggga	gctgtctctt	3300

<210> 69

<211> 1797

<212> DNA

<213> Homo sapiens

<400> 69

ggtcgacggt	atcgataagc	ttgatatcga	attcctgcaa	cagttcttgg	aaaccactc	60
gagagggcca	cgctccatt	caccaggcca	cgcatcaca	gaggcaacac	caggagccaa	120
catgagctcg	gggactgaac	tgctgtggcc	cggagcagcg	ctgctggtgc	tgttgggggt	180
ggcagccagt	ctgtgtgtgc	gctgtcacg	cccagggtga	aagaggtcag	agaaaatcta	240
ccagcagaga	agtctgcgtg	aggaccaaca	gagctttacg	gggtcccgga	cctactcctt	300
ggtcggggcag	gcatggccag	gaccctggc	ggacatggca	cccacaagga	aggacaagct	360
gttgcaattc	tacccagcc	tggaggatcc	agcatcttcc	aggtaccaga	acttcagcaa	420
aggaagcaga	cacgggtcgg	aggaagccta	catagacccc	attgccatgg	agtattacaa	480
ctggggggcgg	ttctcgaagc	ccccagaaga	tgatgatgcc	aattcctacg	agaatgtgtc	540
catttgcaag	cagaaaacca	cagagacagg	tgcccagcag	gagggcatag	gtggcctctg	600
cagaggggac	ctcagcctgt	cactggccct	gaagactggc	cccacttctg	gtctctgtcc	660
ctctgcctcc	ccggaagaag	atgaaggaat	ctgaggatta	tcagaacttc	agcattccat	720
ccattcagtg	gcgcgagtc	aggaaggtca	tggggcaact	ccagagaaga	aagcatcccc	780
tggcccggtg	ggaagcccag	acgaggagga	cggggaaccg	gattacgtga	atggggaggt	840
ggcagccaca	gaagcctagg	gcagaccaag	aagaaaggag	ccaaggcaaa	gagggaccac	900
tgtgtcatg	gaccatcgc	tgccttccaa	ggaccatttc	ccagagctac	tcaactttta	960
agcccctgcc	atggttgctc	ctggaaggag	aaccagccac	cctgaggacc	acctggccat	1020
gcgtgcacag	cctgggaaaa	gacagttact	cacgggagct	gcaggccccg	tcaccaagcc	1080
ctctcccag	ccaggctttg	tggggcaggc	acctggtacc	aagggtaacc	cggctcctgg	1140
tatggacgga	tgcgcaggat	ttaggataag	ctgtcaccca	gtccccataa	caaaaccact	1200
gtccaacact	ggtatctgtg	ttcttttgtg	ctatgaattt	ggattcctaa	ttgctattgt	1260
tggttgctgg	ggttttaaat	gattgataag	cttgtacagt	taacttatag	agggggagcc	1320
atatttaaca	ttctggattt	cagagtagag	atttctgtgt	tgtctcctag	aaagcattac	1380
atgtagttta	tttcagcatc	cttgttgggt	ggggccctgg	ctctcttccc	ctttgggtggg	1440
acctccccct	tctttgggct	tcagttcact	caggaagaaa	tgaggctgtc	gccatcttta	1500
tgtgcttcca	gtggaaatgt	cacttgctac	agacaatagt	gcatgagagt	ctagagaagt	1560
agtgaccaga	acagggcaga	gtagggtccc	tccatggccc	tgaatcctcc	tctgtccag	1620
ggctggcctc	tgcagagctg	attaacacgt	gttgtgactg	tctcatggga	agagctgggg	1680
cccagagggg	ccttgagtca	gaaatgttgc	cagaaaaagt	atctcctcca	accaaaacat	1740
ctcaataaaa	ccatttttagt	tgaaaaaaaa	aaaaaaaaa	aaaaaaaaa	aaaaaaa	1797

<210> 70

<211> 1373

<212> DNA

<213> Homo sapiens

<400> 70

ggcacgaggg	ctgacggcgc	ttttgtctcc	ggtgagtttt	gtggcgggaa	gcttctgcgc	60
------------	------------	------------	------------	------------	------------	----

tggtgcttag	taaccgactt	tcctccggac	tcctgcacga	cctgctccta	cagccggcga	120
tccactcccc	gctgttcccc	cggaggtcca	gaggcctttc	agaaggagaa	ggcagctctg	180
tttctctgca	gaggagtagg	gtcctttcag	ccatgaagca	tgtgttgaac	ctctacctgt	240
taggtgtggt	actgacccta	ctctccatct	tcgttagagt	gatggagtcc	ctagagggct	300
tactagagag	cccctgcctt	gggacctcct	ggaccaccag	aagccaacta	gccaacacag	360
agccacacaa	gggccttcca	gaccatccat	ccagaagcat	gtgataagac	ctccttccat	420
actggccata	ttttggaaca	ctgacctaga	catgtccaga	tgggagtccc	attcctagca	480
gacaagctga	gcaccgttgt	aaccagagaa	ctattactag	gccttgaaaa	acctgtctaa	540
ctggatgctc	attgcctggg	caaggcctgt	ttaggcctgt	tgcgggtggc	catgcctgta	600
atcctagcac	tttgggaggg	tgatgtgggt	ggatcacctg	aggtcaggag	ttcagaccag	660
cctcgccaac	atggcgaaac	cccatctcta	ctaaaaatac	aaaagttagc	tgggtgtggt	720
ggcagaggcc	tgtaatccca	gctccttggg	aggctgaggg	gggagaattg	cttgaacccg	780
gggacggagg	ttgcagttag	ccgagatcgc	actgctgtac	ccagcctggg	ccacagtga	840
agactccatc	tcaaaaaaaa	aagaaaagaa	aaagcctgtt	taatgcacag	gtgtgagtgg	900
attgcttatg	gctatgagat	aggttgatct	cgccttacc	ccggggtctg	gtgtatgctg	960
tgctttcctc	agcagtatgg	ctctgacatc	tcttagatgt	cccaacttca	gctgttggga	1020
gatggtgata	ttttcaaccc	tacttcttaa	acatctgtct	ggggttcctt	tagtcttgaa	1080
tgtcttatgc	tcaattatgt	ggtgttgagc	ctctcttcca	caagagctcc	tccatgtttg	1140
gatagcagtt	gaagagtgtg	tgggtgggct	gttgggatga	gatggagtgt	tcagtgcaca	1200
tttctcattt	tacattttta	agtcgttctc	ccaacatagt	gtgtattggt	ctgaaggggg	1260
tggtgggatg	ccaaagcctg	ctcaagttat	ggacattgtg	gccaccatgt	ggcttaaatg	1320
attttttcta	actaataaag	tgggatatat	atttaaaaaa	aaaaaaaaaa	aaa	1373

<210> 71

<211> 1579

<212> DNA

<213> Homo sapiens

<400> 71

ggcacgagga	tttggagggg	acaaacatcc	aaaccattta	agtcacagca	ctttactccg	60
cagtgtgaat	aacacaggca	ttctcctaca	taatcacagt	acagttaatca	tactctggaa	120
attgaatatc	atctaataata	ctttccatac	ccagattttc	ttagattttcc	caatgatatt	180
tcttactgtc	ctcccttttag	ccttccctctt	tctccattca	ggatttctacc	attacatttc	240
attttcatgt	ctcttcagtc	tctcttttagc	tttgtttttc	tttcttgatg	ttgccacttt	300
taggaggcca	ggccagttgt	tttgtgaaag	atctgttctc	tttgatatgt	ttcatttttg	360
atttgtttca	ttgtttttgc	atgaatggat	tcaggctaaa	cattttttggg	caggactggt	420
tattgtatta	cctagtgatg	tgttcttttc	agtcacatcat	ctggaggcac	ctgatggcag	480
ttttcccaat	attgcgaaat	taagtttgat	tattttgtta	aggtagtgtc	caccagatct	540
ctccatttta	aagacatcct	tttctctaata	tactcagtgg	actgtagagt	gatgctttga	600
aactgaataa	ctaacaactcc	ctaactcagt	gatttagcac	ccgttgattt	tttttttttg	660
cctgaatcaa	atattattat	agtagtttta	aatggtgatt	ttccatttct	attcttttgt	720
tagctgccat	tcttctataa	ttttgtcttt	atattttact	gggtgttaag	attgctattc	780
cagctttcct	ttgtctttta	caccttttcc	cattctttta	tttttttcca	tccttttttg	840
tcctgttttc	caatagatgg	atagaatttt	ctttctctgg	tttaaagggt	atacatttgt	900
gtgtgtgtgt	gtattctaaa	ccatttggcc	ttaaaacata	gagatggtta	ttcctgttga	960
ttaaaaaaa	ctcagtaatg	ttactatctt	ctctttaata	agataggta	tttatttcat	1020
tctgtgttct	tgggtgaggt	ctccccctcac	ccagtcgaagt	tgatgttaat	ctagaatttt	1080
tagtttttta	aattatcgat	atactttctg	tcttttctct	tttttttcac	tctctctgta	1140
tttgtgcttt	ccccctttac	tctctttccc	ttcattctct	ctttctcact	ttcttcttct	1200
atttttcact	cttgggtaga	ttatctttta	gaaaacagca	agatattata	taatttactt	1260
tatatctctc	ttccaaatga	ttaaagtaat	aattaaaaat	ttttgatatg	tgtgatgca	1320
aggataggaa	tcctcttgta	agtggaagaa	ctctaccaca	tgcatgagtc	attagtgtgt	1380
taaacactgg	gaagtggctt	taggtccagc	tgggtgctct	gaagaaggta	ggtttcttca	1440
gttcttttatg	ttaaactgtc	ccttacctta	aaaaaggagt	gaagaatact	gactgcagag	1500
gttttgtgag	gattccggta	acacagaagc	atagaactgg	aaaagaatat	taatatattgc	1560
caaaaaaaaa	aaaaaaaaa					1579

<210> 72
 <211> 1028
 <212> DNA
 <213> Homo sapiens

<400> 72
 gcaacgacaat tgaactgaac cctaaaaatg ctacttcaat tcaccttatg ggtatttgggt 60
 gctatacatt tcccgaatg ccttgggtatc aaagaagaat tgctaaaatg ctgtttgcaa 120
 ctgcctccta gttccaccta tgagaaggta gtatgatgtc ctttggttaag ttagtacgga 180
 ttctctgaac cacagcgccc attctaccat gtgttccaca cattgtggag ctctggattc 240
 agtgaagggg acttgaggca atttccttaa cgatccaatt caactgtgtt atcacaaggc 300
 ttaacactta ttatccttga ctgggtgagt gttttctttt tccccgttag gtgagtggtc 360
 ggtaattctg gaatactgtc atctaaaatg gctcgtggct aaaatctacc ttcattttct 420
 gtttgaaatc taaactatat tgaagtcata aaatagaaca agaaatacag catctgttac 480
 ccagcatgtt ttagctgtat tacacacaat aacagaaaag taaagcagat gcttaagttg 540
 ataaaagaag aacactcatt ataacttcta ttttaaaaag catatgaaag gttcatattc 600
 tctcatattt tcaaggcctt ttgcttttct tgttaaaaat aagatttgag aggaatttct 660
 ggtttaaact tgggtttact catcacaagc ttttcagagt aagaaaacag gcaatcgaaa 720
 aagctgtact tgtattattt acattataac aaggagcctt tttttcttct tgggaagcta 780
 tagtgtagaa attgatgtaa aaaatactta gttgtattct ttacacacag ttgagaaata 840
 ttattaaaat aatgcaccaa tatttataaa tgggtattatt aaaaataatgc ccatttgctg 900
 gacacgggtg ctcatgcctg taatgccagc attttggaaag gccaaaggtg gtggatcagt 960
 tgagcctggg agtttgagac cagcctgggc aacgtggcaa aacccttgca aaaaaaaaaa 1020
 aaaaaaaaaa 1028

<210> 73
 <211> 3674
 <212> DNA
 <213> Homo sapiens

<400> 73
 ggcacgagct caaaagaaat aggggtgattt ttaaaggatt aataaaattc tgaatgtta 60
 agtagaagat tacattgtct agtcttgtat ttctctcttc tgttgccttc tttcattcac 120
 acactctcag tttctcatat ttgtagtcca tttatttggg tatttccctaa gaattattgaa 180
 agtgaagcaa ctatgtgact gtattcttca ggtaaacact gactgcgctt gttggatttt 240
 cctatttttt gtgacttcaa gaataatatg ccctgctgaa tacatgccat ttcacattct 300
 gaaactgggt agagtgggtt ggtgttctgc caacaattgc tagtgggtgtg aattcattca 360
 tatttgccag tattgctcac ttcaaagaaa ctcccttcatc aagcagcca gagctaggcc 420
 agatcaatgc tacaatcatg aagttctcat tgcattgcaat tgtgtaggat tgacaaggaa 480
 ctcagataaa aatttccagg gtgcacttcc agaaccagct tcaacatatg tctacattgc 540
 cccaagtta ataaagtgcc aaccctttac tctctcatc agccagaaat gttagaaatc 600
 caaaatcttg gtgcattatt ttttcataaa cgctaaaaca tttgaagaaa caatttaatt 660
 atttaaaatt caagtatttt attcacatta tttgcaatat ccaaatgttt aaaaattccc 720
 agataattaa ctagctatta cagatctcac ctagaggggt gatgttatga agactccagt 780
 ggactgtact cacaaattga ctggacaccc tatgaaagtg ggtagacctc tcagcggaaa 840
 ataagaaggg cttttaccta cagggcagga cagggtccca tgagagcagt tctgtggaga 900
 tataaaaaga atggaagaag gaatgcctta tagtgatatt gtgacattat atctatatat 960
 ctacacatat ctatctatct atatctacat ctatataatc ttacatttaa aattgtattc 1020
 ctacacatat tagaaactct tctaataaat gaagtaaaaa aattaaaaag aatacaaaata 1080
 ttccagcccc aaatgagaaa tcaaacatat taaaattgtt caagaaaatt tctttgaaca 1140
 cttctgaaag tttttggaaa cttagaaaag agggaaaaaa atccagtgtt actagtaatt 1200
 tccatggtaa tacagataaa atacattctt ttaattctgg gaaattagaa aaagtggggg 1260
 gatctttcca ggaaaaacat gtgtaacatc tgcttatcac tccagctccc tccctcctct 1320
 cctctccacg ttcccttgag taaatgtctg ggaagcatg aagcttgatg caagaaccct 1380
 gttgtactgg cgttttcctc ccctgtgaaa acgttaactac tgttgggagt gaactgagga 1440
 tgtagaaagg tgggtggaacc aaattgtggt caatggaaat aggagaatat ggttctcact 1500

cttgagaaaa	aaacctaaga	ttagcccagg	tagttgcctg	taacttcagt	ttttctgcct	1560
gggtttgata	tagtttaggg	ttggggtag	attaagatct	aaattacatc	aggacaaaga	1620
gacagactat	taactccaca	gttaattaag	gacgtatgtt	ccatgtttat	ttgttaaagc	1680
agtgtgaata	gccttcaagc	atgtgaataa	tcttccatct	tccccgccac	acatacacac	1740
acacactttt	tgtttctttc	aggtagacac	ctttttaa	gcagaactaa	ctgaggcatt	1800
tcagtaactt	tgctttcaaa	tcaataaagt	caaagtgtatg	gaaacatttt	gtgccttact	1860
ctccataccc	cgtgtactca	aattctctac	tgtatgaatt	atgctttaag	tagaattcag	1920
tgccaaggag	aacttgggtga	aataaattat	tttaattttt	tttttatcct	ttacaaagcc	1980
atggatttta	tttgggtgat	gtgtgctctg	tacacaagcc	atttcaatag	gatggagctg	2040
tttaattttt	tccaaagagt	aatagacatg	caaaagtttc	aataaaaaact	gggccattaa	2100
caagtaaat	aataaaactaa	taagcattcc	cttctagggt	tttgccaaac	tgccatcca	2160
ataacaaatt	tgagaatcgt	tgtaaaagct	agttatat	cagagaaatg	attttcatta	2220
ttgaaactgt	tctccctagc	aggccatttt	ccctttttcc	tgagggttta	gcaagtttag	2280
gagagaatag	tcatgaaaag	aaagggagaa	aaagggagaa	gggaagaggt	taaaaagtaa	2340
gtgctcagac	ctatgaacgt	aatccctttg	ctacaaatat	ttaagagcag	ctcagcttgg	2400
ttgaaactga	gttttgtcat	cttccatatt	tgcaggaagg	tattttctga	cttgcaatgc	2460
agctagatgt	aaaattttat	tttatcatcc	tagaaaagcct	tgactagaaa	aatgaataaa	2520
tattgagggt	ttctgtcca	tatctggctt	gcatgtgcca	gaaagcagag	aatagaaaat	2580
gtaatctcca	acatccaagc	atcgaaaccc	aaggggtagg	caattctatg	taggttttgg	2640
acatgaagtt	tggtgcatct	tggtttatgc	tggtcctaac	gctattaaac	ctctctggct	2700
tatagtctct	tcattctatt	agacaagcac	gtatcgaca	cttgcttcgc	acaaggtctt	2760
ttagtttaaca	atttagcagc	tactgtttgt	gttaaacaca	cttttcacca	aatagggtct	2820
gaggcaaacg	agagcaatga	ctattttaa	aaaggctttc	ccagcatcac	ttacacatcc	2880
caaaactaaa	aagatcaact	cttccaactg	agaaaagact	cctggctttg	aatggaaaact	2940
tacagcagag	agtcacaggc	cacggcaaca	acaacgacaa	caacaaacat	ttggaatatt	3000
attctcaact	cacgttttaa	taatacatct	tattattttt	ctagtagaga	aactacaaat	3060
cagcctcttc	aacattttata	tacagtttaa	taagcctctt	gcaagttact	tgttctctca	3120
cctgagggtat	ttttttcttc	cccaccttgc	ccctgttctt	cccttctctt	tctccctttg	3180
caagaggaaa	tatttaacat	atttgggtcc	aacttcaata	atgtaataat	taatacatta	3240
aaagcattta	acttcccttc	tagaaaaatg	cacaggctaa	ggcatagaca	aaacaaagag	3300
aaatgctgag	aaatttgcca	ctggagacaa	gcaatctgaa	taaataattg	ccaaaagttc	3360
tttttatgtc	atatagtgtc	aggatttgaa	ggagctat	ttttttaatg	ttgcaactag	3420
caactcatct	tcggaagaca	cagccaggag	aatgaagtag	aagtgaagg	tttataaatc	3480
catttgtaag	catttatccc	atatatttta	aattcaagaa	aaattgtgtt	tatctttaga	3540
attttgtatt	caatacttta	tgtactatgt	gactcatgct	tctggataaa	taaagcacca	3600
aatatgtatc	tgtaaccaca	atcacacata	ttatatttaa	tatatatcta	tataacaaaa	3660
aaaaaaaaaa	aaaa					3674

<210> 74

<211> 2797

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (853)

<223> n equals a,t,g, or c

<400> 74

ggcagcagag	agcagacaga	attatatgta	gaggacacag	gagatattta	cattgtggat	60
ggagatggag	gattgaataa	cagattgata	aaactgtccc	aagatttcat	gatcctttgg	120
ctgcatggag	aaaatgggac	agggcctgct	aagttcaaca	tacctcacag	tggtacactt	180
gattcagctg	gtcgggtgtg	gggtgctgac	cgaggaaata	aaagaatcca	agtatttgat	240
aaagacactg	gggagtgtt	aggagcatgg	aataattgtt	tcacagaaga	ggaccttctt	300
cagtcagttt	actcctgatg	ggaagtactt	gattgtggcc	cagctgaatc	ttagcaggct	360
ctcagtcgta	gcagcaccoc	cagtgggaag	cattggggag	tggtctgtga	tcagcacaa	420
ccaactagca	gatcaagttt	tgccacatct	cctagaagtc	gacagaaaag	actggagcag	480

tctatgtagc	agaaattgga	gcaaaacaag	tacaaaaata	tgtccctttg	aatagctatg	540
ttccttcatt	tggttcataa	tgtttctttc	ccgggaatat	ttcaagtggc	agttcagatt	600
ctcaattcac	taagtgttta	aaaatgatgt	tcaagcacaa	gaattttatt	ttctagtata	660
aaagatctag	tatcagaaaag	atttgttttt	gtatcattaa	gaatcttata	ttttgttgcc	720
ctcttgggac	ttagttttat	ttgtaagtgc	ataaggatat	tttaatgaaa	ggaaagtaac	780
taaaaaatgg	ggttgggaag	agggactaag	gtggtaacct	cattatttgc	cctggtagac	840
tgattctccc	tgngtaaaaa	aaatgggaat	aaaaatgagc	ttgcatgata	atttattaaa	900
tttcatgtga	agaactccag	acctccagat	tgtgcaacta	acataaaagt	agctgcttga	960
gagattgtaa	ataagatgaa	ctattgatta	atttgagtac	ccacagagtg	ctgtgtcttg	1020
acgacttaaa	aatgaaaaag	catgattgcc	ttttgagtag	cttgcaagtct	agtggggaga	1080
caagcaggca	aacagtcaca	acacagcaaa	agcgaccttg	gagcatagtg	ggacttttgg	1140
agtaggagtg	ctgcatttga	ctgaggggaat	catggatact	tcgcaggaga	agtgaatttg	1200
agctcagact	tgaaaactga	ggaggagcct	accaagggac	aaggaggaga	aaacaataat	1260
ttccaagtaa	agaaggtata	aaaagttaga	agtgtactgt	aaactttgat	aggcttttag	1320
gcctttttta	aagcccaact	tggtctctgt	ccattaccta	taagatatat	aatgtcagtc	1380
agccttttaa	tgtaggaata	aaatggcttg	catctaagca	ctttagtaaa	agagggtttt	1440
acaaataact	aaggatttga	gagcttcctt	ctcttttttt	ttctttttct	ttctttttgt	1500
ttacatgaac	tcaacttatt	cctaacattt	gtctacctca	aagaaatttc	aagattattt	1560
agataacatg	gatatgtgcc	aaatcctttg	agctgttaag	atgataattt	cctgctttcc	1620
tcctacatct	tctcctccca	ctcctcctt	tggtgtgaat	attggcttcc	caattaagac	1680
cttttttttt	tccagtttgt	tttagcttat	tataggtttt	ggaggaaact	tgccattttg	1740
taatctttca	aatcattctt	cacccttcct	cacatcagct	tcctgctttt	ccagtggttt	1800
tactgtaaat	tgtgtagcat	atgacaaatc	ttgagctgac	tttctcttct	acctgttatg	1860
gctggagtat	ttccagacc	tgaagggaact	cacacttggt	ttgatacttg	gatcacatct	1920
ccgtgaggtt	aggaaggtaa	atctaccaac	aggaagccct	gtactctgta	ttccaaggcc	1980
attggtaaat	gtgttggtgc	cactgatcgg	actgtatgac	cttaacaag	tcaccttagt	2040
tttcagtga	atgggaatat	cattgtctcc	tctttcatga	atgctgtgag	aatcagatgt	2100
gcaacaggta	catacttgcc	ctttggaaat	ctaatacctc	tgggatacca	ttaagaggca	2160
ttttaattaa	acaaaagggc	ccttctaaat	gtgctattta	tttgacaata	actatcagat	2220
ttgccttaat	tttgtgttta	tagcatttat	caaaacgtat	cctcatagac	tttatgcaga	2280
ttaatatggt	caattgattt	ggataaaaaga	aagtaatttc	agggtttggt	tttaagccag	2340
gacaagaagt	gcaaatgcct	ctttgaagca	atttaggcta	aactgatttt	gaaatttcaa	2400
aatgttttat	ttacttttgt	tttattaagc	caggacaaga	agtgc aaatg	cctctttgaa	2460
gcaattcagg	ctaggtaaac	cgtttttgcc	atttcaaaac	gttttatttt	actttgwttt	2520
rttrtcagagt	yttawaarvc	ctgctgcaaa	tatttctgaa	tgtctttgta	aaagtgtttg	2580
ttagtgtacc	tgtgattata	gtacttcact	tttttccttt	ggatttaattg	gttaaatgaa	2640
tgagaaatgt	gttatgtttt	ttactaaaaa	gtataaatta	aaatttttga	aagaaaaggc	2700
aatattatct	ggctcccaa	ttaaagtttg	attttattgt	cacaaaaaaa	aaaaaaaaaa	2760
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaa			2797

<210> 75
 <211> 2703
 <212> DNA
 <213> Homo sapiens

<400> 75						
ggcacgagat	ttcctacagg	tgaaacgcc	tcattaggat	tcactgtaac	gttagtgcta	60
ttaaactcac	tagcattttt	attaatggcc	gttatctaca	ctaagctata	ctgcaacttg	120
gaaaaagagg	acctctcaga	aaactcacaa	tctagcatga	ttaagcatgt	cgtttggcta	180
atcttcacca	attgcatctt	tttctgccct	gtggcgtttt	tttcatttgc	accattgatc	240
actgcaatct	ctatcagccc	cgaataaatg	aagctctgta	ctctgatatt	ttttcatttg	300
cctgcttgcc	tgaatccagt	cctgtatgtt	ttctcaacc	caaagtttaa	agaagactgg	360
aagttactga	agcgacgtgt	taccaagaaa	agtggtatcag	tttcagtttc	catcagtagc	420
caaggtgggt	gtctggaaca	ggattttctac	tacgactgtg	gcatgtactc	acatttgtag	480
ggcaacctga	ctgtttgcga	ctgctgcgaa	tcgtttcttt	taacaaagcc	agtatcatgc	540
aaacacttga	taaaatcaca	cagctgtcct	gcattggcag	tggcttcttg	ccaaagacct	600
gagggctact	ggtccgactg	tggcacacag	tcggccact	ctgattatgc	agatgaagaa	660

gattcctttg	tctcagacag	ttctgaccag	gtgcaggcct	gtggacgagc	ctgcttctac	720
cagagtagag	gattcccttt	gggtcgctat	gcttacaatc	taccaagagt	taaagactga	780
actactgtgt	gtgtaaccgt	ttccccgtc	aaccaaaatc	agtgtttata	gagtgaaccc	840
tattctcadc	tttcatctgg	gaagcacttc	tgtaatcact	gcctgggtgc	acttagaaga	900
aggagaggtg	gcagtttatt	tctcaaacca	gtcattttca	aagaacaggt	gcctaaaatta	960
taaattgggtg	aaaaatgcaa	tgtccaagca	atgtatgac	tggttgaaac	aaatatatga	1020
cttgaaaagg	atcttaggtg	tagtagagca	atataatggt	agttttttct	gatccataag	1080
aagcaaat	atacctat	gtgtattaag	cacaagataa	agaacagctg	ttaatat	1140
ttaaaaaat	attttaaaat	gtgattttct	ataactgaag	aaaatatctt	gctaatttta	1200
cctaattgtt	cactcttaat	ctcaggacaa	cttactgcag	ggccaaaaaa	gggactgtcc	1260
cagctagaac	tgtgagagta	tacataggca	ttactttatt	atgtttttac	ttgccatcct	1320
tgacataaga	gaactataaa	ttttgtttta	gcaattttata	aatctaaaac	ctgaagatgt	1380
tttttaaaaca	atattaacag	ctgttaggtt	aaaaaaatag	ctggacattt	gttttcagtc	1440
attatacatt	gctttgggtc	aatcagtaat	tttttcttaa	gtgttttgtg	attacactac	1500
tagaaaaaaa	gtaaaaggct	aattgctgtg	tgggtttagt	cgatttggct	aaactactaa	1560
ctaattgtgg	ggtttaatag	tatctgaggg	atltgggtgc	ttcatgtaat	gttctcatta	1620
atgaatactt	cctaatatcg	ttggctctac	taatat	caatttgcgt	ggatgtcacc	1680
tagcaatagc	ttggattata	tagaaagtaa	actgtggtca	atacttgcac	ttaattagac	1740
gaaacgggga	gtaattatga	cacgaagtac	ttatgtttat	ttcttagtga	gctggattat	1800
cttgaaccctg	tgctattaaa	tggaaatttc	catacatctt	ccccatacta	ttttttataa	1860
aagagcctat	tcaatagctc	agaggttgaa	ctctgggttaa	acaagataat	atgttatttaa	1920
taaaaataga	agaagaaaga	ataaagctta	gtcctgtgtc	tttaaaaatt	aaaaatttta	1980
cttgattccc	atctatgggc	tttagacctt	ttactgggtg	gagtcctaaa	gttataattg	2040
ttcaatatgt	tttttgaaca	gtgtgctaaa	tcaatagcaa	acccactgcc	atattagtta	2100
ttctgaaat	actaaaaaaa	tccagctaga	ttgcagttta	ataattaaac	tgtacatact	2160
gtgcatataa	tgaattttta	tcttatgtaa	attat	gaacacaagt	tgggaaatgt	2220
ggcttctgtt	catttcgttt	aattaaagct	acctcctaaa	ctatagtggc	tgccagtagc	2280
agactgttaa	attgtggttt	atatactttt	tgcattgtaa	atagtccttg	ttgtacattg	2340
tcagtgtaat	aaaaacagaa	tctttgtata	tcaaaatcat	gtagt	ttaaattgtg	2400
gaaggattta	tttacagtgt	gttgtaattt	tgtaaaggcca	actatttaca	agttttaaaa	2460
attgctatca	tgtatattta	cacatctgat	aaatattaaa	tcataacttg	gtaagaaact	2520
cctaattaaa	agggtttttc	caaaattcag	gttatgaaa	at	ttattcattt	2580
aaaaactaga	ataacagata	tataaaagt	ttaatctttg	tgctatatgg	tatgaaatac	2640
aatattgtac	tcagtgtttt	gaattattaa	agtttctaga	aagcaaaaaa	aaaaaaaaaa	2700
aaa						2703

<210> 76

<211> 742

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (707)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (724)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (726)

<223> n equals a,t,g, or c

<400> 76

47

gcgctcgaga	atagtgggtc	ccccggrctg	caggattcgg	cacgagctca	cttcaatyct	60
tctttgagaa	gtttttcctt	tctccgcaac	cagatgtaca	tatttgaact	ctctttgtac	120
ttggagggca	cttctttcgt	ggtagttcct	ttatttttat	taatctctgt	atccttagat	180
agtcctccaa	caaccaagg	ttgggactct	gtcttacata	tctgggtgcc	cctcatagtg	240
cagtaataag	taagttgatt	atatacgagc	tatgtaactt	atatttttta	atgggtggat	300
atcactgagt	tttttttttt	aagaattttt	ttattgaggt	aaacttcaca	taacataaaa	360
ttaactattt	taaagtgaga	agttcagtgc	cacttagtat	tgtaacaat	gttgcataac	420
caccaccttt	atttaaagtt	ccaaaaaaa	tgttctcctc	taaaaggaaa	ccccatccca	480
ttaagcagat	actctccatt	ccttccttcc	tccagccccc	agcaaccacc	aatctgcttt	540
ctgtctctat	ggatttatct	attccttgcta	ttttatataa	atcgaattgt	atgagacctt	600
ttgtgtctgg	cttctttcac	ttagtacaag	tttttgagat	ttattttacat	agtagcatgt	660
atcaacactt	catttttatg	gccaaataaa	attgtattat	gtgtttntag	cacaaaaaaa	720
aaananaaaa	atgacctctg	ag				742

<210> 77

<211> 1825

<212> DNA

<213> Homo sapiens

<400> 77

ggcagcagca	tgtcacatgt	atacctatgt	aacacacctg	cacattctgc	acatgtatcc	60
cagaacttaa	attataataa	taaaaaaaga	ataattgggt	gatggcacat	ccagggttgc	120
caaagacagt	cccagtttat	gctgttgtcc	tggcattatt	aataatgaca	ctgcctttaa	180
ctctcacaat	taatttggat	gataacttat	atggtaactc	tgctaaataa	aaaaaataaa	240
aattaccata	gtaacaggaa	cctacttgaa	atgatgcctc	tgtttctatt	ctggcttgaa	300
ttctgcattc	tttgaggatt	tgtagcctca	tgacagaatc	ctatctacag	gtgatgtatt	360
tcatatgatt	tttggtctatt	tttttaacaa	tctcaagccc	aataatagcc	agtgtataaa	420
ggaatgtagt	tactttctcc	ccactttctg	gcaagttaag	tttagccacc	tgattacaag	480
aagggacatt	cagaggtagg	atggcacaaa	gacacagggt	ccactggaga	tcactggaag	540
cagctgcagc	agggttaaga	gaaggagtc	ccagcgagtc	ttcagtcacc	acacactaac	600
atcatcagtg	aaaagttcct	gggcctgaag	atccagctat	gttgtttcta	gttgactatt	660
ttaagtgaca	gaacttgccc	caagcattga	ccattttggg	tcctcaataa	gcctgattca	720
accagggtca	cccttgaaatc	tgctctccac	ctttccaata	aacctatttt	atgcatcatt	780
cagtgaagta	tttatttatt	tacttttttg	ctgagaaaca	tgactagatt	taggaaaaat	840
gtagaatttt	actttttttt	caatattttc	tgggttttcc	agagttttca	cgtgtttcac	900
accttccctt	gcttcccacc	attccccttt	ctattttgaa	ctagagagac	atgagtttga	960
attctagctg	tgtaacctga	gtcagttatt	taacctcttt	ttgtttctgt	ttctttgtct	1020
gtaaatagca	aaaactacaa	ttaactttag	tcctgtctgt	acaccaaagt	ttatcttgaa	1080
atattataca	tatttatatg	aattactact	gaaatgctct	aagatgccta	tgtgtgaatg	1140
gcattgttgt	aaagattaaa	taatataagg	gaagtgtctg	cttcagtgtc	tggcatataa	1200
taaaagctat	tattttttacg	attatttttc	atcttataga	agaattatcg	ttcttccctt	1260
ccaaagctaa	taaatggaca	tgtgtttatc	agacagaacg	taagagctgc	caaataaata	1320
gggaataggt	gctttcggga	gtctagggaa	ataaaggcca	gggaattgtt	cataaaattt	1380
agtaccata	aatagcctat	aagtagattc	cctagtttat	tctatgcagg	aaaataaagt	1440
tctacggagc	acagattcca	aaactaattg	gtcataaata	tcacctgaaa	gtttagaaaa	1500
tgtagcatca	tggacctctt	ttcataggtt	ctaaatctta	atatctgtgg	gatgggtgcag	1560
gaatctagct	ttgctaagtg	ccctcagatg	actcttgctg	ttctaggcta	aaatacatgt	1620
ggtttggctt	caatggacat	gttctctgaag	aatgtttgga	tgtcacacat	tcataatttag	1680
tatgagagat	gaggtcctcc	tctcatcatt	ttcttaggtt	ctcttctctc	cactccttac	1740
cctcccatca	cttacaataa	atctttttaga	aaattagcta	tacattttgtt	tcattataaa	1800
aaagaaagaa	gataaaaaaa	aaaaa				1825

<210> 78

<211> 1674

<212> DNA

<213> Homo sapiens

<400> 78
ggccacgaga gtatctgcgg cagctgcagg tcctggattt atttctcgat tcgctgtcgg 60
aggagaatga gaccctgggtg gagtttgcta ttggaggcct gtgcaacctg tgcccagaca 120
gggccaacaa ggagcacatc ctgcacgcag gaggtgtccc actcatcatc aactgcctat 180
ccagcccca tgaggagacg gtgctgtctg ccaccaccac gctcatgcac ctgagcccg 240
cgggccgcag ctttctccca gagctgaccg ccacgcccgt ggtgcagtgc atgcttcgct 300
tctccctctc ggccagcgcc aggttccgga acctggcaca gatcttcctg gaggacttct 360
gctccccccg ccaagtggcc gagggccgca gccggcaggc gcaattttgc cctgggtatc 420
ccactgccga ggagcgtggc cccacggcag cgctgatcca tggagactgc gagaccgtgg 480
cacccttact gctggggacc acagtccctga tgtggacgca gggaaacgggg agcacatact 540
gccccattgg tgccttttca gccatctgaa aggcgggttc ttccagcagg acaggcattt 600
acactgatga aacgccactg ggagtgagga agccagactc cagagacacg gagaagatca 660
aactggagct cgtttcatag gctggcactc tcaatccctac atcagggtgcc accaccacca 720
gactcaggcc ctggtgtaag aagcggccaa gtgcctggac ccagaggctt tgcaggacag 780
tgttctcagg agctgggcct gaggtctagg agagctgcct tcgctgcagg aaatcaggga 840
ttatccctta acagaagtgt ctggagtagt ttccaggtat aggaatgaga tgcctcgtgg 900
tgaaaggatc tcaccctggg aagatgtggt gccccctcca gggctctgga ggatggatgc 960
ctccccagg ggctctccaa gctgggcatt tgggcctggt ggatgccaac ctggataacc 1020
tgtggcccag cattgactgt ccaccagcc ttgctgttag gcaccatgac tccaaagatg 1080
aagatgtggt ccctgccctt gagtgacagc cccagggaact taatgtggcc atcgggcatc 1140
aagcacaagg ccatgcaggt gatgatacgt cggaaatagag gcaccagccc tggtaactgc 1200
atcttctccc cttgccaccc catggccccg gctgaaagct tcggccctcc tctgtgtca 1260
ctcaatgatg gggagcccta cccagaaagt gtatcccacg agggcatcag ggacgcagt 1320
agtgttgctc aagggagtca ggaagagacg gcaacgtaaa ggatgtggct ccattgccat 1380
ggtgccccct ggtcaacata agggagcgtg gatccgatgg aaagggtggag ctgagggaaa 1440
atgggggtcc ttgcctctcg tgtaccacct caaggctgac cccttagatg gccagggaat 1500
ggcaggtgct acaaaaatgg taccacgtg ggcatggaaa tggggcagat taggggacca 1560
ctggactcag aggggagggg agggctcacc agcaccgct cagggagcct gtccctttat 1620
gttcccaaat aaagggtcct agaagactaa aaaaaaaaaa aaaaaaaaaa aaaa 1674

<210> 79
<211> 2191
<212> DNA
<213> Homo sapiens

<220>
<221> SITE
<222> (1327)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (1334)
<223> n equals a,t,g, or c

<400> 79
ccttctctaa aaaagcaaac aggcaaaact tcatgagaat cttgatcatg ttaaaatttt 60
atgtccttcg atttctccct acacacacac acacacacac acacacacac acacacacac 120
tcaacatttc ctccacccat atcatcactc cttagcatct ttattccatc aaaactttct 180
acccttgac attctctgtg cagttttgaa aattaccctc tcagcattct ctgttcaccc 240
ccacacactag accctgacct ctagtcaatt ctactacca ggggtgtcca cggttccagc 300
ctcctccatg aagcccagtt ctatgggctc actcctctgg gtaagtggga gccccagct 360
atcatcctca ttgtatagaa aaccaactct gtgatgttac ctgcccctct tccccttctc 420
tcttgaaaga gggctggggg agagggtggg ggactgttta tggccctggc cgggtctgta 480
ttcgtactgg gaggagtatt ggtactctgt ttagaaagaa atggggaggg ggaatgggg 540
tggcctcagc atctccctaa gtcccagcct ttaagtctc ctgttgcatg tcgtcgctgc 600

agcttcgaga	ggagttggat	cgatcttctt	gtggaaacgt	cctcttcaat	ggttacctgc	660
cgccaacagg	taggcactcc	caatggaatg	gaggggcgcg	gaggtgggccc	aaagactaca	720
tttcccataa	ggctgcagct	ctcgggtgcc	tgtgctgtgc	gtcctgagat	acagtgggaa	780
gtgtagttcc	ctatcagatg	cttgggctga	tgcttgaaaa	ggaagttgga	cacagcattt	840
cccatgaaac	aatgggccc	ctaactcttg	aagctcaaaa	agatgtccct	ggaaccccat	900
ggggaatttg	ttatcccggg	tttgggttcc	ttttgttagg	gggggctttg	ggaaaaactg	960
gggattcttc	cgtagggaag	gggaaaaaat	attaaatagg	aagttattga	cattaatgcc	1020
catgatagcc	acccactg	gccatggaag	gtatgcccc	gtgggtattg	gaactaggct	1080
tttctgattg	gtagaagtaa	cagagtaggg	aaatttcac	tacagcttta	tttccctaac	1140
tgcagtcagc	acctgtacct	tcatgaaagt	tgccagatat	aaagatctgt	agtagtactt	1200
ttccaactta	gttttatcct	gttttccga	aaaacaatca	tttatttatt	tatttattta	1260
tttaatttta	tgagacaggg	tctggctttg	tcacccaggg	tggagtgcag	tggtgcgac	1320
ttggctnca	tgcnacctct	gcctctcaga	ttcaagccat	ccttccacct	cagctctgcc	1380
actgagtagc	tgagactaca	agcactcgcc	accatgcccc	gctaattaaa	aaaataataa	1440
tcatttttaa	tgcaagcttt	atattataaa	tacaaagtaa	acatgaaaat	aaaacccaaa	1500
catagcagtg	ttattaaact	ctggcctgta	gcagtggttc	acacctgtaa	tcctagcagt	1560
ttggaggccg	agacaggtgg	attacttgag	acctggagtt	tgagaccagc	ccaggtgaca	1620
cagcaagacc	tcatctctac	taaaaataaa	aaaaaattag	ccaggtgtgg	tggtatgcac	1680
ctgtgggtccc	agctacttag	gatgctggag	tgcgaggatc	gcttgagccc	aggaggtcaa	1740
ggctgcagtg	aactatgac	actcattaca	ccccagcctg	ggtgacagag	cgagatgctg	1800
tctcaaaaaca	aaacaaaacg	aaaaacaact	ctggctagat	gctattgctt	gccaagggtg	1860
cagtcttcca	tttattaaaa	gtgaaaatta	gggccaggca	cattggctca	tgctgtgaat	1920
cccagcactt	tgaggagctg	agggtgggtg	atcacctgag	gtcaggagtt	cgagaccagc	1980
ctggccaaca	tggtgaaacc	ttatctctgc	caaaaatata	aaagattagc	catgtgtcgt	2040
ggtaggggtc	tgtaatctca	gctacttggt	aggctgaggc	aggagaatca	cttgaaccca	2100
ggaggcagag	gttgacagtga	gccaagattg	tgccattgca	ctccagcctg	tgcaacgagc	2160
gaaactccaa	ctcaaaaaaa	aaaaaaaaaa	a			2191

<210> 80

<211> 1335

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1287)

<223> n equals a,t,g, or c

<400> 80

ggatatatcc	agggtgcgg	attttcccc	cttcaggttt	aaatgttcct	gtttttctac	60
ctttccctcg	cagtatacgc	tcaacggcaa	gawagtggaa	gttgccgtca	aacagatcat	120
cgctggaaaa	gccgtggagc	aaggaggtgc	tttctcgaac	cccagagacc	tggtctgtga	180
ccgggacatc	cctgagctgc	agggttctct	agtcagactg	gctggcgtgt	cactcagccg	240
cacccgtgtg	cactgtaact	tttgtgtgct	caagaaatta	tacagaaacc	tacagctgtt	300
gtaaaaggat	gctcgaccca	agtgttctgt	aggcttgggg	agggatcggt	tctctgtttt	360
gttaaactcg	gtgggtacct	ggatcttcca	cacgagtggg	attctggcct	tcagagacca	420
ggagggagtg	tctggggccg	agtgtggcac	tgtggtgaga	gtgtgtgtct	ttgcacacac	480
agtgcagcgg	gaacgggtgg	gctggctggg	gctgaagaca	gacacactcc	tgagccaagg	540
tcttgtcttc	aacctccccg	tcccgttgtc	ccattttgct	ctgtgaagg	gcaaatccct	600
ttcttccctt	cccatctcag	gctctcctgt	tttccctcag	ggtccagtat	gcctttgagc	660
tttagctgtt	agaaaggaac	ccccgtgact	tgacacagct	ttcacagctg	gctgctagga	720
ccggcgggct	gggtgttcac	gtgtgtctgt	gtcatggatg	caatgcaggc	cctggaggac	780
tgtgcgtcac	ccgtcaacca	gagcgtgcct	ccgggccagc	ttccctccaa	ggaatgagtg	840
gattttcatc	aggatctctt	tattgcacag	actgaatggc	tttcatgttt	tctaatgtga	900
attaggcatg	tgaagcagtg	gggtgtccacc	cgtgtccctc	atgggtgagc	cctccagctg	960
tgagcccagg	cagtgtggtc	accgagtggg	gaccctctc	accaggaacc	gcattccctgt	1020
gctgcctcca	cctgagagtt	gctagggggt	tcttgtcgag	atcatgtcat	cagcaccctt	1080

50

aagtcaagtc	acgggtttcc	atagccaggc	agttggtatg	tacaattcag	ttcagcgtat	1140
gaacttgtat	ctctaattctg	atgtccattt	ttatatTTTT	tgaactgag	cacaatgaaa	1200
tcctttcttg	aatcattttc	cttttggatt	ataaaaaat	gggggaaagt	gctatgatga	1260
attttatgca	ataaatgtat	acatgtntgc	acatgcaccc	atgctgaaaa	aaaaaaaaaa	1320
aaaaaaaaaa	aaaaa					1335

<210> 81

<211> 1867

<212> DNA

<213> Homo sapiens

<400> 81

cccacgcgtc	cgggccacag	cagagacagt	ggagggcagt	ggagaggacc	gcgctgtcct	60
gctgtcacca	agagctggag	acaccatctc	ccaccgagag	tcattggcccc	attggccctg	120
cacctcctcg	tcctcgcccc	catctcctc	agcctggtgg	cctcccagga	ctggaaggct	180
gaacgcagcc	aagaccctt	cgagaaatgc	atgcaggatc	ctgactatga	gcagctgctc	240
aaggttggtga	cctgggggct	caatcggacc	ctgaagcccc	agagggtgat	tgtggttggc	300
gctggtgtgg	ccgggctggt	ggccgccaa	gtgctcagcg	atgctggaca	caaggtcacc	360
atcctggagg	cagataacag	gatcgggggc	cgcattctca	cctaccggga	ccagaacacg	420
ggctggattg	gggagctggg	agccatgcgc	atgccagct	ctcacaggat	cctccacaag	480
ctctgccagg	gcctggggct	caacctgacc	aagttcacc	agtacgacaa	gaacacgtgg	540
acggaggtgc	acgaagtga	gctgcgcaac	tatgtggtgg	agaaggtgcc	cgagaagctg	600
ggctacgcct	tgcgtcccca	ggaaaagggc	cactcgcccc	aagacatcta	ccagatggct	660
ctcaaccagg	ccctcaaaaga	cctcaaggca	ctgggctgca	gaaaggcgat	gaagaagttt	720
gaaaggcaca	cgctcttggg	atatcttctc	ggggaggggg	acctgagccg	gccggccgtg	780
cagcttcttg	gagacgtgat	gtccgaggat	ggcttcttct	atctcagctt	cgccgaggcc	840
ctccggggccc	acagctgcct	cagcgacaga	ctccagtaca	gccgcacgtg	gggtggctgg	900
gacctgtctg	cgcgcgcgct	gctgagctcg	ctgtccgggc	ttgtgctgtt	gaacgcgccc	960
gtggtggcga	tgaccagggg	accgcacgat	gtgcacgtgc	agatcgagac	ctctcccccg	1020
gcgcgggaatc	tgaagggtgct	gaaggccgac	gtggtgctgc	tgacggcgag	cggaccggcg	1080
gtgaagcgca	tcaccttctc	gccgcggctg	ccccgccaca	tgcaggaggc	gctgcggagg	1140
ctgcactacg	tgccggccac	caaggtgttc	ctaagcttcc	gcaggccctt	ctggcgcgag	1200
gagcacattg	aaggcggcca	ctcaaacacc	gatcgcccg	cgcgcatgat	tttctaccg	1260
ccgcgcgcgc	agggcgcgct	gctgtgggcc	tcgtacacgt	ggtcggacgc	ggcggcagcg	1320
ttcgccggct	tgagccggga	agaggcggtg	cgcttgccgc	tcgacgacgt	ggcggcattg	1380
cacgggctcg	tcgtgcgcca	gctctgggac	ggcaccggcg	tcgtcaagcg	ttggcgggag	1440
gaccagcaca	gccagggtgg	ctttgtggta	cagccgccc	cgctctggca	aaccgaaaag	1500
gatgactgga	cggtccctta	tgcccgcatc	tactttgccg	gcgagcacac	cgccctaccg	1560
cacggctggg	tgagacgggc	gggtcaagttg	ctgcgcgcgc	ccatcaagat	caacagccgg	1620
aaggggcctg	catcggacac	ggccagcccc	gaggggcacg	catctgacat	ggaggggcag	1680
gggcatgtgc	atggggtggc	cagcagcccc	tcgcatgacc	tggcaaaagg	agaaggcagc	1740
caccctccag	tccaaggcca	gttatctctc	caaaacacga	cccacacgag	gacctcgcat	1800
taaagtattt	tcggaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1860
aaaaaaa						1867

<210> 82

<211> 984

<212> DNA

<213> Homo sapiens

<400> 82

gaattcggca	cgagcccagc	ggaagccaag	ccaccaggcc	ccccagcgtc	cacgaggagc	60
atgaacattg	aggatggcgc	gtgcccgcgg	ctccccgtgc	cccccgctgc	cgcccggtag	120
gatgtccttg	ccccacgggg	cattgtctct	cctctggctc	ttctccccac	ccctgggggc	180
cgggtggagg	ggagtggccg	tgacgtctgc	cgccggaggg	ggctccccgc	cggccacctc	240
ctgccccgtg	gctgtctcct	gcagcaacca	ggccagccgg	gtgatctgca	cacggagaga	300

51

mctggccgag	gtcccagcca	gcatcccggg	caacacgcgg	tacctgaacc	tgcaagagaa	360
cggcatccag	gtgatccgga	cggacacgtt	caagcacctg	cggcacctgg	agattctgca	420
gctgagcaag	aacctgggtg	gcaagatcga	ggtagggcgc	ttcaacgggc	tgcccagcct	480
caacacgctg	gagctttttg	acaacccggc	gaccacgggt	cccacgcagg	ccttcgagta	540
cctgtccaag	ctgcgggagc	tctggctgcg	gaacaacccc	atcgagagca	tcccctecta	600
cgccttcaac	cgcgtgccct	cgctgcggcg	cctggacctg	ggcgagctca	agcggctgga	660
atacatctcg	gaggcggcct	tcgargggct	ggtcaacctg	cgctacctca	acctgggcat	720
gtgcaacctc	aaggacatcc	ccaactgacg	gccctgggtg	gcctggagga	gctggagctg	780
tcgggcaacc	ggctggacct	gatccgcccc	ggctccttcc	agggctctac	cagcctgcgc	840
aagctgtggc	tcatgcacgc	ccaggtagcc	accatcgagc	gcaacgcctt	cgacgacctc	900
aagtcgctgg	aggagctcaa	cctgtcccac	aacaacctga	tgctcgctgc	ccacgacctc	960
ttcacgcccc	tgcaaccgcct	cgta				984

<210> 83
 <211> 2664
 <212> DNA
 <213> Homo sapiens

<400> 83

ggttgctggc	ccaggtgagc	ggcgcgctg	gtccaggtga	gcgggcgcgt	ccccgcgacg	60
gcgctgcctg	cccgaggcgg	ttcacgtaaa	gacagcgaga	tcctgagggc	cagccgggaa	120
ggaggcgctg	atatggagct	ggctgctgcc	aagtcggggg	cccgcgccgc	tgccctagcgc	180
gtccctgggga	ctctgtgggg	acgcgcgccg	cgcgcggct	cggggacccg	tagagcccg	240
cgtgcgcgc	atggccctgc	tctcgcgcc	cgcgctcacc	ctcctgctcc	tcctcatggc	300
cgctgttgtc	aggtgccagg	agcaggccca	gaccacgcag	tgagagagcca	ccctgaagac	360
catccggaac	ggcgttcata	agatagacac	gtacctgaac	gccgccttgg	acctcctggg	420
aggcgaggac	ggtctctgcc	agtataaatg	catgacggat	ctaagccttt	cccacgttat	480
ggttataaac	cctcccacc	gaatggatgt	ggctctccac	tgtttggtgt	tcactctaac	540
attggtatcc	cttccctgac	aaagtgttgc	aaccaacacg	acaggtgcta	tgaracctgt	600
ggcaaaagca	agaatgactg	tgatgaagaa	ttccagtatt	gcctctccaa	gatctgccga	660
gatgtacaga	aaacactagg	actaactcag	catgttcagg	catgtgaaac	aacagtggag	720
ctcttgtttg	acagtgttat	acatttaggt	tgtaaacccat	atctggacag	ccaacgagcc	780
gcattgcagg	gtcattatga	agaaaaaact	gatctttaa	ggagatgccg	acagctagtg	840
acagatgaag	atggaagaac	ataacctttg	acaaataact	aatgttttta	caacataaaa	900
ctgtcttatt	tttgtgaaag	gattattttg	agaccttaaa	ataatttata	tccttgatgtt	960
aaaacctcaa	agcaaaaaaa	gtgagggaga	tagtgagggg	agggcacgct	tgtcttctca	1020
ggtatcttcc	ccagcatttc	tcccttactt	agtatgccaa	atgtcttgac	caatatcaaa	1080
aacaagtgtc	tgtttagcgg	agaattttga	aaagaggaat	atataactca	attttcacaa	1140
ccacatttac	caaaaaaaga	gatcaaatat	aaaattcatc	ataatgtctg	ttcaacatta	1200
tcttattttg	aaaatgggga	aattatcact	tacaagtatt	tgtttactat	gaaattttaa	1260
atacacattt	atgcctagaa	ggaacggact	tttttttctt	attttaatta	cacataatat	1320
gtaattaaag	tacaacataa	tatgttggtt	ctctgtagcc	cgttgagcat	atgagtaagt	1380
cacatttcta	ttaggactac	ttmcaaggac	aagggtttcca	tttttccagt	tgtaaaattg	1440
gaaccatcag	ctgataacct	cgtagggagc	aacccagga	tagctaagtg	ttatgtaata	1500
tgccatagaag	gtgatgtgaa	tgcgattcag	aagcatagcc	actcccattt	tatgagctac	1560
tcacatgaca	aatgtcatct	tttgctataa	cctttgcca	gttagagaaa	agatggattt	1620
aatgagataa	atgaaaagat	atttamccca	atatatcaag	gcactatttg	ctgttatgct	1680
ttgttattta	tttcccagca	cttggtcctt	attgtagatt	ttttaagac	tgtaaccttt	1740
tactaactgt	ggtcttacta	aaatttggtg	ttgatactgc	ttttcaaaaa	gcctttaatt	1800
agagccaaaa	ggatggaaaa	ggcaagatat	aaatgccttt	tatagatctc	ttattttacat	1860
tgaaaatttat	taccatatgt	ttagagcaaa	tccaagaaaa	cttcaacagc	ttctgaagat	1920
gtctatgaat	gttgaaaact	tttcaatctc	ttggaatgct	cagttatggt	cctagaccgg	1980
tctttgctga	ctactggttg	ttaacctttc	cctagcctgg	gacctcaagc	catatatatc	2040
ctttgggtga	cccatggcca	aagtatttaa	gatgaactga	ctttcaaaagt	cagagaagga	2100
cagcataggg	agagggcggt	atttgtaagt	cattacaggt	agaacagggc	agaaggaaaa	2160
gtatgttctg	gagaaaaggc	catgttccca	actttggaga	tatgtcattg	ccgggaacct	2220
agtatcttcc	aacttgaatt	ggtggcagct	gttccagtga	gacaaggcac	atgtatgcct	2280

52

tgtggctaag	tgagcaaaact	gggtttccac	ttaaatgttt	gggaccctca	attgattctt	2340
tatttcaaac	ctttataaaa	ggtacagttt	tgtaagccat	tattaataat	taatgcttat	2400
cggctgggca	cagtggctca	cacctataat	cccagcactt	gggaggctga	ggcgggtgga	2460
tcacttgagg	tcaggagttt	gagaccagct	ggccaacatg	gtgaaacagc	gtctctacta	2520
aaaatacaaa	aatttgccgg	gcgtgggtggc	gcatgcttat	agtctcagct	actcaggaag	2580
ctgagggtacg	agaatcactt	gaaccagga	ggtggagggt	gcagtgagct	gagattgtgc	2640
cactgcactg	cagcctggct	cgag				2664

<210> 84

<211> 1328

<212> DNA

<213> Homo sapiens

<400> 84

cccacgcgtc	cgggccagtg	gaggtccgca	gagtttgggc	gccaggcgag	acggcagggc	60
ttaaagtcc	gggaatcaaa	gatcaactcc	cactgaggac	aaatggacct	gtaattccgg	120
gtgtgacgag	agaacgagat	ttaccttcct	gaattaaaaa	wcwgactccc	tgcgacaagg	180
actgtgtact	gcatgaatga	ggctgagata	gttgatgttg	ctctgggaat	cctgattgag	240
agccgcaaac	aggaaaaggc	ctgcgagcag	cgggccctgg	cgggggctga	taaccagar	300
cactccccctc	cctgctccgt	gtcgctcac	acaagttctg	ggagcagcag	tgaggaagag	360
gacagtggga	aacaggcact	grctccaggc	ctcagccctt	cccagaggcc	ggggggttcc	420
agctctgcct	gtagcaggag	ccctgaggag	gaggagggaag	aggatgtgct	gaaatacgtc	480
cgggagatct	ttttcagcta	gggcataaac	tgtgactga	actgtctgcc	gagagcagct	540
ggaggacagc	tgagcttcca	ctggtgctgc	tgggccgccc	gcctgtggga	atggggctct	600
ctgtgctcct	acctttgtgc	cttcttgggc	ctggcagatt	cacctcaggc	cagaagcccc	660
tggacactcc	gggccttggg	gctgccgttc	tgagtgtgcg	gaaggcagga	ctcaaaatga	720
gatcccatct	gactccctct	gtatgtactg	tgccctctcc	tggtctctga	ggctctggag	780
tcccaattgt	ctgtgttagt	cagtgaccag	gttccaggga	aaatgatgtc	atgtggtggt	840
ccaacttact	ggaaccaaag	agacagtact	ttgcaaagaa	aaggatcact	gccagggtca	900
ctggaattgc	tacagtttag	tcogcatgat	ctctcctgaa	ggaggaagcc	tgtttcaaaa	960
atagtttcca	tcatgagtct	atcaatgagc	tcccacctct	ccagccagcc	tagaaagcaa	1020
acgagctgcc	cacagttctc	tgccctgtct	gggaggttga	ggccacagtg	tatagactgg	1080
taagccagac	aggcctcctc	cgcgaagctg	ctaccttgct	ttcacctgta	ccttggtccc	1140
cgggcagcta	gctataaagc	aagagggaca	ggagcccaga	agagacactg	aggacaagag	1200
atcacaccag	agtacatgtc	tctgcctctg	ttttcagtgt	ggctttggac	aggaatatat	1260
gaataaatca	ctgccataca	ggttttccaa	tacacaagtg	ctagaaaata	cacacaattc	1320
cccaatga						1328

<210> 85

<211> 1342

<212> DNA

<213> Homo sapiens

<400> 85

ggcccgcoca	ggaggtatcc	tgcccttgac	tgcaactctt	gtcgtcttat	gtgggtgttg	60
aattgatctg	tctctgcagc	cagatccagg	ctcctggaag	aacctgtgcc	ggcagctact	120
ggtcatgccca	ggcacacact	gctgcccagg	aggagctgct	gtttgaatta	tctgtgaatg	180
ttgggaagag	gaatgccaga	gctgccggct	gaaaattacc	caaccaagag	aaatctgcag	240
gatggacttt	ctggctcctc	tcttgttcta	cctggcttcg	gtgctgatgg	gtcctgttct	300
tatctgcgtc	tgctcgaaaa	cccatagctt	gaaaggcctg	gccaggggag	gagcacagat	360
atcttctctg	ataattccag	aatgtcttca	gagagccrtg	catggattgc	ttcattacct	420
tttccatacg	agaaaccaca	ccttcattgt	cctgcacctg	gtcttgcaag	ggatggttta	480
tactgagtac	acctggggaa	gtatttggct	actgtcagga	gctggagttg	tccttgcat	540
accttcttct	gccctatctg	ctgctaggtg	taaacctgtt	ttttttcacc	ctgacttggt	600
gaaccaatcc	tggcattata	acaaaagcaa	atgaattatt	atttcttcat	gtttatgaat	660
ttgatgaagt	gatgtttcca	aagaacgtga	ggtgctctac	ttgtgattta	aggaaaaccag	720

53

ctcgatccaa	gcactgcagt	gtgtgtaact	ggtgtgtgca	ccgtttcgac	catcactgtg	780
tttgggtgaa	caactgcac	ggggcctgga	acatcaggta	cttcctcacc	tacgtcttga	840
ccttgacggc	ctcggctgcc	accgtcgcca	ttgtgagcac	cacttttctg	gtccacttgg	900
tggtgatgtc	agatttatac	cagggagactt	acatcgatga	ccttggacac	ctccatgtta	960
tggaacagg	ctttcttatt	cagtacctgt	tcctgacttt	tccacggatt	gtcttcatgc	1020
tggtgtttgt	cgtgggtctg	agcttcctcc	tggtgggcta	cctgttggtt	gtcctgtatc	1080
tggtggccac	caaccagact	actaacgagt	ggtacagagg	tgactgggcc	tggtgccagc	1140
gttgtccct	tgtggcctgg	cctccgtcag	cagagcccca	agtcacccgg	aacattcact	1200
ccatgggct	tgggagcaac	cttcaagaga	tctttctacc	tgcccttcca	tgatcatgaga	1260
ggaagaaaca	agaatgacaa	gtgtatgact	gcctttgagc	tgtagttccc	gtttatttac	1320
acatgtggat	cctcgttttc	ca				1342

<210> 86
<211> 1154
<212> DNA
<213> Homo sapiens

aagacaggaa	aagctccagg	ccgtgggtct	caaagtgtgg	tccttggaca	gcagcaacat	60
cacctaggag	cctgttaggg	aaggcacagc	ctcaggccct	gccccagacc	tgacagaatca	120
gaaactcttg	ggtgaggcct	ggttatctgc	tgtaacagac	cttcacagtgg	gttctgatgc	180
cctctagagc	aggagaacca	ctagcttaga	ggttgacagta	tggttggcat	cttgccattt	240
gtgttagttc	agagggaatgg	ctgaccccca	tgctctcattt	ctaagcttca	ggcagctttt	300
ctcctgggca	gctgtcattc	tggtgagggg	aatcctgggg	actgtggctc	ctcctccctg	360
tcctgtgtgc	cttgatctgg	cagtctaccc	ccttcacttc	cccgtggagg	ctccatgcct	420
agaggtggtc	ttcaaacaga	agaatggcaa	agataattgt	ctcgtgtttt	accctgaccc	480
cattccttta	agagggtcac	ttcttggccc	attcatttaa	aaaccaatgt	catagtctctg	540
tgattccacc	tatcagacag	tgccacgtcc	aaaggcgggg	ctctyacctc	cctgggaaga	600
gagactgttg	ctgtctgtgc	ttcctgtgtt	ctccagtccc	acgctccac	ggacccacgc	660
ccttgagagc	tcctcrgtg	tcccagggt	tctggtgtgt	tcagagacct	ccacactcaa	720
cgaccactgg	tgctgcagaa	gggcccgtgc	ttacattcca	attaacagac	gcttttccca	780
tctaattgct	cttgccctct	cctaacacca	cctcgggagt	gtttatgtct	attctaagtg	840
aatttcactg	tgtagaaaaa	ttcacacctg	ttgtcccagc	gatttgggag	gccggggcgg	900
gtgtatcatt	tgagcccagg	agtttgaggc	tagcctgggc	aggatggtga	aaccccgctc	960
ctataaagaa	attttaaaaa	ttagctgggc	atagtgccac	gtgcctgtag	ttccatctac	1020
tggtggaggc	gggtgtggag	gatcgcatga	gcccgggagt	ttgaggctgc	agtgaagctgt	1080
gatcgacaga	ctgcactcca	gtctgggcaa	cagagcaaga	ccctgtctct	taaaaaaaaa	1140
aaaaaaaaact	cgag					1154

<210> 87
<211> 1197
<212> DNA
<213> Homo sapiens

<220>
<221> SITE
<222> (573)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (1177)
<223> n equals a,t,g, or c

<220>
<221> SITE

<222> (1185)

<223> n equals a,t,g, or c

<400> 87

aagacaggaa	aagctccagg	ccgtggttct	caaagtgtgg	tccctggaca	gcagcaacat	60
cacctaggag	cctgttaggg	aaggcacagc	ctcaggccct	gccccagacc	tgagaaatca	120
gaaactctgg	ggtgaggcct	ggttatctgc	tgtaacagac	cttccagtgg	gttctgatgc	180
cctctagagc	aggagaacca	ctagcttaga	ggttgacagta	tgtttgccat	cttgccattt	240
gtgttagttc	agaggaatgg	ctgaccccca	tgtctcattt	ctaagcttca	ggcagctttt	300
ctcctgggca	gctgtcattc	tggtgagggg	aatcctgggg	actgtgggctc	ctcctccctg	360
tccgtgtgtc	cttgatctgg	cagtctaccc	ccttcatctc	cccgtggagg	ctccatgccw	420
agaggtggtc	ttcaaacaga	agaatggcaa	arataattgt	ctcgtgtttt	accctgaccc	480
cattccctta	agagggtcac	ttcttggccc	attcatthaa	aaaccaatgt	catagttctg	540
tgattccacc	tatcagacag	tgccacgtcc	aangcggggc	tctcacctcc	ctgggaagag	600
agactgttgc	tgtctgtgct	tcctgtgttc	tccagtccca	cgctcccacg	gacccacgcc	660
cttgaggact	ccctcagtgt	cccagggtct	ctggtgtgtt	cagagacctc	cacactcaac	720
gaccactggt	gctgcagaag	ggccggtgct	tacattccaa	ttaacagacg	cttttcccat	780
ctaatagcctc	ttgccttctc	ctaaccaccac	ctcgggagtg	tttatgtcta	ttctaagtga	840
atttcactgt	gtgaaaaaat	tcacacctgt	tatccagca	atttgggagg	ccgagggcgg	900
tgtatcattt	gggcccagga	gtttgagact	agcctgggca	agatggtgaa	accccgcttc	960
tataaagaaa	ttttaaaaaat	tggctgggca	tagtggcgcg	tgctgtagt	tccatctgct	1020
ggggaggctg	gggtgggagg	atcgcatgag	cccgggagtt	tgaggctgca	gtgagctgtg	1080
atcgggcac	tgactccag	tctgggcaac	agagcaaaac	cctgtctctt	aaaaaaaaaa	1140
aaaactcgag	ggggggcccg	gtaccaaat	cgccctnats	agtgnagtgc	tattaca	1197

<210> 88

<211> 910

<212> DNA

<213> Homo sapiens

<400> 88

ggcagagctg	gccttcgact	cgctatgtcc	actaacaata	tgteggaccc	acggaggccc	60
aacaaagtgc	tgaggtgagg	acccagcgt	cgtgggcacg	ggttcggggt	gtgggtgtgg	120
atcggggccc	tggaagcgc	ctgtctatcc	cgggggcagg	acctgagcgc	ccctgaccct	180
cgagcctgtc	gcaggtacaa	gccccgcgcg	agcgaatgta	accggcctt	ggacgacccg	240
acgcccggact	acatgaacct	gctgggcatg	atcttcagca	tggtcggcct	catgcttaag	300
ctgaagtggg	gtgcttgggt	cgctgtctac	tgctccttca	tcagctttgc	caactctcgg	360
agctcggagg	acacgaagca	aatgatgagt	agcttcatgt	gagacttgcc	ctacagaaca	420
agtgactctt	gagtaagggg	tggggggacc	ccagcctggc	catcctagac	tgacacctct	480
ctcctgtctt	catgctgtcc	atctctgccg	tggtgatgtc	ctatctgcag	aatcctcagc	540
ccatgacgcc	cccatggtga	taccagccta	gaagggtcac	atthttggac	ctgtctatcc	600
actaggcctg	ggctttggct	gctaaacctg	ctgccttcag	ctgccatcct	ggacttccct	660
gaatgaggcc	gtctcgggtc	ccccagctgg	atagagggaa	cctggccctt	tcctagggaa	720
caccctaggc	ttacccctcc	tgctccctt	ccctgcctg	ctgctggggg	agatgctgtc	780
catgtttcta	ggggatttca	tttgctttct	cgttgaaacc	tggtgttaat	aaagtthttc	840
actctgaaaa	aaaaaaaaaa	aaaaaaaaac	tygrgggggg	gcccggaaac	caattcscgg	900
gatagtgagt						910

<210> 89

<211> 1076

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1029)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1037)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1040)

<223> n equals a,t,g, or c

<400> 89

ggcagcagggg	gaaagccatg	ctcccaggac	tccttccttg	cagccttaaa	tcggtctgta	60
cggaaaattc	cgcgcccttag	aaacccacgc	ttgggtgtaa	cttattattg	ttcttcctga	120
cctacttcct	gtttatcact	tccgggttca	tcattttggc	atttcggtga	tcgggttgga	180
actattgaag	cccgttttca	ggttcttttc	cccattttcc	ctttgaaagg	aagacttctg	240
gcttctccta	aatctccgtt	ctctgggtta	ggggagtcca	agcctctgtc	atgaggaacg	300
gaaatgcgag	ggcctcgggt	gttactctaa	aatccgcctt	cagcttgac	gccggaagct	360
gcgattcctg	cagcggaaga	ggcgtgatct	ggccttcgac	tcgctatgtc	cactaacaat	420
atgtcggacc	cacggaggcc	gaacaaagt	ctgaggtaca	agcccccgcc	gagcgaatgt	480
aacccggcct	tggacgaccc	gacgccggac	tacatgaacc	tgctgggcat	gatcttcagc	540
atgtgcggcc	tcatgcttaa	gctgaagtgg	tgtgcttggg	tcgctgtcta	ctgctccttc	600
atcagctttg	ccaactctcg	gagctcggag	gacacgaagc	aaatgatgag	tagcttcatg	660
ctgtccatct	ctgccgtggt	gatgtcctat	ctgcagaatc	ctcagcccat	gacgccccca	720
tggtgatacc	agcctagaag	ggtcacattt	tggaccctgt	ctatccacta	ggcctgggct	780
ttggctgcta	aacctgctgc	cttcagctgc	catcctggac	ttccctgaat	gaggccgtct	840
cgggtgcccc	agctggatag	agggaaacctg	gccctttcct	agggaaacacc	ctaggcttac	900
ccctcctgcc	tcctttcccc	tgctgctgc	tgggggagat	gctgtccatg	tttctagggg	960
tattcatttg	ctttctcggt	gaaacctgtt	gttaataaag	tttttctactc	tgaaaaaaaa	1020
aaaaaaaaana	raaaacnecn	ggggggggccc	ggaacccaat	tcscgggata	gtgagt	1076

<210> 90

<211> 1842

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (67)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (98)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (212)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1838)

<223> n equals a,t,g, or c

<400> 90

```

gcgaccgcgc ccttcagcta gctcgctcgc tgcgtctgct tccctgctgc cggctgcgca      60
tggtcttnggc gttggcggcg ctggcggcgg ctcgagcngc ctgcgsagcc ggtaccagca      120
gttgcagaat gaagaagagt ctggagaacc tgaacaggct gcagggtgatg ctctccacc      180
ttacagcagc atttctgcag agagcgca tnatcttgac tacaaggatg agtctgggtt      240
tccaaagccc ccattcttaca atgtagctac aacactgccc agttatgatg aagcggagag      300
gaccaaggct gaagctacta tccctttggt tcctgggaga gatgaggatt ttgtgggtcg      360
ggatgatttt gatgatgctg accagctgag gataggaaat gatgggattt tcatgttaac      420
ttttttcatg gcattcctct ttaactggat tgggtttttc ctgtcttttt gcctgaccac      480
ttcagctgca ggaaggatg gggccatttc aggatttggg ctctctctaa ttaaatggat      540
cctgattgtc aggtttttcca cctattttccc tggatatttt gatggtcagt actggctctg      600
gtgggtgttc cttgttttag gctttctcct gtttctcaga ggatttatca attatgcaaa      660
agttcgggaag atgccagaaa ctttctcaaa tctcccagg accagagttc tctttattta      720
ttaaagatgt tttctggcaa aggccttcct gcatttatga attctctctc aagaagcaag      780
agaacacctg caggaagtga atcaagatgc agaacacaga ggaataatca cctgctttaa      840
aaaaataaag tactgttgaa aagatcattt ctctctattt gttcctaggt gtaaaatttt      900
aatagttaat gcagaattct gtaatcattg aatcattagt ggttaatgtt tgaaaagct      960
cttgcaatca agtctgtgat gtattaataa tgccttatat attgtttgta gtcattttaa      1020
gtagcatgag ccattgtccct gtagtcggta gggggcagtc ttgctttatt catcctccat      1080
ctcaaaatga acttggaatt aaatattgta agatatgtat aatgctggcc attttaaagg      1140
ggttttctca aaagttaaac ttttgttatg actgtgtttt tgcacataat ccattttgc      1200
tgttcaagtt aatctagaaa tttattcaat tctgtatgaa cacctggaag caaaatcata      1260
gtgcaaaaat acatttaagg tgtggtcaaa aataagtctt taatttgtaa ataataagca      1320
ttaatttttt atagcctgta ttcacaattc tgcggtacct tattgtacct aagggtattc      1380
aaaggtgttg tcaactgtata aaacagaaag cactaggata caaatgaagc ttaattacta      1440
aaatgtaatt cttgacactc tttctataat tagcgttctt cacccccacc cccaccccca      1500
cccccttat tttccttttg tctcctgggtg attaggccaa agtctgggag taaggagagg      1560
attaggtact taggagcaaa gaaagaagta gcttggaaact tttgagatga tccctaactat      1620
actgtactac ttgcttttac aatgtgttag cagaaaccag tgggttataa tgtagaatga      1680
tgtgttttct gcccagtggt taattcatct tggtttgcta tgttaaaact gtaaatataa      1740
cagaacatta ataaatatct cttgtgtagc accttttaaa aaaaaaaaaa aaaaaaaaaa      1800
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa                                1842

```

<210> 91

<211> 1963

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (335)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1959)

<223> n equals a,t,g, or c

<400> 91

```

ggatcctcgc ggcggcggcg gtgcttacag cctgagaaga gcgtctcgcc cgggagcggc      60
ggcggccatc gagaccacc caaggcgcgt cccctcggc ctccagcgc tcccaagccg      120
cagcggccgc gcccttcag ctagctcgct cgctcgctct gcttccctgc tgcggctgc      180
gcatggcktt ggcgttggcg gcgctggcgg cggctgagcc gcctgcgcag ccggtaccag      240
cagttgcaga atgaagaaga gtctggagaa cctgaacagg ctgcaggtga tgcctccca      300
ccttacagca gcatttctgc agagagcgca gcatnathtt gactacaagg atgagtctgg      360
gtttccaaag ccccatctt acaatgtagc tacaacactg ccagttatg atgaagcgga      420
gaggaccaag gctgaagcta ctatcccttt gggttcctggg agagatgagg attttgggg      480

```

57

tcgggatgat	tttgatgatg	ctgaccagct	gaggatagga	aatgatggga	ttttcatggt	540
aacttttttc	atggcattcc	tctttaactg	gattgggttt	ttcctgtctt	tttgcccgac	600
cacttcagct	gcaggaaggt	atggggccat	ttcaggattt	ggtctctctc	taattaaatg	660
gacccgtatt	gtcagggttt	ccacctattt	ccctggatat	tttgatgggc	agtactggct	720
ctgggtgggtg	ttccttgttt	taggctttct	cctgtttctc	agaggattta	tcaattatgc	780
aaaagttcgg	aagatgccag	aaactttctc	aaatctcccc	aggaccagag	ttctctttat	840
ttattaaaga	tgttttctgg	caaaggcctt	cctgcattta	tgaattctct	ctcaagaagc	900
aagagaacac	ctgcaggaag	tgaatcaaga	tgcagaacac	agaggaataa	tcacctgctt	960
taaaaaata	aagtactgtt	gaaaagatca	tttctctcta	tttgttccta	ggtgtaaaat	1020
tttaatagtt	aatgcagaat	tctgtaatca	ttgaatcatt	agtggttaat	gtttgaaaaa	1080
gctcttgcaa	tcaagtctgt	gatgtattaa	taatgcctta	tatatgtgtt	gtagtcatct	1140
taagtagcat	gagccatgtc	cctgtagtcg	gtagggggca	gtcttgcttt	attcatcctc	1200
catctcaaaa	tgaacttgga	attaaatatt	gtaagatatg	tataatgctg	gccattttta	1260
aggggttttc	tcaaaaagtt	aacttttgtt	atgactgtgt	ttttgcacat	aatccatatt	1320
tgctgttcaa	gttaatctag	aaattttatt	aattctgtat	gaacacctgg	aagcaaaatc	1380
atagtgcata	aatacattta	aggtgtgggc	aaaaataagt	ctttaattgg	tataataata	1440
gcattaattt	tttatagcct	gtattcacia	ttctgcggta	ccttattgta	cctaagggat	1500
tctaaaggtg	ttgtcactgt	ataaaacaga	aagcactagg	atacaaatga	agcttaatta	1560
ctaaaatgta	attcttgaca	ctctttctat	aattagcggt	cttcaccccc	accccccccc	1620
ccacccccct	tattttcctt	ttgtctcctg	gtgattaggc	caaagtctgg	gagtaaggag	1680
aggattaggt	acttaggagc	aaagaaagaa	gtagcttgga	acttttgaga	tgatccctaa	1740
catactgtac	tacttgcttt	tacaatgtgt	tagcagaaac	cagtgggtta	taatgtagaa	1800
tgatgtgctt	tctgcccacg	tggttaattca	tcttggtttg	ctatgttaaa	actgtaaata	1860
caacagaaca	ttataaata	tctcttgtgt	agcaccttta	aaaaaaaaaa	aaaaaaaaaa	1920
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaa		1963

<210> 92
 <211> 1487
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1470)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1487)
 <223> n equals a,t,g, or c

<400> 92						
gcgaccgcgc	ccctttcagc	tagctcgtct	gctcgtctct	cttccctgct	gccggctgcg	60
catggckwtg	gcgttggcgg	cgctggcggc	ggtcgagccg	gcctgcgcag	ccggtaccag	120
cagttgcaga	atgaagaaga	gtctggagaa	cctgaacagg	ctgcagggtg	tgctcctcca	180
ccttacagca	gcatttctgc	agagagcgca	gttttccacc	tatttccctg	gatattttga	240
tggtcagtag	tggtctcgtg	gggtgttctt	tggttttagc	tttctcctgt	ttctcagagg	300
atztatcaat	tatgcaaaag	ttcggaagat	gccagaaact	ttctcaaatc	tccccaggac	360
cagagttctc	tttattttat	aaagatgttt	tctggcaaa	gccttcctgc	atztatgaat	420
tctctctcaa	gaagcaagag	aacacctgca	ggaagtgaat	caagatgcag	aacacagagg	480
aataatcacc	tgcttttaaa	aaataaagta	ctgttgaaaa	gatcatttct	ctctatttgt	540
tcctaggtgt	aaaattttta	tagttaatgc	agaattctgt	aatcattgaa	tcattagtgg	600
ttaatgtttg	aaaaagctct	tgcaatcaag	tctgtgatgt	attaataatg	ccttatatat	660
tggttttagt	catttttaag	agcatgagcc	atgtccctgt	agtcggtagg	gggcagtcct	720
gctttattca	tcctccatct	caaaatgaac	ttggaattaa	atattgttaag	atatgtataa	780
tgctggccat	tttaaaaggg	ttttctcaaa	agttaaactt	ttgttatgac	tggtgttttg	840
cacataatcc	atatttgcgt	ttcaagttaa	tctagaaatt	tattcaattc	tgatatgaaca	900

58

cctggaagca	aaatcatagt	gcaaaaatac	atttaagggtg	tgggtcaaaaa	taagtcttta	960
attggtaaat	aataagcatt	aattttttat	agcctgtatt	cacaattctg	cggtacctta	1020
ttgtacctaa	gggattctaa	agggtgtgtc	actgtataaa	acagaaagca	ctaggataca	1080
aatgaagctt	aattactaaa	atgtaattct	tgacactctt	tctataatta	gcgttcttca	1140
ccccacccc	cacccccacc	ccccttattt	tccttttgtc	tcctgggtgat	taggccaag	1200
tctgggagta	aggagaggat	taggtactta	ggagcaaaga	aagaagtagc	ttgggaacttt	1260
tgagatgata	cctaacatac	tgtactactt	gctttttaca	tgtgttagca	gaaaccagtg	1320
ggttataatg	tagaatgatg	tgctttctgc	ccaagtggta	attcatcttg	gtttgctatg	1380
ttaaaactgt	aaatacaaca	gaacattaat	aaatatctct	tgtgtagcac	ctttaaaaaa	1440
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	ccccgggggg	ggcccn		1487

<210> 93

<211> 1653

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (67)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (212)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1636)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1653)

<223> n equals a,t,g, or c

<400> 93

gagaccgagc	ccttcagcta	gctcgtcgc	tcgctctgct	tccttgcctgc	cggtcgcgca	60
tggcttnggc	gttggcggcg	ctggcggcgg	ctcgagccgc	ctgcgsagcc	ggtaccagca	120
gttgacgaat	gaagaagagt	ctggagaacc	tgaacaggct	gcaggtgatg	ctcctccacc	180
ttacagcagc	atttctgcag	agagcgcaca	tnattttgac	tacaaggatg	agtctggggt	240
tccaaaagccc	ccatcttaca	atgtagctac	aacactgccc	agttatgatg	aagcggagag	300
gaccaaggct	gaagctacta	tccttttggg	tcctgggaga	gatgaggatt	ttgtgggtcg	360
ggatgatatt	gatgatgctg	accagctgag	gataggaaat	gatgggattt	tcattgttaac	420
ttttttcatg	gcattcctct	ttaactggat	tgggtttttc	ctgtcttttt	gcctgaccac	480
ttcagctgca	ggaaggatg	gggccatttc	aggatttggg	ctctctctaa	ttaaatggat	540
cctgattgtc	aggttttcca	cctatttccc	tgcatattatg	aattctctct	caagaagcaa	600
gagaacacct	gcaggaagtg	aatcaagatg	cagaacacag	aggaataatc	acctgcttta	660
aaaaaataaa	gtactgttga	aaagatcatt	tctctctatt	tgttcctagg	tgtaaaattt	720
taatagttaa	tgcagaattc	tgtaatcatt	gaatcattag	tgggttaatt	ttgaaaaagc	780
tcttgcaatc	aagtcgtgta	tgtattaata	atgccttata	tattgtttgt	agtcatttta	840
agtacatga	gccatgtccc	tgtagtcggg	agggggcagt	cttgctttat	tcattctcca	900
tctcaaaatg	aacttggaat	taaaatttgt	aagatatgta	taatgctggc	catttttaag	960
gggtttttctc	aaaagttaaa	cttttggtat	gactgtgttt	ttgcacataa	tccatatttg	1020
ctgttcaagt	taatctagaa	atttattcaa	ttctgtatga	acacctggaa	gcaaaatcat	1080
agtgcaaaaa	tacatttaag	gtgtgggtcaa	aaataagtct	ttaattggta	aataataagc	1140
attaattttt	tatagcctgt	attcacaatt	ctcgggtacc	ttattgtacc	taagggtacc	1200

59

taaagggtgtt	gtcactgtat	aaaacagaaa	gcactaggat	acaaatgaag	cttaattact	1260
aaaaatgtaat	tcttgacact	ctttctataa	ttagcgttct	tcacccccac	ccccaccccc	1320
acccccctta	ttttcctttt	gtctcctggt	gattaggcca	aagtctggga	gtaaggagag	1380
gattaggtag	ttaggagcaa	agaaagaagt	agcttggaac	ttttgagatg	atccctaaca	1440
tactgtacta	cttgctttta	caatgtgtta	gcagaaacca	gtgggttata	atgtagaatg	1500
atgtgctttc	tgcccaagtg	gtaattcatt	ttgggttgct	atgttaaaac	tgtaaatata	1560
acagaacatt	aataaatatc	tcttgtgtag	caccttttaw	aaaaaaaaaa	aaaaaaaaaa	1620
aaaaaaaaaa	aaaaancccc	ggggggggcc	ccn			1653

<210> 94

<211> 1830

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (67)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (97)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (211)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1813)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1830)

<223> n equals a,t,g, or c

<400> 94

gcgaccgcgc	ccttcagcta	gctcgtctgc	tcgctctgct	tcctctgctgc	cggtctgcga	60
tggcttnggc	gttggcggcg	ctggcggcgg	tcgagcngcc	tgcsagccg	gtaccagcag	120
ttgcagaatg	aagaagagtc	tggagaacct	gaacaggctg	cagggtgatgc	tcctccacct	180
tacagcagca	tttctgcaga	gagcgcacat	nattttgact	acaaggatga	gtctggggtt	240
ccaaagcccc	catcttacaa	tgtagctaca	acactgcccc	gttatgatga	agcggagagg	300
accaaggctg	aagctactat	ccctttgggt	cctgggagag	atgaggattt	tgtgggtcgg	360
gatgattttg	atgatgctga	ccagctgagg	ataggaaatg	atgggatttt	catgttaact	420
tttttcatgg	cattcctctt	taactggatt	gggtttttcc	tgtctttttg	cctgaccact	480
tcagctgcag	gaaggtagtg	ggccatttca	ggatttggtc	tctctctaata	taaatggatc	540
ctgattgtca	ggttttccac	ctatttccct	ggatattttg	atgggtcagta	ctgggtctgg	600
tgggtgttcc	ttgttttagg	ctttctcctg	tttctcagag	gatttatcaa	ttatgcaaaa	660
gttcggaaga	tgccagaaac	tttctcaaat	ctccccagga	ccagagttct	ctttatttat	720
taaagatgtt	ttctggcaaa	ggccttctctg	catttatgaa	ttctctctca	agaagcaaga	780
gaacacctgc	aggaagtga	tcaagatgca	gaacacagag	gaataatcac	ctgctttaaa	840
aaaataaagt	actgttgaaa	agatcatttc	tctctatttg	ttcctagggtg	taaaatttta	900
atagttaatg	cagaattctg	taatcattga	atcatttagt	gttaattgttt	gaaaaagctc	960
ttgcaatcaa	gtctgtgatg	tattaataat	gccttatata	ttgtttgtag	tcatttttaag	1020

60

tagcatgagc	catgtccctg	tagtcggtag	ggggcagctc	tgctttattc	atcctccatc	1080
tcaaaatgaa	cttggaatta	aatattgtaa	gatatgtata	atgctggcca	ttttaaagg	1140
gttttctcaa	aagttaaact	tttgttatga	ctgtgttttt	gcacataatc	catatttgct	1200
gttcaagtta	atctagaaat	ttattcaatt	ctgtatgaac	acctggaagc	aaaatcatag	1260
tgcaaaaata	catttaaggt	gtggtcaaaa	ataagtcttt	aattggtaaa	taataagcat	1320
taatttttta	tagcctgtat	tcacaattct	gcggtacctt	attgtacctt	agggattcta	1380
aagggtgtgt	cactgtataa	aacagaaagc	actaggatac	aaatgaagct	taattactaa	1440
aatgtaattc	ttgacactct	ttctataatt	agcgtctctc	acccccaccc	ccacccccac	1500
cccccttatt	ttccttttgt	ctcctgggtg	ttaggccaaa	gtctgggagt	aaggagagga	1560
ttagggtactt	aggagcaaa	aaagaagtag	cttggaaactt	ttgagatgat	ccctaacata	1620
ctgtactact	tgctttttaca	atgtgttagc	agaaaccagt	gggttataat	gtagaatgat	1680
gtgctttctg	cccaagtggg	aattcatctt	gggttgctat	gttaaaactg	taataacaac	1740
agaacattaa	taaataatctc	ttgtgtagca	ccttttaaaa	aaaaaaaaaa	aaaaaaaaaa	1800
aaaaaaaaaa	aancccgggg	gggggccccn				1830

<210> 95
 <211> 1134
 <212> DNA
 <213> Homo sapiens

<400> 95						
tccatctaca	gtcctcacac	aggtattcag	gaataccagg	atggcgtgcc	caagattcca	60
acagcctgta	ttacgggtga	agatgcagaa	atgatgtcaa	gaatggcttc	tcatgggatc	120
aaaattgtca	ttcagctaaa	gatgggggca	aagacctacc	cagatactga	ttccttcaac	180
actgtagcag	agatcactgg	gagcaaatat	ccagaacagg	ttgtactggg	cagtggacat	240
ctggacagct	gggatgttgg	gcagggtgcc	atggatgatg	gcggtggagc	ctttatatca	300
tgggaagcac	tctcacttat	taaagatctt	gggctgcgtc	caaagaggac	tctgcggtcg	360
gtgctctgga	ctgcagaaga	acaagggtga	gttgggtgct	tccagtatta	tcagttacac	420
aaggtaata	tttccaaacta	cagtctgggtg	atggagtctg	acgcagggaac	cttcttacct	480
actgggctgc	aattcactgg	cagtgaagag	gccaggggcat	catggaggag	gttatgagcc	540
tgctgcagcc	cctcaatata	actcaggtcc	tgagccatgg	agaagggaca	gacatcaact	600
tttgatccca	agctggagtg	cctggagcca	gtctacttga	tgacttatac	aagtatttct	660
tottccatca	ctcccacgga	gacaccatga	ctgtcatgga	tccaaagcag	atgaatgttg	720
ctgctgctgt	ttgggctgtt	gtttcttatg	ttgttgacga	catggaagaa	atgctgccta	780
ggctcctagaa	acagtaagaa	agaaacgttt	tcattgctct	ggccagggaat	cctgggtctg	840
caacttttga	aaactcctct	tcacataaca	atttcattcca	attcatcttc	aaagcacaac	900
tctatttcat	gctttctgtt	attatcttct	ttgatacttt	ccaaattctc	tgcatctctg	960
aaaaaggaat	cattctcccc	tccttccccac	cacatagaat	caacatatgg	tagggattac	1020
agtgggggca	tttctttata	tcacctctta	aaaacattgt	ttccacttta	aaagtaaac	1080
cttaataaat	ttttggaaga	tctctgaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaa	1134

<210> 96
 <211> 1772
 <212> DNA
 <213> Homo sapiens

<400> 96						
tcgacccacg	cgtccgggag	gatccccagc	cgggtcccaa	gcctgtgcct	gagcctgagc	60
ctgagcctga	gccgagccgg	gagccgggtc	cgggggctcc	gggctgtggg	accgctgggc	120
ccccagcgat	ggcgaccctg	tggggaggcc	ttcttcggct	tggtcctctg	ctcagcctgt	180
cgtgcctggc	gctttccgtg	ctgctgctgg	cgactgtca	gacgccgcca	agaatttcga	240
ggatgtcaga	tgtaaatgta	tctgccctcc	ctataaagaa	aaattctggg	catattttata	300
ataagaacat	atctcagaaa	gattgtgatt	gccttcatgt	tgtggagccc	atgcctgtgc	360
gggggcctga	tgtagaagca	tactgtctac	gctgtgaatg	caaatatgaa	gaaagaagct	420
ctgtocacaat	caaggttacc	attataattt	atctctccat	tttgggcctt	ctacttctgt	480
acatgggtata	tcttactctg	gttgagccca	tactgaagag	gcgcctcttt	ggacatgcac	540

61

agttgataca	gagtgatgat	gatattgggg	atcaccagcc	ttttgcaaat	gcacacgatg	600
tgctagcccg	ctcccgcagt	cgagccaacg	tgctgaacaa	ggtagaatat	ggcacagcag	660
cgctggaagc	ttcaagtcca	agagcagcga	aaagtctgtc	tttgaccggc	atgttgtcct	720
cagctaattg	gggaattgaa	ttcaagggtga	ctagaaagaa	acaggcagac	aactggaaag	780
gaactgactg	ggttttgctg	ggtttcattt	taataccttg	ttgatttcac	caactggtgc	840
tggaagattc	aaaactggaa	gkaaaaaactt	gcttgatttt	tttttcttgt	taacgtaata	900
atagagacat	ttttaaaagc	acacagctca	aagtcagcca	ataagtcttt	tcctatttgt	960
gacttttact	aataaaaaata	aatctgcctg	taaaaataat	taaaaaatcc	tttacctgga	1020
acaagcactc	tctttttcac	cacatagttt	taacttgact	ttccaagata	attttcaggg	1080
tttttgttgt	tggtgttttt	tgtttgtttg	ttttggtggg	agaggggagg	gatgcctggg	1140
aagtgggttaa	caactttttt	caagtcactt	tactaaacaa	acttttgtaa	atagacctta	1200
ccttctattt	tcgagtttca	tttataattt	gcagtgtagc	cagcctcctc	aaagagctga	1260
cttactcatt	tgacttttgc	actgactgta	ttatctgggt	atctgctgtg	tctgcacttc	1320
atgggtaaacg	ggatctaaaa	tgccctgggtg	cttttcacaa	aaagcagatt	ttcttcattg	1380
actgtgatgt	ctgatgcaat	gcactcctaga	acaaaactggc	catttgctag	tttactctaa	1440
agactaaaca	tagtcttggt	gtgtgtgggc	ttactcatct	tctagtacct	ttaaggacaa	1500
atcctaagga	cttggaacct	tgcaataaag	aaattttatt	ttaaacccaa	gcctccctgg	1560
attgataata	tatacacatt	tgtcagcatt	tccgggtcgtg	gtgagaggca	gctgtttgag	1620
ctccaatgtg	tgacagcttg	aactagggct	gggggtgtgg	gtgcctcttc	tgaagggtct	1680
aaccattatt	ggataactgg	ctttttttct	tcctcttggt	aatgtaacaa	taaaaataat	1740
ttttgaaaca	tcaaaaaaaa	aaaaaaaaaa	aa			1772

<210> 97

<211> 2381

<212> DNA

<213> Homo sapiens

<400> 97

ccacgcgtcc	cgcaaggcca	gttctagtgt	agagagaaaa	aggagccggc	agcggctctt	60
acgcgtcccg	gggctgcgcg	ccactctctc	ggccggtaac	gcgggtgctt	gcggctgtcg	120
tcaagcgcgg	cgttgggccc	gcggggcggg	gctgaggggc	tgccatggcg	gcggcggggc	180
ggctcccgag	ctcctgggccc	ctcttctcgc	cgctcctcgc	agggcttgca	ctactgggag	240
tcggggccggt	cccagcgcgg	gcgctgcaca	acgtcacggc	cgagctcttt	ggggccgagg	300
cctggggcac	ccttgccggt	ttcggggacc	tcaactccga	caagcagacg	gatctcttcg	360
tgctgcggga	aagaaatgac	ttaatcgtct	ttttggcaga	ccagaatgca	ccctatttta	420
aacccaaagt	aaaggatatc	ttcaagaatc	acagtgcatc	gataacaagt	gtagtccctg	480
gggattatga	tggagattct	caaatggatg	tccttctgac	atatcttccc	aaaaattatg	540
ccaagagtga	attaggagct	gttatcttct	ggggacaaaa	tcaaacatta	gatcctaaca	600
atatgaccat	actcaatagg	acttttcaag	atgagccact	aattatggat	ttcaatgggtg	660
atctaattcc	tgatattttt	ggtatcacaa	atgaatccaa	ccagccacag	atactattag	720
gagggaaatt	atcatggcat	ccagcattga	ccactacaag	taaaatgcga	attccacatt	780
ctcatgcatt	tattgatctg	actgaagatt	ttacagcaga	tttattcctg	acgacattga	840
atgccaccac	tagtaccttc	cagtttgaaa	tatgggaaaa	tttggatgga	aacttctctg	900
tcagtactat	attggaaaaa	cctcaaaaata	tgatgggtgt	tggacagtca	gcatttgacc	960
actttgatgg	agatggacac	atggatcatt	tactgccagg	ctgtgaagat	aaaaattgcc	1020
aaaagagtac	catctactta	gtgagatctg	ggatgaagca	gtgggttcca	gtcctacaag	1080
atttcagcaa	taagggcaca	ctctggggct	ttgtgccatt	tgtggatgaa	cagcaaccaa	1140
ctgaaatacc	aattccaatt	acccttcata	ttggagacta	caatatggat	ggctatccag	1200
acgctctggt	cataactaag	aacacatctg	gaagcaacca	gcaggccttt	ttactggaga	1260
acgtcccttg	taataatgca	agctgtgaag	aggcgcgtcg	aatgttttaa	gtctactggg	1320
agctgcacaga	cctaaatcaa	atgaaggatg	ccatgggtgc	caccttcttt	gacatttacg	1380
aagatggaat	cttggaacatt	gtagtgtctaa	gtaaaggata	tacaaagaat	gatttttgcca	1440
ttcatacact	aaaaaataac	tttgaagcag	atgcttattt	tgttaaagtt	attgttctta	1500
gtgggtctgtg	ttctaattgac	tgctctcgta	gataacaccc	tttggagtga	atcaacctgg	1560
accttatatc	atgtatacaa	ctgtagatgc	aaatgggtat	ctgaaaaatg	gatcagctgg	1620
ccaactcagc	caatccgcac	attttagctct	ccaactacca	tacaacgtgc	ttgggttagg	1680
toggagcgca	aatttttctg	accatctcta	cgttggtatt	ccccgtccat	ctggagaaaa	1740

atctatacga	aaacaagagt	ggactgcaat	cattccaaat	ccccagctaa	ttgtcattcc	1800
ataccctcac	aatgtccctc	gaagttggag	tgccaaactg	tatcttacac	caagtaatat	1860
tgttctgctt	actgctatag	ctctcatcgg	tgtctgtgtt	ttcatcttgg	caataattgg	1920
cattttacat	tggcaggaaa	agaaagcaga	tgatagagaa	aaacgacaag	aagcccaccg	1980
gtttcatttt	gatgctatgt	gacttgccct	taatattaca	taatggaatg	gctgttcact	2040
tgattagtgt	aaacacaaat	tctggcttga	aaaaataggg	gagattaaat	attatttata	2100
aatgatgtat	cccatggtaa	ttattggaaa	gtattcaaat	aaatatgggt	tgaatatgtc	2160
acaaggtctt	tttttttaaa	gcactttgta	tataaaaatt	tgggttctct	attctgtagt	2220
gctgtacatt	tttgttcctt	tgtggaatgt	gttgcatgta	ctccagtgtt	tgtgtattta	2280
taatcttatt	tgcattcatga	tgatggaaaa	agttgtgtaa	ataaaaaata	ttaaatgagc	2340
aggaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	a		2381

<210> 98
 <211> 1955
 <212> DNA
 <213> Homo sapiens

<400> 98						
ggcagcagtg	ccatccctgt	atttgctgcc	atgctcttcc	ttttctccat	ggctacactg	60
ttgaggacca	gcttcagtga	ccctggagtg	attcctcggg	cgctaccaga	tgaagcagct	120
ttcatagaaa	tggagataga	agctaccaat	ggtgcggtgc	cccagggcca	gcgaccaccg	180
cctcgtatca	agaatttcca	gataaacaac	cagattgtga	aactgaaata	ctgttacaca	240
tgaagatct	tccggcctcc	ccgggcctcc	cattgcagca	tctgtgacaa	ctgtgtggag	300
cgcttcgacc	atcactgccc	ctgggtgggg	aattgtgttg	gaaagaggaa	ctaccgctac	360
ttctacctct	tcactccttc	tctctccctc	ctcacaatct	atgtcttcgc	cttcaacatc	420
gtctatgttg	ccctcaaate	tttgaaaatt	ggcttcttgg	agacattgaa	aggaaactcc	480
tggaaactgt	ctagaagtcc	tcatttgctt	ctttacactc	tgggtccgtcg	tgggactgac	540
tggatttcat	actttcctcg	tggctctcaa	ccagacaacc	aatgaaagac	atcaaaggat	600
catggacagg	gaagaatcgc	gtccagaatc	cctacagcca	tggcaatatt	gtgaagaact	660
gctgtgaagt	gctgtgtggc	cccttgcccc	ccagtgtgct	ggatcgaagg	ggtattttgc	720
cactggagga	aagtgggaagt	cgacctccca	gtactcaaga	gaccagtagc	agcctcttgc	780
cacagagccc	agccccca	gaacacctga	actcaaatga	gatgccggag	gacagcagca	840
ctcccgaaga	gatgccacct	ccagagcccc	cagagccacc	acaggaggca	gctgaagctg	900
agaagtagcc	tatctatgga	agagactttt	gtttgtgttt	aattagggct	atgagagatt	960
tcaggtgaga	agttaaacct	gagacagaga	gcaagtaagc	tgtccctttt	aactgttttt	1020
ctttgggtct	tagtcaccca	gttgcacact	ggcattttct	tgtctgcaagc	ttttttaaat	1080
ttctgaactc	aaggcagtg	cagaagatgt	cagtcacctc	tgataactgg	aaaaatgggt	1140
ctcttggggc	ctggcactgg	ttctccatgg	cctcagccac	aggggtccct	tggacccccct	1200
ctcttccctc	cagatcccag	ccctcctgct	tgggggtcact	gggtctcatc	tggggctaaa	1260
agttttctgag	actggctcaa	atcctcccaa	gctgtctgac	gtgctgagtc	cagaggcagt	1320
cacagagacc	tctggccagg	ggatccctaac	tgggttcttg	gggtcttcag	gactgaagag	1380
gagggagagt	ggggtcagaa	gattctctctg	gccaccaagt	gccagcattg	cccacaaatc	1440
cttttaggaa	tgggacaggt	accttccact	agttgtatatt	attagtgtag	cttctccttt	1500
gtctcccatc	cactctgaca	ccttaagccc	cactcttttc	ccattagata	tatgtaagta	1560
gttgtagtag	agataataat	tgacatttct	cgtagactac	ccagaaactt	ttttaatacc	1620
tgtgccattc	tcaataagaa	tttatgagat	gccagcggca	tagcccttca	cactctctgt	1680
ctcatctctc	ctcctttctc	attagccccct	tttaatttgt	ttttcctttt	gactcctgct	1740
cccattagga	gcaggaatgg	cagtaataaa	agtctgcact	ttggtcattt	cttttctctca	1800
gaggaagcct	gagtgtctcac	ttaaacacta	tcccctcaga	ctccctgtgt	gaggcctgca	1860
gaggccctga	atgcacaaat	gggaaccacaa	ggcacagaga	ggctctctctc	tcctctctctc	1920
tcccccgatg	taccctcaaa	aaaaaaaaaa	aaaaa			1955

<210> 99
 <211> 1958
 <212> DNA
 <213> Homo sapiens

<400> 99

ccacgcgtcc	ggggcggtcc	tggtcgtgag	aggggagccc	caggggagct	ggggcagcat	60
gactgggggtg	ataaatggcc	ggaaatttgg	cgtggccaca	ctcaacacca	gcgtgatgca	120
ggaggcacac	tccgggggtca	gcagcatcca	cagcagcatc	cgccatgtcc	cagcaaacgt	180
ggggcctctg	atgcgggtgc	tcgtgggtcac	catcgcccc	atctactggg	ccctggccag	240
agagagtggg	gaagccctga	atggccactc	tctgactggg	ggcaagtcc	ggcagagtca	300
cacgtggagt	ttgtacag	gagctgctca	cgatgaccca	gtggccccggg	gtctggatcc	360
cgatggcctc	ctgctcctcg	acgtgggtgt	caatggcggt	gtccccggac	gagcctggct	420
gacgcagatc	ttcaagtga	ggactttgaa	gaagcactac	gtgcaaacaa	gggcctggcc	480
agctgttcgt	gggctccaca	cagcgcttct	tccaggcgcg	cctccccctg	ttcctacgct	540
gcaaccacag	catccagtac	aacgcggccc	ggggccccc	gccccagctg	gtgcagcacc	600
tgccggcctc	agctatcagc	tcggcctttg	atccagaggc	cgaggccctg	cgcttccagc	660
tcgtacacag	cctgcaggcg	gaggagaacg	aggtcggctg	ccccgagggc	tttgagctgg	720
actcccagg	agcgttttgt	gtggatgtgg	acgagtgtgc	gtgggatgct	cacctctgcc	780
gagaggggaca	gcgctgtgtg	aacctgtctg	ggtcctaccg	ctgcctcccc	gactgtgggc	840
ctggcttccg	ggtggctgat	ggggccggct	gtgaaaatgt	ggacgaatgc	ctggaagggg	900
ttggacgact	gtcactacaa	ccagctctgc	gagaaccccc	caggcgggtca	ccgctgcagc	960
tgccccagg	gttaccggat	gcagggcccc	agcctgccct	gcctagatgt	caatgagtg	1020
ctgcagctgc	ccaaggcctg	cgccctaccg	tgccacaacc	tccagggcag	ctaccgctgc	1080
ctgtgcccc	caggccagac	cctccttcgc	gacggcaagg	cctgcacctc	actggagcgg	1140
aatggacaaa	atgtgaccac	cgtcagccac	cgaggccctc	tattgccctg	gctgcggccc	1200
tgggcctcga	tccccggtag	ctcctaccac	gcctgggtct	ctctccgtcc	gggtcccatg	1260
gccttgagca	gtgtgggccc	ggcctgggtg	cctcctgggt	tcattcaggca	gaacggagtc	1320
tgacacagacc	ttgacgagtg	ccgcgtgagg	aacctgtgtc	agcacgcctg	ccgcaaacat	1380
gagggcagct	accagtgcct	gtgccccgcc	ggctaccgtc	tgctccccag	cgggaagaac	1440
tgccaggaca	tcaacgagtg	cgaggaggag	agcatcgagt	gtggaccggg	ccagatgtgc	1500
ttcaacaccc	gtggcagcta	ccagtgtgtg	gacacaccc	gtcctgccac	ctaccggcag	1560
ggccccagcc	ctgggacgtg	cttcggggcg	tgctgcagg	actgcggcac	gggcggccct	1620
tctacgctgc	agtaccggct	gctgccgctg	cccctggggc	tgccgcgcc	ccacgacgtg	1680
gcccgcctca	ccgccttctc	cgagggtcgg	gtccccgcc	accgcaccga	gtcagcatg	1740
ctggagcccc	acccccgcag	ccccttcggc	ctgcgtccgc	tgccgcgggg	ccttggcgcg	1800
gtctacaccc	gtcgcgcgct	cacccgcgcc	ggcctctacc	ggctcaccgt	gcgtgctgcg	1860
gcaccgcgcc	accaaagcgt	cttcgtcttg	ctcatgcggc	tgtcccccta	cccctactaa	1920
acgggagagg	gcattggcgg	ccgctctaga	ggatccct			1958

<210> 100

<211> 2444

<212> DNA

<213> Homo sapiens

<400> 100

ttacgccaag	ctggcacgag	caatgaaaga	gttaatctct	ttggctgggc	ctacagatga	60
catacagagt	acagtccccc	aggttcatgc	tttaaatatc	cttagagcat	tggtcagaga	120
tacgcgcctg	ggagaaaata	ttattcctta	tggttgctgat	ggagctaagg	ctgcaattct	180
gggttttaca	tcaccgggtct	gggcagtgcg	aaattcatcc	acacttctct	ttagtgcctt	240
gatcacaaga	atthttggag	ttaaaagggc	aaaggatgaa	cattccaaaa	caaatagaat	300
gacagggaga	gagtttttct	ctcgtttccc	agaactctat	ccttttcttc	tcaaacagtt	360
ggaaactgta	gccaatacag	tagacagtga	tatgggagaa	ccaaatcgtc	atccaagcat	420
gtttctctta	cttttgggtg	tggagagact	ctacgcttcc	ccgatggatg	gtacttcttc	480
tgctctcagc	atgggacctt	ttgttccctt	cattatgagg	tgtgggtcact	cacctgtcta	540
ccactcccgt	gaaatggcag	ctcgtgcctt	ggctccattt	gttatgatag	atcacattcc	600
taataccatt	cgaactctgt	tgtccacact	ccccagctgc	actgaccagt	gtttccgggc	660
aaaaccacat	tcattggggac	acttctccag	gtttttccat	ttgttgcaag	cctactcaga	720
ctccaaaaca	cggaaacgaat	tcagacttcc	agcacgagct	gactgacatc	actgtttgta	780
ccaaagccaa	actctggctg	gccaaagagg	aaaatccatg	tttggtgacc	agagctgtat	840
atattgatat	tctcttcccta	ttgacttgct	gcctcaacag	atctgcaaaag	gacaaccagc	900

cagttcttggga	gagtccttggc	ttctgggaag	aaattcaaag	ggaattatct	caggatcaga	960
agctgataac	gggattccct	tgggccttca	aggtgccagg	cctgcccag	tacctccaga	1020
gcctcaccag	actagccatt	gctgcagtgt	gggccgcggc	agccaagagt	ggagagcggg	1080
agacgaatgt	ccccatctct	ttctctcagc	tgtagaatc	tgccctccct	gaagtgcgct	1140
cactaacact	ggaagccctc	ttggaaaagt	tcttagcagc	agactctgga	cttgggagaga	1200
agggcgtgcc	acccttgctg	tgcaacatgg	gagagaagtt	cttattgttg	gccatgaagg	1260
aaaatcaccc	agaatgcttc	tgcaagatac	tgaaaattct	acactgcatg	gaccctggtg	1320
agtggcttcc	ccagacggag	cactgtgtcc	atctgacccc	aaaggagttc	ttgatctgga	1380
cgatggatat	tgcttccaat	gaaagatctg	aaattcagag	tgtagctctg	agacttgctt	1440
ccaaagtcat	ttcccaccac	atgcagacat	gtgtggagaa	cagggaattg	atagctgctg	1500
agctgaagca	gtgggttcag	ctggtcacat	tgatcatgtg	agaccatctt	cctacagagt	1560
ctaggctggc	cgctggtgaa	gtcctcacca	gtactacacc	acttttcttc	accaaccccc	1620
atcctattct	tgagttgcag	gatacacttg	ctctctggaa	gtgtgtcctt	acccttctgc	1680
agagttagga	gcaagctgtt	agagatgcag	ccacggaaac	cgtagacaact	gccatgtcac	1740
aagaaaaatac	ctgccagtc	acagagtttg	ccttctgcc	ggtggatgcc	tccatcgctc	1800
tggccctggc	cctggccgtc	ctgtgtgac	tgctccagca	gtgggaccag	ttggccctg	1860
gactgcccat	cctgctggga	tggtgttgg	gagagagtga	tgacctcgtg	gcctgtgtgg	1920
agagcatgca	tcaggtggaa	gaagactacc	tgtttgaaaa	agcagaagtc	aacttttggg	1980
ccgagacctt	gatctttgtg	aaatacctct	gcaagcacct	cttctgtctc	ctctcaaagt	2040
ccggctggcg	tcaccaagc	cctgagatgc	tctgtcacct	tcaaaggatg	gtgtcagagc	2100
agtgccacct	cctgtctcag	ttcttcagag	agcttccacc	agctgctgag	tttgtgaaga	2160
cagtggagtt	cacaagacta	cgcatccaag	aggaaaggac	tttggcttgc	ttgaggctgc	2220
tggccctttt	ggaaggaaag	gaaggggaag	acaccctagt	tctcagtggt	tgggactctt	2280
atgcagaatc	gaggcagtta	actcttccaa	gaacagaagc	ggcatgttga	agaaaatctg	2340
ggggattggg	atgggggtat	gtgtggattt	ttcctccact	aaatctgcag	gaaacatggt	2400
gaacataaat	tcaaaaattt	tatccccaaa	aaaaaaaaaa	aaaa		2444

<210> 101

<211> 2709

<212> DNA

<213> Homo sapiens

<400> 101

ggcacgagat	ttcctacagg	tgaaacgcca	tcattaggat	tcaactgtaac	gttagtgcta	60
ttaaactcac	tagcattttt	attaatggcc	gttatctaca	ctaagctata	ctgcaacttg	120
gaaaaagagg	acctctcaga	aaactcacaa	tctagcatga	ttaaagcatgt	cgcttggcta	180
atcttcacca	attgcattct	ttctgcctt	gtggcggttt	tttcatttgc	accattgatc	240
actgcaatct	ctatcagccc	cgaaataatg	aagctctgta	ctctgatatt	ttttccatgc	300
ctgcttgcct	gaatccagtc	ctgtatgttt	tcttcaaccc	aaagtttaaa	gaagactgga	360
agttactgaa	gcgacgtgtt	accaagaaaa	gtggatcagt	ttcagtttcc	atcagtagcc	420
aagggtggtg	tctggaacag	gattttctact	acgactgtgg	catgtactca	catttgcagg	480
gcaacctgac	tgtttgcgac	tgctgcgaat	cgtttctttt	aacaaagcca	gtatcatgca	540
aacacttgat	aaaatcacac	agctgtcctg	cattggcagt	ggcttcttgc	caaagacctg	600
agggctactg	gtccgactgt	ggcacacatt	cggtccactc	tgattatgca	gatgaagaag	660
attcctttgt	ctcagacagt	tctgaccagg	tgcaggcctg	tgacagagcc	tgcttctacc	720
agagttagagg	attccctttg	gtgcgctatg	cttacaatct	accaagagtt	aaagactgaa	780
ctactgtgtg	tgtaaccgtt	tcctccgtca	accaaaatca	gtgtttatag	agtgaacctt	840
attctcatct	ttcatctggg	aagcacttct	gtaatcactg	cctggtgtca	cttagaagaa	900
ggagaggtgg	cagttttatt	ctcaaaccag	tcattttcaa	agaacagggtg	cctaaattat	960
aaattggtga	aaaatgcaat	gtccaagcaa	tgtatgatct	gtttgaaaca	aatatatgac	1020
ttgaaaagga	tcttaggtgt	agtagagcaa	tataatgtta	gttttttctg	atccataaga	1080
agcaaattta	tacctatttg	tgtattaagc	acaagataaa	gaacagctgt	taatattttt	1140
taaaaattct	attttttaaaa	tgtgattttc	tataactgaa	gaaaaatatc	ttgctaattt	1200
tacctaatgt	ttcatccttt	aatctcagga	caacttactg	cagggccaaa	aaagggactg	1260
tcccagctag	acctgtgaga	gtatacatag	gcattacttt	attatgtttt	cacttgccat	1320
ccttgacata	agagaactat	aaattttgtt	taagcaattt	ataaatctaa	aacctgaaga	1380
tgtttttaaa	acaatattaa	cagctgttag	gttataaaaa	tagctggaca	ttgtttttca	1440

65

gtcattatac	attgcttttg	tccaatcagt	aattttttct	taagtgtttt	gtgattacac	1500
tactagaaaa	aaagtaaaag	gctaattgct	gtgtgggttt	agtcgatttg	gctaaactac	1560
taactaatgt	gggggtttta	tagtatctga	gggatttggg	ggcttcatgt	aatgttctca	1620
ttaatgaata	cttcctaata	tcgttggctc	tactaatatt	ttccaatttg	ctgggatgtc	1680
acctagcaat	agcttggatt	atatagaaag	taaactgtgg	tcaatacttg	catttaatta	1740
gacgaaacgg	ggagtaatta	tgacacgaag	tacttaatgt	ttatttctta	gtgagctgga	1800
ttatcttgaa	cctgtgctat	taaatggaaa	tttccataca	tcttccccat	actatttttt	1860
ataaaagagc	ctattcaata	gctcagaggt	tgaactctgg	ttaaacaaga	taatatgtta	1920
ttataaaaaa	tagaagaaga	aagaataaag	cttagtcctg	tgtcttttaa	aaattaaaaa	1980
ttttacttga	ttccccatct	atgggcttta	gacctattac	tgggtggagt	cttaaagtta	2040
taattgttca	atatgttttt	tgaacagtgt	gctaaatcaa	tagcaaacc	actgccatat	2100
tagttattct	gaatatacta	aaaaaatcca	gctagattgc	agtttaataa	ttaaactgta	2160
catactgtgc	atataatgaa	tttttatctt	atgtaaatta	tttttagaac	acaagtggg	2220
aaatgtggct	tctgttcatt	tcgtttaatt	aaagctacct	cctaaactat	agtggctgcc	2280
agtagcagac	tgtaaatttg	tggtttatat	actttttgca	ttgtaaatag	tctttgttgt	2340
acattgtcag	tgtaataaaa	acagaatctt	tgtatatcaa	aatcatgtag	tttgtataaa	2400
atgtgggaag	gatttattta	cagtgtgttg	taattttgta	aggccaacta	tttacaagtt	2460
ttaaaaattg	ctatcatgta	tatttacaca	tctgataaat	attaaatcat	aacttggtta	2520
gaaactccta	attaaaagg	tttttccaaa	attcaggtta	ttgaaaactt	ttcattttat	2580
tcattttaaaa	actagaataa	cagatatata	aaagtgttaa	tctttgtgct	atatgggtatg	2640
aaatacaata	ttgtactcag	tgttttgaat	tattaaagtt	tctagaaagc	aaaaaaaaaa	2700
aaaaaaaaaa						2709

<210> 102

<211> 1722

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (401)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (695)

<223> n equals a,t,g, or c

<400> 102

gggacgcgc	tgctctgctg	tcaccaagag	ctggagacac	catctccac	cgagagtcac	60
ggccccattg	gccctgcacc	tctctgtcct	cgtcccatc	ctcctcagcc	tggtggcctc	120
ccaggactgg	aaggctgaac	gcagccaaga	ccccttcgag	aaatgcatgc	aggatcctga	180
ctatgagcag	ctgctcaagg	tcaccatcct	ggaggcagat	aacaggatcg	ggggccgcat	240
cttcacctac	cgggaccaga	wyacgggctg	gattggggag	ctgggagcca	tgcgcatgcc	300
cagctctcac	aggatcctcc	acaagctctg	ccagggcctg	gggctcaacc	tgaccaagtt	360
caccagtagc	gacaagaaca	cgtggacgga	ggtgcacgaa	ntgaagctgc	gcaactatgt	420
ggtggagaag	gtgcccgaag	agctgggcta	cgccttgctg	ccccaggaaa	agggccactc	480
gcccgaagac	atctaccaga	tggtctctca	ccaggccctc	aaagacctca	aggcactggg	540
ctgcagaaag	gcgatgaaga	agtttgaaag	gcacacgctc	ttggaatc	ttctcgggga	600
ggggaaacctg	agccggcccg	ccgtgcagct	tctgggagac	gtgatgtccg	aggatggctt	660
cttctatctc	agcttcgccc	aggccctccg	ggccnacagc	tgctcagcg	acagactcca	720
gtacagcccg	atcgtgggtg	gctgggacct	gctgccgcgc	gegctgctga	gctcgtgtgc	780
cgggcttgtg	ctgttgaaag	cgcccggtgt	ggcgatgacc	cagggaccgc	acgatgtgca	840
cgtgcagatc	gagacctctc	ccccggcgcg	gaatctgaag	gtgctgaagg	ccgacgtggg	900
gctgctgacg	gcgagcggac	cggcggtgaa	gcgcacacc	ttctcgccgc	gctgccccgc	960
cacatgcagg	aggcgctgcg	gaggctgcac	tacgtgccgg	ccaccaaggt	gttcctaagc	1020
ttccgcaggc	ccttctggcg	cgaggagcac	attgaaggcg	gccactcaaa	caccgatcgc	1080

66

```

ccgtcgcgca tgattttcta cccgccgcgc cgcgagggcg cgctgctgct ggctcgtac 1140
acgtgggtcgg acgcggcggc agcgttcgcc ggcttgagcc gggaagaggg gttgcgcttg 1200
gcgctcgacg acgtggcggc attgcacggg cctgtcgtgc gccagctctg ggacggcacc 1260
ggcgtcgtca agcgttgggc ggaggaccag cacagccagg gtggctttgt ggtacagmcg 1320
ccggcgctct ggcaaaccga aaaggatgac tggacgggcc cttatggccg catctacttt 1380
gccggcgagc acaccgccta cccgcacggc tgggtggaga cggcggtcaa gtcggcgctg 1440
cgcgccgcca tcaagatcaa cagccggaag gggcctgcat cggacacggc cagccccgag 1500
gggcacgcat ctgacatgga ggggcagggg catgtgcatg ggggtggccag cagccccctg 1560
catgacctgg caaaggaaga aggcagccac cctccagtcc aaggccagtt atctctccaa 1620
aacacgaccc acacgaggac ctgcatttaa agtattttcg gaaaaaaaaa aaaaaaaaaa 1680
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaagggcgg cc 1722

```

<210> 103

<211> 106

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (14)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (29)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 103

```

Met Gly Ser Leu Ser Gly Cys Ala Leu Pro Phe Cys Leu Xaa Val Phe
  1             5             10             15

```

```

Phe Leu Thr Val Ser Pro Ser Ala Val Gly Leu Leu Xaa Phe Ala Gly
          20             25             30

```

```

Gly Pro Leu Gln Thr Leu Phe Ala Trp Val Ser Pro Val Glu Ala Ala
  35             40             45

```

```

Glu Gln Gln Arg Leu Leu Pro Val Leu Ser Ser Gly Ser Phe Val Ser
  50             55             60

```

```

Glu Gly Thr Cys Gln Met Pro Ala Arg Ala Leu Leu Tyr Glu Val Ser
  65             70             75             80

```

```

Val Gly Pro Tyr Trp Glu Ile Pro Pro Ser Gln Asp Thr Arg Arg Ser
          85             90             95

```

```

Gly Thr Tyr Leu Arg Arg Gln Ser Asp Pro
          100             105

```

<210> 104

<211> 86

<212> PRT

<213> Homo sapiens

<400> 104

Met Thr Leu Pro Ser Arg Ala Leu Ala Ser Leu Gly Val Gly Val Trp

1 5 10 15
 Gly Met Leu Arg Leu Asn Gln Val Thr Val Ser Cys Gly Gly Ser Arg
 20 25 30
 Trp Ser Ser Arg Val Ala Leu Gly Ala Phe Ser Trp Val Cys Gly Val
 35 40 45
 Ala Leu Val Leu Gln Pro Ser Gly Gly Gly Leu Gly Leu Thr Ser Pro
 50 55 60
 Ser Glu Gly Cys Trp Glu Gly Glu Leu Ala Leu Ala Val Leu Arg Ala
 65 70 75 80
 Pro Gly Gly Ser Pro Ser
 85
 <210> 105
 <211> 302
 <212> PRT
 <213> Homo sapiens
 <400> 105
 Met Ala Arg Ala Arg Gly Ser Pro Cys Pro Pro Leu Pro Pro Gly Arg
 1 5 10 15
 Met Ser Trp Pro His Gly Ala Leu Leu Phe Leu Trp Leu Phe Ser Pro
 20 25 30
 Pro Leu Gly Ala Gly Gly Gly Gly Val Ala Val Thr Ser Ala Ala Gly
 35 40 45
 Gly Gly Ser Pro Pro Ala Thr Ser Cys Pro Val Ala Cys Ser Cys Ser
 50 55 60
 Asn Gln Ala Ser Arg Val Ile Cys Thr Arg Arg Asp Leu Ala Glu Val
 65 70 75 80
 Pro Ala Ser Ile Pro Val Asn Thr Arg Tyr Leu Asn Leu Gln Glu Asn
 85 90 95
 Gly Ile Gln Val Ile Arg Thr Asp Thr Phe Lys His Leu Arg His Leu
 100 105 110
 Glu Ile Leu Gln Leu Ser Lys Asn Leu Val Arg Lys Ile Glu Val Gly
 115 120 125
 Ala Phe Asn Gly Leu Pro Ser Leu Asn Thr Leu Glu Leu Phe Asp Asn
 130 135 140
 Arg Leu Thr Thr Val Pro Thr Gln Ala Phe Glu Tyr Leu Ser Lys Leu
 145 150 155 160
 Arg Glu Leu Trp Leu Arg Asn Asn Pro Ile Glu Ser Ile Pro Ser Tyr
 165 170 175
 Ala Phe Asn Arg Val Pro Ser Leu Arg Arg Leu Asp Leu Gly Glu Leu

68

180	185	190
Lys Arg Leu Glu Tyr Ile Ser Glu Ala Ala Phe Glu Gly Leu Val Asn		
195	200	205
Leu Arg Tyr Leu Asn Leu Gly Met Cys Asn Leu Lys Asp Ile Pro Asn		
210	215	220
Leu Thr Ala Leu Val Arg Leu Glu Glu Leu Glu Leu Ser Gly Asn Arg		
225	230	235 240
Leu Asp Leu Ile Arg Pro Gly Ser Phe Gln Gly Leu Thr Ser Leu Arg		
245	250	255
Lys Leu Trp Leu Met His Ala Gln Val Ala Thr Ile Glu Arg Asn Ala		
260	265	270
Phe Asp Asp Leu Lys Ser Leu Glu Glu Leu Asn Leu Ser His Asn Asn		
275	280	285
Leu Met Ser Leu Pro His Asp Leu Phe Thr Pro Leu His Arg		
290	295	300

<210> 106

<211> 56

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (56)

<223> Xaa equals stop translation

<400> 106

Met Pro Ser Ser Trp Leu Pro Gly Cys Phe Val Leu Leu Cys Leu Val
1 5 10 15

Ala Val Gly Cys Gln Leu Arg Glu Trp Gly Val Gly Gly Val Ser Ala
20 25 30

Val Gly Leu Leu Ala Leu Pro His Leu Gln Val Leu Gly Met Arg Gly
35 40 45

Arg Gly Leu Ile Ser Gly Gly Xaa
50 55

<210> 107

<211> 189

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (94)

<223> Xaa equals any of the naturally occurring L-amino acids

69

<400> 107

Met Ala Leu Leu Ser Arg Pro Ala Leu Thr Leu Leu Leu Leu Met
1 5 10 15

Ala Ala Val Val Arg Cys Gln Glu Gln Ala Gln Thr Thr Asp Trp Arg
20 25 30

Ala Thr Leu Lys Thr Ile Arg Asn Gly Val His Lys Ile Asp Thr Tyr
35 40 45

Leu Asn Ala Ala Leu Asp Leu Leu Gly Gly Glu Asp Gly Leu Cys Gln
50 55 60

Tyr Lys Cys Ser Asp Gly Ser Lys Pro Phe Pro Arg Tyr Gly Tyr Lys
65 70 75 80

Pro Ser Pro Pro Asn Gly Cys Gly Ser Pro Leu Phe Gly Xaa His Leu
85 90 95

Asn Ile Gly Ile Pro Ser Leu Thr Lys Cys Cys Asn Gln His Asp Arg
100 105 110

Cys Tyr Glu Thr Cys Gly Lys Ser Lys Asn Asp Cys Asp Glu Glu Phe
115 120 125

Gln Tyr Cys Leu Ser Lys Ile Cys Arg Asp Val Gln Lys Thr Leu Gly
130 135 140

Leu Thr Gln His Val Gln Ala Cys Glu Thr Thr Val Glu Leu Leu Phe
145 150 155 160

Asp Ser Val Ile His Leu Gly Cys Lys Pro Tyr Leu Asp Ser Gln Arg
165 170 175

Ala Ala Cys Arg Cys His Tyr Glu Glu Lys Thr Asp Leu
180 185

<210> 108

<211> 61

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (61)

<223> Xaa equals stop translation

<400> 108

Met Gly Asn Cys Gln Ala Gly His Asn Leu His Leu Cys Leu Ala His
1 5 10 15

His Pro Pro Leu Val Cys Ala Thr Leu Ile Leu Leu Leu Gly Leu
20 25 30

Ser Gly Leu Gly Leu Gly Ser Phe Leu Leu Thr His Arg Thr Gly Leu
35 40 45

70

Arg Thr Leu Thr Ser Pro Arg Thr Gly Ser Leu Phe Xaa
 50 55 60

<210> 109

<211> 128

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (47)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (90)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 109

Met Arg Leu Glu Ser Leu Cys His Leu Cys Leu Ala Cys Leu Phe Phe
 1 5 10 15

Arg Leu Pro Ala Thr Arg Thr Val Tyr Cys Met Asn Glu Ala Glu Ile
 20 25 30

Val Asp Val Ala Leu Gly Ile Leu Ile Glu Ser Arg Lys Gln Xaa Lys
 35 40 45

Ala Cys Glu Gln Pro Ala Leu Ala Gly Ala Asp Asn Pro Glu His Ser
 50 55 60

Pro Pro Cys Ser Val Ser Pro His Thr Ser Ser Gly Ser Ser Ser Glu
 65 70 75 80

Glu Glu Asp Ser Gly Lys Gln Ala Leu Xaa Pro Gly Leu Ser Pro Ser
 85 90 95

Gln Arg Pro Gly Gly Ser Ser Ser Ala Cys Ser Arg Ser Pro Glu Glu
 100 105 110

Glu Glu Glu Glu Asp Val Leu Lys Tyr Val Arg Glu Ile Phe Phe Ser
 115 120 125

<210> 110

<211> 69

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (50)

<223> Xaa equals any of the naturally occurring L-amino acids

71

<220>

<221> SITE

<222> (69)

<223> Xaa equals stop translation

<400> 110

Met Pro His Phe Leu Asp Trp Phe Val Pro Val Tyr Leu Val Ile Ser
 1 5 10 15

Val Leu Ile Leu Val Gly Phe Gly Ala Cys Ile Tyr Tyr Phe Glu Pro
 20 25 30

Gly Leu Gln Glu Ala His Lys Trp Arg Met Gln Arg Pro Leu Val Asp
 35 40 45

Arg Xaa Leu Arg Lys Thr Leu Met Val Arg Asp Asn Leu Ala Phe Gly
 50 55 60

Gly Pro Glu Val Xaa
 65

<210> 111

<211> 123

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (123)

<223> Xaa equals stop translation

<400> 111

Met Ile Gly Gly Ile Thr Cys Ile Leu Ser Leu Ile Cys Ala Leu Ala
 1 5 10 15

Leu Ala Tyr Leu Asp Gln Arg Ala Glu Arg Ile Leu His Lys Glu Gln
 20 25 30

Gly Lys Thr Gly Glu Val Ile Lys Leu Thr Asp Val Lys Asp Phe Ser
 35 40 45

Leu Pro Leu Trp Leu Ile Phe Ile Ile Cys Val Cys Tyr Tyr Val Ala
 50 55 60

Val Phe Pro Phe Ile Gly Leu Gly Lys Val Phe Phe Thr Glu Lys Phe
 65 70 75 80

Gly Phe Ser Ser Gln Ala Ala Ser Ala Ile Asn Ser Val Val Tyr Val
 85 90 95

Ile Ser Ala Pro Met Ser Pro Val Phe Gly Leu Leu Val Asp Lys Thr
 100 105 110

Gly Lys Asn Ile Ile Trp Val Leu Cys Ala Xaa
 115 120

```
<210> 112
<211> 83
<212> PRT
<213> Homo sapiens
```

```
<220>  
<221> SITE  
<222> (83)  
<223> Xaa equals stop translation
```

```
<400> 112
Met Glu Lys Gln Cys Cys Ser His Pro Val Ile Cys Ser Leu Ser Thr
   1               5              10             15

Met Tyr Thr Phe Leu Leu Gly Ala Ile Phe Ile Ala Leu Ser Ser Ser
          20                25              30

Arg Ile Leu Leu Val Lys Tyr Ser Ala Asn Glu Gly Lys Leu Arg Leu
      35                 40              45

Gly Ile Cys Met Glu His Phe His Leu Ile Thr His Leu Ser Leu Ala
    50                55              60

Phe Gly Ser Val Ile Tyr Asn Met Glu Ile Ile Met Pro Phe Ala Ser
    65                70              75             80

Cys Glu Xaa
```

```
<210> 113
<211> 345
<212> PRT
<213> Homo sapiens
```

```
<220> .
<221> SITE
<222> (53)
<223> Xaa equals any of the naturally occurring L-amino acids
```

```
<220>  
<221> SITE  
<222> (345)  
<223> Xaa equals stop translation
```

```

<400> 113
Met Asp Phe Leu Val Leu Phe Leu Phe Tyr Leu Ala Ser Val Leu Met
  1              5              10              15
Gly Leu Val Leu Ile Cys Val Cys Ser Lys Thr His Ser Leu Lys Gly
      20              25              30
Leu Ala Arg Gly Gly Ala Gln Ile Phe Ser Cys Ile Ile Pro Glu Cys
      35              40              45
Leu Gln Arg Ala Xaa His Gly Leu Leu His Tyr Leu Phe His Thr Arg
      50              55              60

```

73

Asn His Thr Phe Ile Val Leu His Leu Val Leu Gln Gly Met Val Tyr
 65 70 75 80
 Thr Glu Tyr Thr Trp Glu Val Phe Gly Tyr Cys Gln Glu Leu Glu Leu
 85 90 95
 Ser Leu His Tyr Leu Leu Leu Pro Tyr Leu Leu Leu Gly Val Asn Leu
 100 105 110
 Phe Phe Phe Thr Leu Thr Cys Gly Thr Asn Pro Gly Ile Ile Thr Lys
 115 120 125
 Ala Asn Glu Leu Leu Phe Leu His Val Tyr Glu Phe Asp Glu Val Met
 130 135 140
 Phe Pro Lys Asn Val Arg Cys Ser Thr Cys Asp Leu Arg Lys Pro Ala
 145 150 155 160
 Arg Ser Lys His Cys Ser Val Cys Asn Trp Cys Val His Arg Phe Asp
 165 170 175
 His His Cys Val Trp Val Asn Asn Cys Ile Gly Ala Trp Asn Ile Arg
 180 185 190
 Tyr Phe Leu Ile Tyr Val Leu Thr Leu Thr Ala Ser Ala Ala Thr Val
 195 200 205
 Ala Ile Val Ser Thr Thr Phe Leu Val His Leu Val Val Met Ser Asp
 210 215 220
 Leu Tyr Gln Glu Thr Tyr Ile Asp Asp Leu Gly His Leu His Val Met
 225 230 235 240
 Asp Thr Val Phe Leu Ile Gln Tyr Leu Phe Leu Thr Phe Pro Arg Ile
 245 250 255
 Val Phe Met Leu Gly Phe Val Val Val Leu Ser Phe Leu Leu Gly Gly
 260 265 270
 Tyr Leu Leu Phe Val Leu Tyr Leu Ala Ala Thr Asn Gln Thr Thr Asn
 275 280 285
 Glu Trp Tyr Arg Gly Asp Trp Ala Trp Cys Gln Arg Cys Pro Leu Val
 290 295 300
 Ala Trp Pro Pro Ser Ala Glu Pro Gln Val His Arg Asn Ile His Ser
 305 310 315 320
 His Gly Leu Arg Ser Asn Leu Gln Glu Ile Phe Leu Pro Ala Phe Pro
 325 330 335
 Cys His Glu Arg Lys Lys Gln Glu Xaa
 340 345

<210> 114

<211> 181

<212> PRT

74

<213> Homo sapiens

<220>

<221> SITE

<222> (110)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 114

Met Ala Asp Pro His Val Ser Phe Leu Ser Phe Arg Gln Leu Phe Ser
1 5 10 15

Trp Ala Ala Val Ile Leu Leu Arg Gly Ile Leu Gly Thr Val Ala Pro
20 25 30

Pro Pro Cys Pro Cys Val Leu Asp Leu Ala Val Tyr Pro Leu His Leu
35 40 45

Pro Val Glu Ala Pro Cys Leu Glu Val Val Phe Lys Gln Lys Asn Gly
50 55 60

Lys Asp Asn Cys Leu Val Phe Tyr Pro Asp Pro Ile Pro Leu Arg Gly
65 70 75 80

Ser Leu Leu Gly Pro Phe Ile Lys Asn Gln Cys His Ser Ser Val Ile
85 90 95

Pro Leu Ser Asp Ser Ala Thr Ser Lys Ala Arg Ala Leu Xaa Leu Pro
100 105 110

Gly Arg Glu Thr Val Leu Ser Val Leu Pro Val Phe Ser Ser Pro Thr
115 120 125

Leu Pro Arg Thr His Ala Leu Gly Asp Ser Leu Gly Val Pro Gly Leu
130 135 140

Leu Val Cys Ser Glu Thr Ser Thr Leu Asn Asp His Trp Cys Cys Arg
145 150 155 160

Arg Ala Gly Ala Tyr Ile Pro Ile Asn Arg Arg Phe Ser His Leu Met
165 170 175

Pro Leu Ala Phe Ser
180

<210> 115

<211> 116

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (116)

<223> Xaa equals stop translation

<400> 115

Met Pro Ser Ser Ser Ser Gly Leu Gly Ser Pro Ser Arg Pro Pro Ser
1 5 10 15

75

Ser Phe Leu Cys Leu Leu Leu Leu Leu Leu Pro Pro Ala Ala Leu Ala
 20 25 30
 Leu Leu Leu Phe Phe Leu Asp Phe Phe Pro Pro Arg Ala Ala Val Ser
 35 40 45
 Pro Phe Leu Pro Asp His Cys Ser Ala Arg Gln Pro Arg Val Trp Arg
 50 55 60
 Arg Glu Thr Leu Asn Arg Ser Ala Ser Gly Leu Gly Cys Trp Ala Arg
 65 70 75 80
 Ser Thr Glu Gln Gly Ala Val Gly Val Ala Thr Gly Thr Val Leu Asp
 85 90 95
 Ile Ser Leu Pro Ala Ser Cys Leu Ser Leu Trp Pro Pro Gly Pro Ser
 100 105 110
 Gly Gly Ile Xaa
 115

<210> 116
 <211> 71
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (71)
 <223> Xaa equals stop translation

<400> 116
 Met Asn Leu Leu Gly Met Ile Phe Ser Met Cys Gly Leu Met Leu Lys
 1 5 10 15
 Leu Lys Trp Cys Ala Trp Val Ala Val Tyr Cys Ser Phe Ile Ser Phe
 20 25 30
 Ala Asn Ser Arg Ser Ser Glu Asp Thr Lys Gln Met Met Ser Ser Phe
 35 40 45
 Met Leu Ser Ile Ser Ala Val Val Met Ser Tyr Leu Gln Asn Pro Gln
 50 55 60
 Pro Met Thr Pro Pro Trp Xaa
 65 70

<210> 117
 <211> 64
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (64)

76

<223> Xaa equals stop translation

<400> 117

Met Arg Asp Leu Ser Phe Leu Tyr Thr Leu Leu Trp Leu Pro Glu Ile
 1 5 10 15

Trp Gln Ala Leu Ala Gly Gly Ile Arg Leu Asp Glu Val Glu Leu Leu
 20 25 30

Glu Asn Glu Ala Val Leu Gly Glu Glu Met Arg Leu Tyr Arg Lys Ile
 35 40 45

Asn Glu Val Val Leu Ser Gly Asn Glu Val Val Leu Gly Gly Lys Xaa
 50 55 60

<210> 118

<211> 335

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (335)

<223> Xaa equals stop translation

<400> 118

Met Gly Ile Phe Pro Gly Ile Ile Leu Ile Phe Leu Arg Val Lys Phe
 1 5 10 15

Ala Thr Ala Ala Val Ile Val Ser Gly Val Ser Lys His Leu His Cys
 20 25 30

Ile Ser His Gln Lys Ser Thr Thr Val Ser His Glu Met Ser Gly Leu
 35 40 45

Asn Trp Lys Pro Phe Val Tyr Gly Gly Leu Ala Ser Ile Val Ala Glu
 50 55 60

Phe Gly Thr Phe Pro Val Asp Leu Thr Lys Thr Arg Leu Gln Val Gln
 65 70 75 80

Gly Gln Ser Ile Asp Ala Arg Phe Lys Glu Ile Lys Tyr Arg Gly Met
 85 90 95

Phe His Ala Leu Phe Arg Ile Cys Lys Glu Glu Gly Val Leu Ala Leu
 100 105 110

Tyr Ser Gly Ile Ala Pro Ala Leu Leu Arg Gln Ala Ser Tyr Gly Thr
 115 120 125

Ile Lys Ile Gly Ile Tyr Gln Ser Leu Lys Arg Leu Phe Val Glu Arg
 130 135 140

Leu Glu Asp Glu Thr Leu Leu Ile Asn Met Ile Cys Gly Val Val Ser

145					150					155					160
Gly	Val	Ile	Ser	Ser	Thr	Ile	Ala	Asn	Pro	Thr	Asp	Val	Leu	Lys	Ile
				165					170					175	
Arg	Met	Gln	Ala	Gln	Gly	Ser	Leu	Phe	Gln	Gly	Ser	Met	Ile	Gly	Ser
			180					185					190		
Phe	Ile	Asp	Ile	Tyr	Gln	Gln	Glu	Gly	Thr	Arg	Gly	Leu	Trp	Arg	Gly
		195					200					205			
Val	Val	Pro	Thr	Ala	Gln	Arg	Ala	Ala	Ile	Val	Val	Gly	Val	Glu	Leu
	210					215					220				
Pro	Val	Tyr	Asp	Ile	Thr	Lys	Lys	His	Leu	Ile	Leu	Ser	Gly	Met	Met
225					230					235					240
Gly	Asp	Thr	Ile	Leu	Thr	His	Phe	Val	Ser	Ser	Phe	Thr	Cys	Gly	Leu
				245					250					255	
Ala	Gly	Ala	Leu	Ala	Ser	Asn	Pro	Val	Asp	Val	Val	Arg	Thr	Arg	Met
			260					265					270		
Met	Asn	Gln	Arg	Ala	Ile	Val	Gly	His	Val	Asp	Leu	Tyr	Lys	Gly	Thr
		275					280					285			
Val	Asp	Gly	Ile	Leu	Lys	Met	Trp	Lys	His	Glu	Gly	Phe	Phe	Ala	Leu
	290					295					300				
Tyr	Lys	Gly	Phe	Trp	Pro	Asn	Trp	Leu	Arg	Leu	Gly	Pro	Trp	Asn	Ile
305					310					315					320
Ile	Phe	Phe	Ile	Thr	Tyr	Glu	Gln	Leu	Lys	Arg	Leu	Gln	Ile	Xaa	
				325					330					335	

```
<210> 119
<211> 221
<212> PRT
<213> Homo sapiens
```

```
<220>
<221> SITE
<222> (51)
<223> Xaa equals any of the naturally occurring L-amino acids
```

```

<400> 119
Met Ala Leu Ala Leu Ala Ala Leu Ala Ala Val Glu Pro Ala Cys Gly
  1                      5                      10                      15

Ser Arg Tyr Gln Gln Leu Gln Asn Glu Glu Glu Ser Gly Glu Pro Glu
                      20                      25                      30

Gln Ala Ala Gly Asp Ala Pro Pro Pro Tyr Ser Ser Ile Ser Ala Glu
          35                      40                      45

Ser Ala Xaa Tyr Phe Asp Tyr Lys Asp Glu Ser Gly Phe Pro Lys Pro
    50                      55                      60

```

Pro Ser Tyr Asn Val Ala Thr Thr Leu Pro Ser Tyr Asp Glu Ala Glu
 65 70 75 80
 Arg Thr Lys Ala Glu Ala Thr Ile Pro Leu Val Pro Gly Arg Asp Glu
 85 90 95
 Asp Phe Val Gly Arg Asp Asp Phe Asp Asp Ala Asp Gln Leu Arg Ile
 100 105 110
 Gly Asn Asp Gly Ile Phe Met Leu Thr Phe Phe Met Ala Phe Leu Phe
 115 120 125
 Asn Trp Ile Gly Phe Phe Leu Ser Phe Cys Leu Thr Thr Ser Ala Ala
 130 135 140
 Gly Arg Tyr Gly Ala Ile Ser Gly Phe Gly Leu Ser Leu Ile Lys Trp
 145 150 155 160
 Ile Leu Ile Val Arg Phe Ser Thr Tyr Phe Pro Gly Tyr Phe Asp Gly
 165 170 175
 Gln Tyr Trp Leu Trp Trp Val Phe Leu Val Leu Gly Phe Leu Leu Phe
 180 185 190
 Leu Arg Gly Phe Ile Asn Tyr Ala Lys Val Arg Lys Met Pro Glu Thr
 195 200 205
 Phe Ser Asn Leu Pro Arg Thr Arg Val Leu Phe Ile Tyr
 210 215 220

<210> 120

<211> 473

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (473)

<223> Xaa equals stop translation

<400> 120

Met Lys Phe Leu Ile Phe Ala Phe Phe Gly Gly Val His Leu Leu Ser
 1 5 10 15
 Leu Cys Ser Gly Lys Ala Ile Cys Lys Asn Gly Ile Ser Lys Arg Thr
 20 25 30
 Phe Glu Glu Ile Lys Glu Glu Ile Ala Ser Cys Gly Asp Val Ala Lys
 35 40 45
 Ala Ile Ile Asn Leu Ala Val Tyr Gly Lys Ala Gln Asn Arg Ser Tyr
 50 55 60
 Glu Arg Leu Ala Leu Leu Val Asp Thr Val Gly Pro Arg Leu Ser Gly
 65 70 75 80

79

Ser Lys Asn Leu Glu Lys Ala Ile Gln Ile Met Tyr Gln Asn Leu Gln
 85 90 95
 Gln Asp Gly Leu Glu Lys Val His Leu Glu Pro Val Arg Ile Pro His
 100 105 110
 Trp Glu Arg Gly Glu Glu Ser Ala Val Met Leu Glu Pro Arg Ile His
 115 120 125
 Lys Ile Ala Ile Leu Gly Leu Gly Ser Ser Ile Gly Thr Pro Pro Glu
 130 135 140
 Gly Ile Thr Ala Glu Val Leu Val Val Thr Ser Phe Asp Glu Leu Gln
 145 150 155 160
 Arg Arg Ala Ser Glu Ala Arg Gly Lys Ile Val Val Tyr Asn Gln Pro
 165 170 175
 Tyr Ile Asn Tyr Ser Arg Thr Val Gln Tyr Arg Thr Gln Gly Ala Val
 180 185 190
 Glu Ala Ala Lys Val Gly Ala Leu Ala Ser Leu Ile Arg Ser Val Ala
 195 200 205
 Ser Phe Ser Ile Tyr Ser Pro His Thr Gly Ile Gln Glu Tyr Gln Asp
 210 215 220
 Gly Val Pro Lys Ile Pro Thr Ala Cys Ile Thr Val Glu Asp Ala Glu
 225 230 235 240
 Met Met Ser Arg Met Ala Ser His Gly Ile Lys Ile Val Ile Gln Leu
 245 250 255
 Lys Met Gly Ala Lys Thr Tyr Pro Asp Thr Asp Ser Phe Asn Thr Val
 260 265 270
 Ala Glu Ile Thr Gly Ser Lys Tyr Pro Glu Gln Val Val Leu Val Ser
 275 280 285
 Gly His Leu Asp Ser Trp Asp Val Gly Gln Gly Ala Met Asp Asp Gly
 290 295 300
 Gly Gly Ala Phe Ile Ser Trp Glu Ala Leu Ser Leu Ile Lys Asp Leu
 305 310 315 320
 Gly Leu Arg Pro Lys Arg Thr Leu Arg Leu Val Leu Trp Thr Ala Glu
 325 330 335
 Glu Gln Gly Gly Val Gly Ala Phe Gln Tyr Tyr Gln Leu His Lys Val
 340 345 350
 Asn Ile Ser Asn Tyr Ser Leu Val Met Glu Ser Asp Ala Gly Thr Phe
 355 360 365
 Leu Pro Thr Gly Leu Gln Phe Thr Gly Ser Glu Lys Ala Arg Ala Ile
 370 375 380
 Met Glu Glu Val Met Ser Leu Leu Gln Pro Leu Asn Ile Thr Gln Val

[illegible]

Asp Arg His Val Val Leu Ser Xaa
165

<210> 122
<211> 47
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (47)
<223> Xaa equals stop translation

<400> 122
Met Lys Phe Ile Leu Trp Arg Arg Phe Arg Trp Ala Ile Ile Leu Phe
1 5 10 15
Ile Ile Leu Phe Ile Leu Leu Leu Phe Leu Ala Ile Phe Ile Tyr Ala
20 25 30
Phe Pro Asn Tyr Ala Ala Met Lys Leu Val Lys Pro Phe Ser Xaa
35 40 45

<210> 123
<211> 108
<212> PRT
<213> Homo sapiens

<400> 123
Met His Gln Asp Trp Leu Cys Asn Leu Gly Trp Pro Leu Leu Ser Leu
1 5 10 15
Trp Ala Ala Glu Ser Ala Pro His Val Ala Met Ala Ser Ala Thr Ala
20 25 30
Gln Leu Trp Ser Arg Pro Cys Gly Arg Thr His Met Val Ser Leu Ala
35 40 45
Leu Gly His Gln Glu Thr Gly Leu Trp Leu Cys Ser Ala Phe Gly Cys
50 55 60
Val Val Asp Ser Pro Trp Ala Ser Val Cys Pro Ser Val Lys Gly Gln
65 70 75 80
Leu Thr Val Cys Gly Ile Leu Pro Arg Val Pro Val Cys Val Tyr Val
85 90 95
Cys Ala Cys Val Arg Val Ser Met Cys Val His Ile
100 105

<210> 124
<211> 60
<212> PRT
<213> Homo sapiens

<400> 124

Met Arg Gly Cys Val Pro Ala Phe Leu Leu His Val Leu Ser Leu Arg
 1 5 10 15
 Arg Ala Cys Cys Thr Gln Ala Ala Gln Val Phe Thr Ala Gln Leu Pro
 20 25 30
 Gly Arg Gln Val Ala Arg Arg Arg Gly Gly Trp His Glu Gln Gln Gly
 35 40 45
 Gly Pro Met Leu Cys Ser Ser His His Ser Arg Thr
 50 55 60

<210> 125

<211> 248

<212> PRT

<213> Homo sapiens

<400> 125

Met Ala Met Leu Pro Leu Val Leu His Trp Phe Phe Ile Glu Trp Tyr
 1 5 10 15
 Ser Gly Lys Lys Ser Ser Ser Ala Leu Phe Gln His Ile Thr Ala Leu
 20 25 30
 Phe Glu Cys Ser Met Ala Ala Ile Ile Thr Leu Leu Val Ser Asp Pro
 35 40 45
 Val Gly Val Leu Tyr Ile Arg Ser Cys Arg Val Leu Met Leu Ser Asp
 50 55 60
 Trp Tyr Thr Met Leu Tyr Asn Pro Ser Pro Asp Tyr Val Thr Thr Val
 65 70 75 80
 His Cys Thr His Glu Ala Val Tyr Pro Leu Tyr Thr Ile Val Phe Ile
 85 90 95
 Tyr Tyr Ala Phe Cys Leu Val Leu Met Met Leu Leu Arg Pro Leu Leu
 100 105 110
 Val Lys Lys Ile Ala Cys Gly Leu Gly Lys Ser Asp Arg Phe Lys Ser
 115 120 125
 Ile Tyr Ala Ala Leu Tyr Phe Phe Pro Ile Leu Thr Val Leu Gln Ala
 130 135 140
 Val Gly Gly Gly Leu Leu Tyr Tyr Ala Phe Pro Tyr Ile Ile Leu Val
 145 150 155 160
 Leu Ser Leu Val Thr Leu Ala Val Tyr Met Ser Ala Ser Glu Ile Glu
 165 170 175
 Asn Cys Tyr Asp Leu Leu Val Arg Lys Lys Arg Leu Ile Val Leu Phe
 180 185 190
 Ser His Trp Leu Leu His Ala Tyr Gly Ile Ile Ser Ile Ser Arg Val

83

195 200 205
 Asp Lys Leu Glu Gln Asp Leu Pro Pro Leu Ala Leu Val Pro Thr Pro
 210 215 220
 Ala Leu Phe Tyr Leu Phe Thr Ala Lys Phe Thr Glu Pro Ser Arg Ile
 225 230 235 240
 Leu Ser Glu Gly Ala Asn Gly His
 245

 <210> 126
 <211> 248
 <212> PRT
 <213> Homo sapiens

 <400> 126
 Met Glu Lys Ile Pro Glu Ile Gly Lys Phe Gly Glu Lys Ala Pro Pro
 1 5 10 15
 Ala Pro Ser His Val Trp Arg Pro Ala Ala Leu Phe Leu Thr Leu Leu
 20 25 30
 Cys Leu Leu Leu Leu Ile Gly Leu Gly Val Leu Ala Ser Met Phe His
 35 40 45
 Val Thr Leu Lys Ile Glu Met Lys Lys Met Asn Lys Leu Gln Asn Ile
 50 55 60
 Ser Glu Glu Leu Gln Arg Asn Ile Ser Leu Gln Leu Met Ser Asn Met
 65 70 75 80
 Asn Ile Ser Asn Lys Ile Arg Asn Leu Ser Thr Thr Leu Gln Thr Ile
 85 90 95
 Ala Thr Lys Leu Cys Arg Glu Leu Tyr Ser Lys Glu Gln Glu His Lys
 100 105 110
 Cys Lys Pro Cys Pro Arg Arg Trp Ile Trp His Lys Asp Ser Cys Tyr
 115 120 125
 Phe Leu Ser Asp Asp Val Gln Thr Trp Gln Glu Ser Lys Met Ala Cys
 130 135 140
 Ala Ala Gln Asn Ala Ser Leu Leu Lys Ile Asn Asn Lys Asn Ala Leu
 145 150 155 160
 Glu Phe Ile Lys Ser Gln Ser Arg Ser Tyr Asp Tyr Trp Leu Gly Leu
 165 170 175
 Ser Pro Glu Glu Asp Ser Thr Arg Gly Met Arg Val Asp Asn Ile Ile
 180 185 190
 Asn Ser Ser Ala Trp Val Ile Arg Asn Ala Pro Asp Leu Asn Asn Met
 195 200 205
 Tyr Cys Gly Tyr Ile Asn Arg Leu Tyr Val Gln Tyr Tyr His Cys Thr

84

210

215

220

Tyr Lys Gln Arg Met Ile Cys Glu Lys Met Ala Asn Pro Val Gln Leu
 225 230 235 240

Gly Ser Thr Tyr Phe Arg Glu Ala
 245

<210> 127
 <211> 612
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (245)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (246)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (249)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 127
 Met Ala Ala Ala Gly Arg Leu Pro Ser Ser Trp Ala Leu Phe Ser Pro
 1 5 10 15

Leu Leu Ala Gly Leu Ala Leu Leu Gly Val Gly Pro Val Pro Ala Arg
 20 25 30

Ala Leu His Asn Val Thr Ala Glu Leu Phe Gly Ala Glu Ala Trp Gly
 35 40 45

Thr Leu Ala Ala Phe Gly Asp Leu Asn Ser Asp Lys Gln Thr Asp Leu
 50 55 60

Phe Val Leu Arg Glu Arg Asn Asp Leu Ile Val Phe Leu Ala Asp Gln
 65 70 75 80

Asn Ala Pro Tyr Phe Lys Pro Lys Val Lys Val Ser Phe Lys Asn His
 85 90 95

Ser Ala Leu Ile Thr Ser Val Val Pro Gly Asp Tyr Asp Gly Asp Ser
 100 105 110

Gln Met Asp Val Leu Leu Thr Tyr Leu Pro Lys Asn Tyr Ala Lys Ser
 115 120 125

Glu Leu Gly Ala Val Ile Phe Trp Gly Gln Asn Gln Thr Leu Asp Pro
 130 135 140

Asn Asn Met Thr Ile Leu Asn Arg Thr Phe Gln Asp Glu Pro Leu Ile

85

145		150		155		160
Met Asp Phe Asn Gly Asp Leu Ile Pro Asp Ile Phe Gly Ile Thr Asn						
	165			170		175
Glu Ser Asn Gln Pro Gln Ile Leu Leu Gly Gly Asn Leu Ser Trp His						
	180		185			190
Pro Ala Leu Thr Thr Thr Ser Lys Met Arg Ile Pro His Ser His Ala						
	195		200			205
Phe Ile Asp Leu Thr Glu Asp Phe Thr Ala Asp Leu Phe Leu Thr Thr						
	210		215			220
Leu Asn Ala Thr Thr Ser Thr Phe Gln Phe Glu Ile Trp Glu Asn Leu						
	225		230		235	240
Asp Gly Asn Phe Xaa Xaa Ser Thr Xaa Leu Glu Lys Pro Gln Asn Met						
	245			250		255
Met Val Val Gly Gln Ser Ala Phe Ala Asp Phe Asp Gly Asp Gly His						
	260			265		270
Met Asp His Leu Leu Pro Gly Cys Glu Asp Lys Asn Cys Gln Lys Ser						
	275		280			285
Thr Ile Tyr Leu Val Arg Ser Gly Met Lys Gln Trp Val Pro Val Leu						
	290		295			300
Gln Asp Phe Ser Asn Lys Gly Thr Leu Trp Gly Phe Val Pro Phe Val						
	305		310		315	320
Asp Glu Gln Gln Pro Thr Glu Ile Pro Ile Pro Ile Thr Leu His Ile						
		325		330		335
Gly Asp Tyr Asn Met Asp Gly Tyr Pro Asp Ala Leu Val Ile Leu Lys						
	340			345		350
Asn Thr Ser Gly Ser Asn Gln Gln Ala Phe Leu Leu Glu Asn Val Pro						
	355			360		365
Cys Asn Asn Ala Ser Cys Glu Glu Ala Arg Arg Met Phe Lys Val Tyr						
	370		375			380
Trp Glu Leu Thr Asp Leu Asn Gln Ile Lys Asp Ala Met Val Ala Thr						
	385		390		395	400
Phe Phe Asp Ile Tyr Glu Asp Gly Ile Leu Asp Ile Val Val Leu Ser						
		405		410		415
Lys Gly Tyr Thr Lys Asn Asp Phe Ala Ile His Thr Leu Lys Asn Asn						
	420			425		430
Phe Glu Ala Asp Ala Tyr Phe Val Lys Val Ile Val Leu Ser Gly Leu						
	435			440		445
Cys Ser Asn Asp Cys Pro Arg Lys Ile Thr Pro Phe Gly Val Asn Gln						
	450		455			460

Pro Gly Pro Tyr Ile Met Tyr Thr Thr Val Asp Ala Asn Gly Tyr Leu
 465 470 475 480
 Lys Asn Gly Ser Ala Gly Gln Leu Ser Gln Ser Ala His Leu Ala Leu
 485 490 495
 Gln Leu Pro Tyr Asn Val Leu Gly Leu Gly Arg Ser Ala Asn Phe Leu
 500 505 510
 Asp His Leu Tyr Val Gly Ile Pro Arg Pro Ser Gly Glu Lys Ser Ile
 515 520 525
 Arg Lys Gln Glu Trp Thr Ala Ile Ile Pro Asn Ser Gln Leu Ile Val
 530 535 540
 Ile Pro Tyr Pro His Asn Val Pro Arg Ser Trp Ser Ala Lys Leu Tyr
 545 550 555 560
 Leu Thr Pro Ser Asn Ile Val Leu Leu Thr Ala Ile Ala Leu Ile Gly
 565 570 575
 Val Cys Val Phe Ile Leu Ala Ile Ile Gly Ile Leu His Trp Gln Glu
 580 585 590
 Lys Lys Ala Asp Asp Arg Glu Lys Arg Gln Glu Ala His Arg Phe His
 595 600 605
 Phe Asp Ala Met
 610

<210> 128

<211> 447

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (8)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (28)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (309)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (333)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 128

87

Met Glu Leu Ser Gln Met Ser Xaa Leu Met Gly Leu Ser Val Leu Leu
 1 5 10 15
 Gly Leu Leu Ala Leu Met Ala Thr Ala Ala Val Xaa Arg Gly Trp Leu
 20 25 30
 Arg Ala Gly Glu Glu Arg Ser Gly Arg Pro Ala Cys Gln Lys Ala Asn
 35 40 45
 Gly Phe Pro Pro Asp Lys Ser Ser Gly Ser Lys Lys Gln Lys Gln Tyr
 50 55 60
 Gln Arg Ile Arg Lys Glu Lys Pro Gln Gln His Asn Phe Thr His Arg
 65 70 75 80
 Leu Leu Ala Ala Ala Leu Lys Ser His Ser Gly Asn Ile Ser Cys Met
 85 90 95
 Asp Phe Ser Ser Asn Gly Lys Tyr Leu Ala Thr Cys Ala Asp Asp Arg
 100 105 110
 Thr Ile Arg Ile Trp Ser Thr Lys Asp Phe Leu Gln Arg Glu His Arg
 115 120 125
 Ser Met Arg Ala Asn Val Glu Leu Asp His Ala Thr Leu Val Arg Phe
 130 135 140
 Ser Pro Asp Cys Arg Ala Phe Ile Val Trp Leu Ala Asn Gly Asp Thr
 145 150 155 160
 Leu Arg Val Phe Lys Met Thr Lys Arg Glu Asp Gly Gly Tyr Thr Phe
 165 170 175
 Thr Ala Thr Pro Glu Asp Phe Pro Lys Lys His Lys Ala Pro Val Ile
 180 185 190
 Asp Ile Gly Ile Ala Asn Thr Gly Lys Phe Ile Met Thr Ala Ser Ser
 195 200 205
 Asp Thr Thr Val Leu Ile Trp Ser Leu Lys Gly Gln Val Leu Ser Thr
 210 215 220
 Ile Asn Thr Asn Gln Met Asn Asn Thr His Ala Ala Val Ser Pro Cys
 225 230 235 240
 Gly Arg Phe Val Ala Ser Cys Gly Phe Thr Pro Asp Val Lys Val Trp
 245 250 255
 Glu Val Cys Phe Gly Lys Lys Gly Glu Phe Gln Glu Val Val Arg Ala
 260 265 270
 Phe Glu Leu Lys Gly His Ser Ala Ala Val His Ser Phe Ala Phe Ser
 275 280 285
 Asn Asp Ser Arg Arg Met Ala Ser Val Ser Lys Asp Gly Thr Trp Lys
 290 295 300
 Leu Trp Asp Thr Xaa Val Glu Tyr Lys Lys Lys Gln Asp Pro Tyr Leu

305						310						315					320
Leu	Lys	Thr	Gly	Arg	Phe	Glu	Glu	Ala	Ala	Gly	Ala	Xaa	Pro	Cys	Arg		
				325					330					335			
Leu	Ala	Leu	Ser	Pro	Asn	Ala	Gln	Val	Leu	Ala	Leu	Ala	Ser	Gly	Ser		
			340					345					350				
Ser	Ile	His	Leu	Tyr	Asn	Thr	Arg	Arg	Gly	Glu	Lys	Glu	Glu	Cys	Phe		
		355					360					365					
Glu	Arg	Val	His	Gly	Glu	Cys	Ile	Ala	Asn	Leu	Ser	Phe	Asp	Ile	Thr		
	370					375					380						
Gly	Arg	Phe	Leu	Ala	Ser	Cys	Gly	Asp	Arg	Ala	Val	Arg	Leu	Phe	His		
385					390					395					400		
Asn	Thr	Pro	Gly	His	Arg	Ala	Met	Val	Glu	Glu	Met	Gln	Gly	His	Leu		
				405					410					415			
Lys	Arg	Ala	Ser	Asn	Glu	Ser	Thr	Arg	Gln	Arg	Leu	Gln	Gln	Gln	Leu		
			420					425					430				
Thr	Gln	Ala	Gln	Glu	Thr	Leu	Lys	Ser	Leu	Gly	Ala	Leu	Lys	Lys			
	435						440					445					

```
<210> 129
<211> 291
<212> PRT
<213> Homo sapiens
```

```

<400> 129
Met Leu Phe Leu Phe Ser Met Ala Thr Leu Leu Arg Thr Ser Phe Ser
  1                      5                      10                      15

Asp Pro Gly Val Ile Pro Arg Ala Leu Pro Asp Glu Ala Ala Phe Ile
      20                      25                      30

Glu Met Glu Ile Glu Ala Thr Asn Gly Ala Val Pro Gln Gly Gln Arg
      35                      40                      45

Pro Pro Pro Arg Ile Lys Asn Phe Gln Ile Asn Asn Gln Ile Val Lys
      50                      55                      60

Leu Lys Tyr Cys Tyr Thr Cys Lys Ile Phe Arg Pro Pro Arg Ala Ser
      65                      70                      75                      80

His Cys Ser Ile Cys Asp Asn Cys Val Glu Arg Phe Asp His His Cys
      85                      90                      95

Pro Trp Val Gly Asn Cys Val Gly Lys Arg Asn Tyr Arg Tyr Phe Tyr
      100                      105                      110

Leu Phe Ile Leu Ser Leu Ser Leu Leu Thr Ile Tyr Val Phe Ala Phe
      115                      120                      125

Asn Ile Val Tyr Val Ala Leu Lys Ser Leu Lys Ile Gly Phe Leu Glu

```

89

130 135 140
 Thr Leu Lys Glu Thr Pro Gly Thr Val Leu Glu Val Leu Ile Cys Phe
 145 150 155 160
 Phe Thr Leu Trp Ser Val Val Gly Leu Thr Gly Phe His Thr Phe Leu
 165 170 175
 Val Ala Leu Asn Gln Thr Thr Asn Glu Asp Ile Lys Gly Ser Trp Thr
 180 185 190
 Gly Lys Asn Arg Val Gln Asn Pro Tyr Ser His Gly Asn Ile Val Lys
 195 200 205
 Asn Cys Cys Glu Val Leu Cys Gly Pro Leu Pro Pro Ser Val Leu Asp
 210 215 220
 Arg Arg Gly Ile Leu Pro Leu Glu Glu Ser Gly Ser Arg Pro Pro Ser
 225 230 235 240
 Thr Gln Glu Thr Ser Ser Ser Leu Leu Pro Gln Ser Pro Ala Pro Thr
 245 250 255
 Glu His Leu Asn Ser Asn Glu Met Pro Glu Asp Ser Ser Thr Pro Glu
 260 265 270
 Glu Met Pro Pro Pro Glu Pro Pro Glu Pro Pro Gln Glu Ala Ala Glu
 275 280 285
 Ala Glu Lys
 290

<210> 130
 <211> 78
 <212> PRT
 <213> Homo sapiens

<400> 130
 Met Val Arg Lys Trp Leu Thr Phe Val Glu His Leu Leu Cys Ala Trp
 1 5 10 15
 Pro Arg Leu Gly Ala Phe Val Pro Arg Val Thr Pro Ser Glu Cys Ser
 20 25 30
 Ser Leu Pro His Ser Asn Trp Gly Val Gly Gly Arg Ala Ala Gln Leu
 35 40 45
 Thr Gly Ala Glu Leu Lys Thr His Ser Trp Val Cys Leu Gly Trp Ala
 50 55 60
 Val Leu Val Ala Pro Val Ala Asn Thr Arg Ala Pro Phe Thr
 65 70 75

<210> 131
 <211> 333
 <212> PRT

90

<213> Homo sapiens

<220>

<221> SITE

<222> (97)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 131

```

Met Leu Met Phe Ala Val Ile Val Ala Ser Ser Gly Leu Leu Leu Met
 1              5              10              15

Ile Glu Arg Gly Ile Leu Ala Glu Met Lys Pro Leu Pro Leu His Pro
      20              25              30

Pro Gly Arg Glu Gly Thr Ala Trp Arg Gly Lys Ala Pro Lys Pro Gly
      35              40              45

Gly Leu Ser Leu Arg Ala Gly Asp Ala Asp Leu Gln Val Arg Gln Asp
 50              55              60

Val Arg Asn Arg Thr Leu Arg Ala Val Cys Gly Gln Pro Gly Met Pro
 65              70              75              80

Arg Asp Pro Trp Asp Leu Pro Val Gly Gln Arg Arg Thr Leu Leu Arg
      85              90              95

Xaa Ile Leu Val Ser Asp Arg Tyr Arg Phe Leu Tyr Cys Tyr Val Pro
      100              105              110

Lys Val Ala Cys Ser Asn Trp Lys Arg Val Met Lys Val Leu Ala Gly
      115              120              125

Val Leu Asp Ser Val Asp Val Arg Leu Lys Met Asp His Arg Ser Asp
      130              135              140

Leu Val Phe Leu Ala Asp Leu Arg Pro Glu Glu Ile Arg Tyr Arg Leu
      145              150              155              160

Gln His Tyr Phe Lys Phe Leu Phe Val Arg Glu Pro Leu Glu Arg Leu
      165              170              175

Leu Ser Ala Tyr Arg Asn Lys Phe Gly Glu Ile Arg Glu Tyr Gln Gln
      180              185              190

Arg Tyr Gly Ala Glu Ile Val Arg Arg Tyr Arg Ala Gly Ala Gly Pro
      195              200              205

Ser Pro Ala Gly Asp Asp Val Thr Phe Pro Glu Phe Leu Arg Tyr Leu
      210              215              220

Val Asp Glu Asp Pro Glu Arg Met Asn Glu His Trp Met Pro Val Tyr
      225              230              235              240

His Leu Cys Gln Pro Cys Ala Val His Tyr Asp Phe Val Gly Ser Tyr
      245              250              255

Glu Arg Leu Glu Ala Asp Ala Asn Gln Val Leu Glu Trp Val Arg Ala
      260              265              270

```

91

Pro Pro His Val Arg Phe Pro Ala Arg Gln Ala Trp Tyr Arg Pro Ala
 275 280 285

Ser Pro Glu Ser Leu His Tyr His Leu Cys Ser Ala Pro Arg Ala Leu
 290 295 300

Leu Gln Asp Val Leu Pro Lys Tyr Ile Leu Asp Phe Ser Leu Phe Ala
 305 310 315 320

Tyr Pro Leu Pro Asn Val Thr Lys Glu Ala Cys Gln Gln
 325 330

<210> 132

<211> 164

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (126)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 132

Met Leu Pro Leu Leu Ile Ile Cys Leu Leu Pro Ala Ile Glu Gly Lys
 1 5 10 15

Asn Cys Leu Arg Cys Trp Pro Glu Leu Ser Ala Leu Ile Asp Tyr Asp
 20 25 30

Leu Gln Ile Leu Trp Val Thr Pro Gly Pro Pro Thr Glu Leu Ser Gln
 35 40 45

Ser Ile His Ser Leu Phe Leu Glu Asp Asn Asn Phe Leu Lys Pro Trp
 50 55 60

Tyr Leu Asp Arg Asp His Leu Glu Glu Glu Thr Ala Lys Phe Phe Thr
 65 70 75 80

Gln Val His Gln Ala Ile Lys Thr Leu Arg Asp Asp Lys Thr Val Leu
 85 90 95

Leu Glu Glu Ile Tyr Thr His Lys Asn Leu Phe Thr Glu Arg Leu Asn
 100 105 110

Lys Ile Ser Asp Gly Leu Lys Glu Lys Gly Ala Pro Pro Xaa Ser Met
 115 120 125

Asn Ala Phe Pro Ala Pro Ser Pro Thr Cys Thr Pro Glu Pro Leu Gly
 130 135 140

Ser Val Cys Leu Pro Ser Thr Ser Val Ser Leu Pro Ser His Leu Pro
 145 150 155 160

Gly Ser Leu Gln

<210> 133
 <211> 245
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (245)
 <223> Xaa equals stop translation

<400> 133
 Met Val Ala Val Gly Val Tyr Ala Arg Leu Met Lys His Ala Glu Ala
 1 5 10 15
 Ala Leu Ala Cys Leu Ala Val Asp Pro Ala Ile Leu Leu Ile Val Val
 20 25 30
 Gly Val Leu Met Phe Leu Leu Thr Phe Cys Gly Cys Ile Gly Ser Leu
 35 40 45
 Arg Glu Asn Ile Cys Leu Leu Gln Thr Phe Ser Leu Cys Leu Thr Ala
 50 55 60
 Val Phe Leu Leu Gln Leu Ala Ala Gly Ile Leu Gly Phe Val Phe Ser
 65 70 75 80
 Asp Lys Ala Arg Gly Lys Val Ser Glu Ile Ile Asn Asn Ala Ile Val
 85 90 95
 His Tyr Arg Asp Asp Leu Asp Leu Gln Asn Leu Ile Asp Phe Gly Gln
 100 105 110
 Lys Lys Phe Ser Cys Cys Gly Gly Ile Ser Tyr Lys Asp Trp Ser Gln
 115 120 125
 Asn Met Tyr Phe Asn Cys Ser Glu Asp Asn Pro Ser Arg Glu Arg Cys
 130 135 140
 Ser Val Pro Tyr Ser Cys Cys Leu Pro Thr Pro Asp Gln Ala Val Ile
 145 150 155 160
 Asn Thr Met Cys Gly Gln Gly Met Gln Ala Phe Asp Tyr Leu Glu Ala
 165 170 175
 Ser Lys Val Ile Tyr Thr Asn Gly Cys Ile Asp Lys Leu Val Asn Trp
 180 185 190
 Ile His Ser Asn Leu Phe Leu Leu Gly Gly Val Ala Leu Gly Leu Ala
 195 200 205
 Ile Pro Gln Leu Val Gly Ile Leu Leu Ser Gln Ile Leu Val Asn Gln
 210 215 220
 Ile Lys Asp Gln Ile Lys Leu Gln Leu Tyr Asn Gln Gln His Arg Ala
 225 230 235 240
 Asp Pro Trp Tyr Xaa

245

<210> 134
 <211> 56
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (56)
 <223> Xaa equals stop translation

<400> 134
 Met Gly Thr Val Gly Leu Trp Pro Ser Trp Leu Trp Leu Pro Ala Ser
 1 5 10 15
 Trp Pro Leu Thr Ser Cys Gly Val Thr Arg Arg Arg Leu Arg Gly Pro
 20 25 30
 Gly Leu Arg Arg Thr Ser Gln Thr Gly Arg His Thr Ser Pro Cys Pro
 35 40 45
 Thr Ala Thr Trp Ala Glu Ser Xaa
 50 55

<210> 135
 <211> 55
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (47)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (51)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (55)
 <223> Xaa equals stop translation

<400> 135
 Met Ser Ile Val Met Ser Pro Leu Leu Leu Pro Ile Cys Tyr Leu Asn
 1 5 10 15
 Leu Leu Leu Phe Phe Val Asn Leu Ala Lys Asn Leu Ser Ile Leu Phe
 20 25 30
 Val Ser Ser Lys Lys Tyr Thr Phe Val Phe Met Ile Ser Leu Xaa Phe
 35 40 45
 Phe His Xaa Tyr Phe Ile Xaa

94

50

55

<210> 136
 <211> 89
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (89)
 <223> Xaa equals stop translation

<400> 136
 Met Ala Ile Ile Ser Phe Glu Leu Leu Phe Leu Met Asn Leu Pro Thr
 1 5 10 15
 Val Asn Ser Ser Asn Phe Lys Leu Ile Ile Pro Glu Asp Val Thr Leu
 20 25 30
 Ser Phe Val Ser His Leu Asp Ile Thr Val Asn His Phe Val Phe Leu
 35 40 45
 Ser Thr Phe Glu Leu Ala Gly Val Ile Glu Gly Lys Pro Leu Pro Asp
 50 55 60
 Ser Lys Ser Asp Leu Cys Pro Ile Leu Gly Gln Leu Trp Phe His Ile
 65 70 75 80
 Leu Leu Phe Phe Ile Phe Trp Val Xaa
 85

<210> 137
 <211> 62
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (62)
 <223> Xaa equals stop translation

<400> 137
 Met Arg Leu Pro Ile Ala Pro His Leu Gln Tyr Phe Met Trp Ser Val
 1 5 10 15
 Leu Leu Phe Leu Val Ile Leu Val Asp Met Lys Trp His Leu Ser Val
 20 25 30
 Ala Phe His Tyr Ile Ser Leu Met Thr Asn Gly Ile Leu Ser Pro Phe
 35 40 45
 Gln Cys Leu Leu Ala Ile His Val Ser Leu Phe Phe Val Xaa
 50 55 60

<210> 138

95

<211> 106
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (106)
<223> Xaa equals stop translation

<400> 138

Met Cys Leu Leu Pro Gly Gly Val Leu Leu Ile Trp Ser Cys Ala Ser
1 5 10 15
Gly Thr Pro Ala Ser His Thr Lys Asp Trp Gly Arg Cys Lys Phe Ser
20 25 30
Ala Ala Thr Lys Arg Thr Ala Glu Ser Asn Leu Glu Ser Thr Gln Leu
35 40 45
Met Leu Ala Ser Gln Ile Asp Pro Leu Leu Ala Glu Cys Trp His Leu
50 55 60
Cys Ala Ser Val Ser Ser Ser Val Asn Gly Gly Asp Lys Lys Cys Val
65 70 75 80
His Thr Ser Arg Ala Val Gly Arg Ile Lys Leu Cys Ser Asp Thr Ile
85 90 95
Arg Ala Cys Ser Gly Trp Tyr Leu Gln Xaa
100 105

<210> 139
<211> 52
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (52)
<223> Xaa equals stop translation

<400> 139

Met Ser His Ser Val Phe Ala His Tyr Ile Phe Asn Ile Leu Leu Leu
1 5 10 15
Leu Leu Leu Leu Leu Leu Ile Gly Phe Leu Tyr Ser Met Pro Phe Ile
20 25 30
Tyr Lys Asp Thr Lys Lys Thr His Val Cys Asn Phe Asn Asn Ile Phe
35 40 45
Pro Ile Leu Xaa
50

<210> 140
<211> 119

96

<212> PRT

<213> Homo sapiens

<400> 140

```

Met Lys Trp Arg Arg Lys Ser Ala Tyr Trp Lys Ala Leu Lys Val Phe
  1              5              10              15

Lys Leu Pro Val Glu Phe Leu Leu Leu Thr Val Pro Val Val Asp
      20              25              30

Pro Asp Lys Asp Asp Gln Asn Trp Lys Arg Pro Leu Asn Cys Leu His
      35              40              45

Leu Val Ile Ser Pro Leu Val Val Val Leu Thr Leu Gln Ser Gly Thr
      50              55              60

Tyr Gly Val Tyr Glu Ile Gly Gly Leu Val Pro Val Trp Val Val Val
      65              70              75              80

Val Ile Ala Gly Thr Ala Leu Ala Ser Val Thr Phe Phe Ala Thr Ser
      85              90              95

Asp Ser Gln Pro Pro Arg Leu His Trp Leu Phe Ala Phe Leu Gly Phe
      100             105             110

Leu Thr Ser Ala Leu Trp Ile
      115

```

<210> 141

<211> 59

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (59)

<223> Xaa equals stop translation

<400> 141

```

Met Cys Ser Gly Ser Phe Lys Glu Leu Tyr Leu Val Pro Ile Ser Leu
  1              5              10              15

Phe Ser Thr Cys Val Leu Gly Phe Tyr Phe His Asn Phe Leu Leu Leu
      20              25              30

Ile Ile Leu Phe Ser Ile Leu Leu Arg Lys Ile Thr Gly Lys Leu Phe
      35              40              45

Phe Thr Tyr Tyr His Phe Ser Cys Gly Val Xaa
      50              55

```

<210> 142

<211> 100

<212> PRT

<213> Homo sapiens

97

```
<210> 143
<211> 65
<212> PRT
<213> Homo sapiens
```

```
<220>  
<221> SITE  
<222> (65)  
<223> Xaa equals stop translation
```

```

<400> 143
Met Ala Val Leu Leu Ile Thr Ile Leu Leu Phe Leu Cys Leu Gly Tyr
  1             5             10             15
Tyr Arg Val Ile Thr Glu Ile Ser Arg Lys Thr Pro Ala Cys Arg Met
          20             25             30
Phe Thr Ser Ser Leu Ser Ser Trp Tyr Ile Met Arg Lys Leu Tyr Asp
  35             40             45
Thr Pro Gly Glu Val Phe Leu Ser His Ala Ile Val Xaa Phe Leu Lys
  50             55             60
Xaa
  65

```


<210> 144
 <211> 67
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (67)
 <223> Xaa equals stop translation

<400> 144
 Met Leu Asn Gln Pro Cys Ile Leu Gly Met Lys Pro Thr Trp Leu Trp
 1 5 10 15
 Trp Ile Ser Phe Leu Met Cys Cys Trp Val Trp Leu Ala Ser Val Leu
 20 25 30
 Leu Gly Ile Phe Ala Ser Ile Phe Ile Arg Asp Ile Gly Leu Glu Phe
 35 40 45
 Ser Phe Phe Val Met Cys Leu Pro Gly Phe Gly Ile Arg Val Met Leu
 50 55 60
 Ala Ser Xaa
 65

<210> 145
 <211> 59
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (59)
 <223> Xaa equals stop translation

<400> 145
 Met Thr Ala Met Ser Ile His Leu Phe Cys Thr Ala Leu Ser Cys Gly
 1 5 10 15
 Ser Ser Gly Gln Cys Asn Lys Ala Ile Lys Arg Asn Lys Ile Ser Asn
 20 25 30
 Asp Trp Lys Asp Val Asn Val Ser Ser Phe Ile Glu Asn Met Ile His
 35 40 45
 Arg Tyr Thr Tyr Thr Asn Ala Leu Asn Ser Xaa
 50 55

<210> 146
 <211> 56
 <212> PRT
 <213> Homo sapiens

99

<220>
<221> SITE
<222> (56)
<223> Xaa equals stop translation

<400> 146
Met Ser His Cys Thr Trp Pro Val Cys Leu Phe Cys Leu Val Pro Pro
1 5 10 15
Pro Met Gly Asp Leu Lys Glu Val Cys Leu Pro His Arg Cys Pro Gly
20 25 30
Arg Thr Ala Cys Cys Ser Tyr Ser Glu Pro His Leu Gln Thr Glu Glu
35 40 45
Asp Arg Arg Thr Leu Ile Cys Xaa
50 55

<210> 147
<211> 66
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (66)
<223> Xaa equals stop translation

<400> 147
Met Thr Asn Gly His Gln Val Leu Leu Leu Leu Leu Thr Ser Ala
1 5 10 15
Val Ala Ala Gly Pro Trp Pro Gln Val His Ala Gly Gln Trp Gly Trp
20 25 30
Met Cys Leu Pro Pro Gly Leu Pro Ser Val Gln Ala Arg Ser Gly Leu
35 40 45
Gly Gly Leu Pro Gly Gly Pro Gln Trp Val Pro Gly Gly Ala Arg Gly
50 55 60
Tyr Xaa
65

<210> 148
<211> 328
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (328)
<223> Xaa equals stop translation

<400> 148
Met Ala Cys Arg Lys Leu Ala Val Ala His Pro Leu Leu Leu Arg

100

1	5	10	15
His Leu Pro Met Ile Ala Ala Leu Leu His Gly Arg Thr His Leu Asn	20	25	30
Phe Gln Glu Phe Arg Gln Gln Asn His Leu Ser Cys Phe Leu His Val	35	40	45
Leu Gly Leu Leu Glu Leu Leu Gln Pro His Val Phe Arg Ser Glu His	50	55	60
Gln Gly Ala Leu Trp Asp Cys Leu Leu Ser Phe Ile Arg Leu Leu Leu	65	70	75
Asn Tyr Arg Lys Ser Ser Arg His Leu Ala Ala Phe Ile Asn Lys Phe	85	90	95
Val Gln Phe Ile His Lys Tyr Ile Thr Tyr Asn Ala Pro Ala Ala Ile	100	105	110
Ser Phe Leu Gln Lys His Ala Asp Pro Leu His Asp Leu Ser Phe Asp	115	120	125
Asn Ser Asp Leu Val Met Leu Lys Ser Leu Leu Ala Gly Leu Ser Leu	130	135	140
Pro Ser Arg Asp Asp Arg Thr Asp Arg Gly Leu Asp Glu Glu Gly Glu	145	150	155
Glu Glu Ser Ser Ala Gly Ser Leu Pro Leu Val Ser Val Ser Leu Phe	165	170	175
Thr Pro Leu Thr Ala Ala Glu Met Ala Pro Tyr Met Lys Arg Leu Ser	180	185	190
Arg Gly Gln Thr Val Glu Asp Leu Leu Glu Val Leu Ser Asp Ile Asp	195	200	205
Glu Met Ser Arg Arg Arg Pro Glu Ile Leu Ser Phe Phe Ser Thr Asn	210	215	220
Leu Gln Arg Leu Met Ser Ser Ala Glu Glu Cys Cys Arg Asn Leu Ala	225	230	235
Phe Ser Leu Ala Leu Arg Ser Met Gln Asn Ser Pro Ser Ile Ala Ala	245	250	255
Ala Phe Leu Pro Thr Phe Met Tyr Cys Leu Gly Ser Gln Asp Phe Glu	260	265	270
Val Val Gln Thr Ala Leu Arg Asn Leu Pro Glu Tyr Ala Leu Leu Cys	275	280	285
Gln Glu His Ala Ala Val Leu Leu His Arg Ala Phe Leu Val Gly Met	290	295	300
Tyr Gly Gln Met Asp Pro Ser Ala Gln Ile Ser Glu Ala Leu Arg Ile	305	310	315
			320

101

Leu His Met Glu Ala Val Met Xaa
325

<210> 149

<211> 90

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (10)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (13)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (90)

<223> Xaa equals stop translation

<400> 149

Met Gly Phe Leu Gln Leu Leu Val Val Xaa Val Leu Xaa Ser Glu His
1 5 10 15

Arg Val Ala Gly Ala Ala Glu Val Phe Gly Asn Ser Ser Glu Gly Leu
20 25 30

Ile Glu Phe Ser Val Gly Lys Phe Arg Tyr Phe Glu Leu Asn Arg Pro
35 40 45

Phe Pro Glu Glu Ala Ile Leu His Asp Ile Ser Ser Asn Val Thr Phe
50 55 60

Leu Ile Phe Gln Ile His Ser Gln Tyr Gln Asn Thr Thr Val Ser Phe
65 70 75 80

Ser Pro Arg Arg Arg Ser Pro Thr Met Xaa
85 90

<210> 150

<211> 149

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (149)

<223> Xaa equals stop translation

<400> 150

Met Ala Gly Ser Pro Leu Leu Trp Gly Pro Arg Ala Gly Gly Val Gly
1 5 10 15

102

Leu Leu Val Leu Leu Leu Leu Gly Leu Phe Arg Pro Pro Pro Ala Leu
 20 25 30
 Cys Ala Arg Pro Val Lys Glu Pro Arg Gly Leu Ser Ala Ala Ser Pro
 35 40 45
 Pro Leu Ala Arg Leu Ala Leu Leu Ala Ala Ser Gly Gly Gln Cys Pro
 50 55 60
 Glu Val Arg Arg Arg Gly Arg Cys Arg Pro Gly Ala Gly Ala Gly Ala
 65 70 75 80
 Ser Ala Gly Ala Glu Arg Gln Glu Arg Ala Arg Ala Glu Ala Gln Arg
 85 90 95
 Leu Arg Ile Ser Arg Arg Ala Ser Trp Arg Ser Cys Cys Ala Ser Gly
 100 105 110
 Ala Pro Pro Ala Thr Leu Ile Arg Leu Trp Ala Trp Thr Thr Thr Pro
 115 120 125
 Thr Arg Leu Gln Arg Ser Ser Leu Ala Leu Cys Ser Ala Pro Ala Leu
 130 135 140
 Thr Leu Pro Pro Xaa
 145

<210> 151
 <211> 391
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (391)
 <223> Xaa equals stop translation

<400> 151
 Met Leu Pro Thr Trp Trp Ile Val Ser Ser Trp Leu Val Trp Gly Val
 1 5 10 15
 Ile Leu Phe Val Tyr Leu Val Ile Arg Ala Leu Arg Leu Trp Arg Thr
 20 25 30
 Ala Lys Leu Gln Val Thr Leu Lys Lys Tyr Ser Val His Leu Glu Asp
 35 40 45
 Met Ala Thr Asn Ser Arg Ala Phe Thr Asn Leu Val Arg Lys Ala Leu
 50 55 60
 Arg Leu Ile Gln Glu Thr Glu Val Ile Ser Arg Gly Phe Thr Leu Val
 65 70 75 80
 Ser Ala Ala Cys Pro Phe Asn Lys Ala Gly Gln His Pro Ser Gln His
 85 90 95

103

Leu Ile Gly Leu Arg Lys Ala Val Tyr Arg Thr Leu Arg Ala Asn Phe
 100 105 110
 Gln Ala Ala Arg Leu Ala Thr Leu Tyr Met Leu Lys Asn Tyr Pro Leu
 115 120 125
 Asn Ser Glu Ser Asp Asn Val Thr Asn Tyr Ile Cys Val Val Pro Phe
 130 135 140
 Lys Glu Leu Gly Leu Gly Leu Ser Glu Glu Gln Ile Ser Glu Glu Glu
 145 150 155 160
 Ala His Asn Phe Thr Asp Gly Phe Ser Leu Pro Ala Leu Lys Val Leu
 165 170 175
 Phe Gln Leu Trp Val Ala Gln Ser Ser Glu Phe Phe Arg Arg Leu Ala
 180 185 190
 Leu Leu Leu Ser Thr Ala Asn Ser Pro Pro Gly Pro Leu Leu Thr Pro
 195 200 205
 Ala Leu Leu Pro His Arg Ile Leu Ser Asp Val Thr Gln Gly Leu Pro
 210 215 220
 His Ala His Ser Ala Cys Leu Glu Glu Leu Lys Arg Ser Tyr Glu Phe
 225 230 235 240
 Tyr Arg Tyr Phe Glu Thr Gln His Gln Ser Val Pro Gln Cys Leu Ser
 245 250 255
 Lys Thr Gln Gln Lys Ser Arg Glu Leu Asn Asn Val His Thr Ala Val
 260 265 270
 Arg Ser Leu Gln Leu His Leu Lys Ala Leu Leu Asn Glu Val Ile Ile
 275 280 285
 Leu Glu Asp Glu Leu Glu Lys Leu Val Cys Thr Lys Glu Thr Gln Glu
 290 295 300
 Leu Val Ser Glu Ala Tyr Pro Ile Leu Glu Gln Lys Leu Lys Leu Ile
 305 310 315 320
 Gln Pro His Val Gln Ala Ser Asn Asn Cys Trp Glu Glu Ala Ile Ser
 325 330 335
 Gln Val Asp Lys Leu Leu Arg Arg Asn Thr Asp Lys Lys Gly Lys Pro
 340 345 350
 Glu Ile Ala Cys Glu Asn Pro His Cys Thr Val Ser Thr Phe Glu Ala
 355 360 365
 Ala Tyr Ser Thr His Cys Arg Gln Arg Ser Asn Pro Arg Gly Ala Gly
 370 375 380
 Ile Arg Ser Leu Cys Arg Xaa
 385 390

104

<210> 152
 <211> 99
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (99)
 <223> Xaa equals stop translation

<400> 152
 Met Thr Thr Arg Gln Pro Thr Ala Val Ser Trp Pro Cys Trp Leu Met
 1 5 10 15
 Ser Ser Ser Leu Ser Thr Ala Cys Leu Ala Trp Thr Leu Thr Gly Ser
 20 25 30
 Leu Ala Arg Glu Ala Thr Arg Arg Ala Arg Ser Leu Ser Pro Thr Trp
 35 40 45
 Asn Cys Ser Ala Arg Gln Val Pro Pro Ser Pro Pro His Ser Gly Leu
 50 55 60
 Gly Arg Arg Gly Trp Ala His Cys His Leu Thr Cys Leu Leu Val Thr
 65 70 75 80
 Gln Leu Phe Arg Val Gly Arg Ile His Pro Ile Leu Ser Leu Pro Leu
 85 90 95
 Val Thr Xaa

<210> 153
 <211> 61
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (61)
 <223> Xaa equals stop translation

<400> 153
 Met Ser His Cys Ala Arg Pro Thr Phe Leu Thr Leu Leu Leu Ala Ser
 1 5 10 15
 Cys Phe Trp Ala Ala Ala Ile Pro Asn Arg Asn Val Ile Leu Ser Val
 20 25 30
 Ser Phe Arg Pro Leu His Met Gln Phe Thr Leu Ser Ile Leu Val Phe
 35 40 45
 Ile Leu Arg Ile Leu Ile Leu Leu Arg Ser Phe Leu Xaa
 50 55 60

<210> 154

105

<211> 393
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (393)
 <223> Xaa equals stop translation

<400> 154

Met	Glu	Trp	Trp	Ala	Ser	Ser	Pro	Leu	Arg	Leu	Trp	Leu	Leu	Leu	Phe
1				5					10					15	
Leu	Leu	Pro	Ser	Ala	Gln	Gly	Arg	Gln	Lys	Glu	Ser	Gly	Ser	Lys	Trp
		20					25						30		
Lys	Val	Phe	Ile	Asp	Gln	Ile	Asn	Arg	Ser	Leu	Glu	Asn	Tyr	Glu	Pro
		35					40					45			
Cys	Ser	Ser	Gln	Asn	Cys	Ser	Cys	Tyr	His	Gly	Val	Ile	Glu	Glu	Asp
	50					55					60				
Leu	Thr	Pro	Phe	Arg	Gly	Ile	Ser	Arg	Lys	Met	Met	Ala	Glu	Val	
65					70				75					80	
Val	Arg	Arg	Lys	Leu	Gly	Thr	His	Tyr	Gln	Ile	Thr	Lys	Asn	Arg	Leu
				85					90					95	
Tyr	Arg	Glu	Asn	Asp	Cys	Met	Phe	Pro	Ser	Arg	Cys	Ser	Gly	Val	Glu
		100						105					110		
His	Phe	Ile	Leu	Glu	Val	Ile	Gly	Arg	Leu	Pro	Asp	Met	Glu	Met	Val
		115					120					125			
Ile	Asn	Val	Arg	Asp	Tyr	Pro	Gln	Val	Pro	Lys	Trp	Met	Glu	Pro	Ala
	130						135				140				
Ile	Pro	Val	Phe	Ser	Phe	Ser	Lys	Thr	Ser	Glu	Tyr	His	Asp	Ile	Met
145				150						155				160	
Tyr	Pro	Ala	Trp	Thr	Phe	Trp	Glu	Gly	Gly	Pro	Ala	Val	Trp	Pro	Ile
			165					170						175	
Tyr	Pro	Thr	Gly	Leu	Gly	Arg	Trp	Asp	Leu	Phe	Arg	Glu	Asp	Leu	Val
		180					185						190		
Arg	Ser	Ala	Ala	Gln	Trp	Pro	Trp	Lys	Lys	Lys	Asn	Ser	Thr	Ala	Tyr
		195					200					205			
Phe	Arg	Gly	Ser	Arg	Thr	Ser	Pro	Glu	Arg	Asp	Pro	Leu	Ile	Leu	Leu
	210					215					220				
Ser	Arg	Lys	Asn	Pro	Lys	Leu	Val	Asp	Ala	Glu	Tyr	Thr	Lys	Asn	Gln
225				230						235				240	
Ala	Trp	Lys	Ser	Met	Lys	Asp	Thr	Leu	Gly	Lys	Pro	Ala	Ala	Lys	Asp
			245						250					255	

106

Val His Leu Val Asp His Cys Lys Tyr Lys Tyr Leu Phe Asn Phe Arg
260 265 270

Gly Val Ala Ala Ser Phe Arg Phe Lys His Leu Phe Leu Cys Gly Ser
275 280 285

Leu Val Phe His Val Gly Asp Glu Trp Leu Glu Phe Phe Tyr Pro Gln
290 295 300

Leu Lys Pro Trp Val His Tyr Ile Pro Val Lys Thr Asp Leu Ser Asn
305 310 315 320

Val Gln Glu Leu Leu Gln Phe Val Lys Ala Asn Asp Asp Val Ala Gln
325 330 335

Glu Ile Ala Glu Arg Gly Ser Gln Phe Ile Arg Asn His Leu Gln Met
340 345 350

Asp Asp Ile Thr Cys Tyr Trp Glu Asn Leu Leu Ser Glu Tyr Ser Lys
355 360 365

Phe Leu Ser Tyr Asn Val Thr Arg Arg Lys Gly Tyr Asp Gln Ile Ile
370 375 380

Pro Lys Met Leu Lys Thr Glu Leu Xaa
385 390

<210> 155

<211> 75

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (75)

<223> Xaa equals stop translation

<400> 155

Met Leu Ile Leu Phe Leu Ser Val Cys Leu Phe Val Phe Leu Leu Thr
1 5 10 15

Val Arg Ala Leu Cys Cys Arg Ser Ala Gly Val Trp Leu Arg Ser Thr
20 25 30

Pro Asp Pro Val Cys Leu Gly Phe Ala Arg Gly Gly Cys Arg Ile Ala
35 40 45

Met Ile Ala Ala Cys Phe Ser Ser Gly Ser Phe Val Pro Glu Gly His
50 55 60

Pro Pro Asp Ala Ser Gln Ser Ser Pro Val Xaa
65 70 75

<210> 156

<211> 82

<212> PRT

107

<213> Homo sapiens

<220>

<221> SITE

<222> (82)

<223> Xaa equals stop translation

<400> 156

Met	Trp	Pro	Leu	Leu	Ala	Val	Ser	Pro	Phe	Gly	Leu	Val	Trp	Ala	Ser
1				5					10					15	

Ser	Gln	Ser	Gly	Ser	Leu	Leu	Leu	Arg	Ala	Ser	Ile	Pro	Arg	Gln	His
			20					25						30	

Ser	Arg	Arg	Ala	Trp	His	Phe	Tyr	Ser	Glu	Val	Trp	Gln	Ser	His	Ser
			35				40						45		

Val	Ala	Ser	Val	Leu	Leu	Tyr	Leu	Leu	Val	Arg	Ala	Ile	Thr	Lys	Met
	50					55					60				

Cys	Ile	Gly	Ser	Lys	Lys	Arg	Asp	Ile	Thr	Pro	Thr	Thr	Arg	Trp	Lys
	65				70					75					80

Lys Xaa

<210> 157

<211> 54

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (49)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (54)

<223> Xaa equals stop translation

<400> 157

Met	Ser	His	His	Ala	Gly	Leu	Gly	Gly	Gly	Ile	Leu	Phe	Ser	Leu	Lys
1				5					10					15	

Ile	Ser	Phe	Phe	Ile	Ala	Leu	Ala	Val	Val	Gly	Gly	Ser	Arg	Gly	Val
				20				25					30		

Asn	Asp	Cys	Gln	Leu	Gly	Gly	Cys	Arg	Val	Gly	Ser	Cys	Pro	Arg	Val
			35				40					45			

Xaa	Val	Arg	Val	Ala	Xaa
	50				

<210> 158

<211> 103

108

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (103)

<223> Xaa equals stop translation

<400> 158

Met Thr Val Arg Arg Leu Ser Leu Leu Cys Arg Asp Leu Trp Ala Leu
 1 5 10 15

Trp Leu Leu Leu Lys Ala Gly Ala Val Arg Gly Ala Arg Ala Gly Pro
 20 25 30

Arg Leu Pro Gly Arg Cys Cys Gly Ala Thr Cys Gly Asp Ala Gly Arg
 35 40 45

Gly Trp Thr Phe Trp Ala Gln Pro Cys Pro Gln Lys Leu Leu Gly Gln
 50 55 60

Lys Pro Gly Ala Gly Gly Cys Arg Gly Trp Val Leu Gly Trp Val Pro
 65 70 75 80

Pro Arg Pro Glu Glu Pro Cys Ser Leu Ala Gly Lys Val Cys Thr Gly
 85 90 95

Leu Ala Arg Trp Met Val Xaa
 100

<210> 159

<211> 575

<212> PRT

<213> Homo sapiens

<400> 159

Met Arg Val Leu Val Val Thr Ile Ala Pro Ile Tyr Trp Ala Leu Ala
 1 5 10 15

Arg Glu Ser Gly Glu Ala Leu Asn Gly His Ser Leu Thr Gly Gly Lys
 20 25 30

Phe Arg Gln Glu Ser His Val Glu Phe Ala Thr Gly Glu Leu Leu Thr
 35 40 45

Met Thr Gln Trp Pro Gly Val Trp Ile Pro Met Ala Ser Cys Ser Ser
 50 55 60

Thr Trp Trp Ser Met Ala Leu Ser Pro Asp Ser Leu Ala Asp Ala Asp
 65 70 75 80

Leu Gln Val Gln Asp Phe Glu Glu His Tyr Val Gln Thr Gly Pro Gly
 85 90 95

Gln Leu Phe Val Gly Ser Thr Gln Arg Phe Phe Gln Gly Gly Leu Pro
 100 105 110

109

Ser Phe Leu Arg Cys Asn His Ser Ile Gln Tyr Asn Ala Ala Arg Gly
 115 120 125
 Pro Gln Pro Gln Leu Val Gln His Leu Arg Ala Ser Ala Ile Ser Ser
 130 135 140
 Ala Phe Asp Pro Glu Ala Glu Ala Leu Arg Phe Gln Leu Ala Thr Ala
 145 150 155 160
 Leu Gln Ala Glu Glu Asn Glu Val Gly Cys Pro Glu Gly Phe Glu Leu
 165 170 175
 Asp Ser Gln Gly Ala Phe Cys Val Asp Val Asp Glu Cys Ala Trp Asp
 180 185 190
 Ala His Leu Cys Arg Glu Gly Gln Arg Cys Val Asn Leu Leu Gly Ser
 195 200 205
 Tyr Arg Cys Leu Pro Asp Cys Gly Pro Gly Phe Arg Val Ala Asp Gly
 210 215 220
 Ala Gly Cys Glu Asp Val Asp Glu Cys Leu Glu Gly Leu Asp Asp Cys
 225 230 235 240
 His Tyr Asn Gln Leu Cys Glu Asn Thr Pro Gly Gly His Arg Cys Ser
 245 250 255
 Cys Pro Arg Gly Tyr Arg Met Gln Gly Pro Ser Leu Pro Cys Leu Asp
 260 265 270
 Val Asn Glu Cys Leu Gln Leu Pro Lys Ala Cys Ala Tyr Gln Cys His
 275 280 285
 Asn Leu Gln Gly Ser Tyr Arg Cys Leu Cys Pro Pro Gly Gln Thr Leu
 290 295 300
 Leu Arg Asp Gly Lys Ala Cys Thr Ser Leu Glu Arg Asn Gly Gln Asn
 305 310 315 320
 Val Thr Thr Val Ser His Arg Gly Pro Leu Leu Pro Trp Leu Arg Pro
 325 330 335
 Trp Ala Ser Ile Pro Gly Thr Ser Tyr His Ala Trp Val Ser Leu Arg
 340 345 350
 Pro Gly Pro Met Ala Leu Ser Ser Val Gly Arg Ala Trp Cys Pro Pro
 355 360 365
 Gly Phe Ile Arg Gln Asn Gly Val Cys Thr Asp Leu Asp Glu Cys Arg
 370 375 380
 Val Arg Asn Leu Cys Gln His Ala Cys Arg Asn Thr Glu Gly Ser Tyr
 385 390 395 400
 Gln Cys Leu Cys Pro Ala Gly Tyr Arg Leu Leu Pro Ser Gly Lys Asn
 405 410 415
 Cys Gln Asp Ile Asn Glu Cys Glu Glu Glu Ser Ile Glu Cys Gly Pro

110

420

425

430

Gly Gln Met Cys Phe Asn Thr Arg Gly Ser Tyr Gln Cys Val Asp Thr
 435 440 445
 Pro Cys Pro Ala Thr Tyr Arg Gln Gly Pro Ser Pro Gly Thr Cys Phe
 450 455 460
 Arg Arg Cys Ser Gln Asp Cys Gly Thr Gly Gly Pro Ser Thr Leu Gln
 465 470 475 480
 Tyr Arg Leu Leu Pro Leu Pro Leu Gly Val Arg Ala His His Asp Val
 485 490 495
 Ala Arg Leu Thr Ala Phe Ser Glu Val Gly Val Pro Ala Asn Arg Thr
 500 505 510
 Glu Leu Ser Met Leu Glu Pro Asp Pro Arg Ser Pro Phe Ala Leu Arg
 515 520 525
 Pro Leu Arg Ala Gly Leu Gly Ala Val Tyr Thr Arg Arg Ala Leu Thr
 530 535 540
 Arg Ala Gly Leu Tyr Arg Leu Thr Val Arg Ala Ala Ala Pro Arg His
 545 550 555 560
 Gln Ser Val Phe Val Leu Leu Ile Ala Val Ser Pro Tyr Pro Tyr
 565 570 575

<210> 160

<211> 643

<212> PRT

<213> Homo sapiens

<400> 160

Met Gly Glu Pro Asn Arg His Pro Ser Met Phe Leu Leu Leu Leu Val
 1 5 10 15
 Leu Glu Arg Leu Tyr Ala Ser Pro Met Asp Gly Thr Ser Ser Ala Leu
 20 25 30
 Ser Met Gly Pro Phe Val Pro Phe Ile Met Arg Cys Gly His Ser Pro
 35 40 45
 Val Tyr His Ser Arg Glu Met Ala Ala Arg Ala Leu Val Pro Phe Val
 50 55 60
 Met Ile Asp His Ile Pro Asn Thr Ile Arg Thr Leu Leu Ser Thr Leu
 65 70 75 80
 Pro Ser Cys Thr Asp Gln Cys Phe Arg Gln Asn His Ile His Gly Thr
 85 90 95
 Leu Leu Gln Val Phe His Leu Leu Gln Ala Tyr Ser Asp Ser Lys His
 100 105 110
 Gly Thr Asn Ser Asp Phe Gln His Glu Leu Thr Asp Ile Thr Val Cys

111

115	120	125
Thr Lys Ala Lys Leu Trp Leu Ala Lys Arg Gln Asn Pro Cys Leu Val		
130	135	140
Thr Arg Ala Val Tyr Ile Asp Ile Leu Phe Leu Leu Thr Cys Cys Leu		
145	150	155
Asn Arg Ser Ala Lys Asp Asn Gln Pro Val Leu Glu Ser Leu Gly Phe		
165	170	175
Trp Glu Glu Val Arg Gly Ile Ile Ser Gly Ser Glu Leu Ile Thr Gly		
180	185	190
Phe Pro Trp Ala Phe Lys Val Pro Gly Leu Pro Gln Tyr Leu Gln Ser		
195	200	205
Leu Thr Arg Leu Ala Ile Ala Ala Val Trp Ala Ala Ala Ala Lys Ser		
210	215	220
Gly Glu Arg Glu Thr Asn Val Pro Ile Ser Phe Ser Gln Leu Leu Glu		
225	230	235
Ser Ala Phe Pro Glu Val Arg Ser Leu Thr Leu Glu Ala Leu Leu Glu		
245	250	255
Lys Phe Leu Ala Ala Ala Ser Gly Leu Gly Glu Lys Gly Val Pro Pro		
260	265	270
Leu Leu Cys Asn Met Gly Glu Lys Phe Leu Leu Leu Ala Met Lys Glu		
275	280	285
Asn His Pro Glu Cys Phe Cys Lys Ile Leu Lys Ile Leu His Cys Met		
290	295	300
Asp Pro Gly Glu Trp Leu Pro Gln Thr Glu His Cys Val His Leu Thr		
305	310	315
Pro Lys Glu Phe Leu Ile Trp Thr Met Asp Ile Ala Ser Asn Glu Arg		
325	330	335
Ser Glu Ile Gln Ser Val Ala Leu Arg Leu Ala Ser Lys Val Ile Ser		
340	345	350
His His Met Gln Thr Cys Val Glu Asn Arg Glu Leu Ile Ala Ala Glu		
355	360	365
Leu Lys Gln Trp Val Gln Leu Val Ile Leu Ser Cys Glu Asp His Leu		
370	375	380
Pro Thr Glu Ser Arg Leu Ala Val Val Glu Val Leu Thr Ser Thr Thr		
385	390	395
Pro Leu Phe Leu Thr Asn Pro His Pro Ile Leu Glu Leu Gln Asp Thr		
405	410	415
Leu Ala Leu Trp Lys Cys Val Leu Thr Leu Leu Gln Ser Glu Glu Gln		
420	425	430

112

```
<210> 161
<211> 191
<212> PRT
<213> Homo sapiens
```

```
<220>
<221> SITE
<222> (191)
<223> Xaa equals stop translation
```

<400> 161
Met Ser Ser Gly Thr Glu Leu Leu Trp Pro Gly Ala Ala Leu Leu Val
1 5 10 15

113

Leu Leu Gly Val Ala Ala Ser Leu Cys Val Arg Cys Ser Arg Pro Gly
 20 25 30
 Ala Lys Arg Ser Glu Lys Ile Tyr Gln Gln Arg Ser Leu Arg Glu Asp
 35 40 45
 Gln Gln Ser Phe Thr Gly Ser Arg Thr Tyr Ser Leu Val Gly Gln Ala
 50 55 60
 Trp Pro Gly Pro Leu Ala Asp Met Ala Pro Thr Arg Lys Asp Lys Leu
 65 70 75 80
 Leu Gln Phe Tyr Pro Ser Leu Glu Asp Pro Ala Ser Ser Arg Tyr Gln
 85 90 95
 Asn Phe Ser Lys Gly Ser Arg His Gly Ser Glu Glu Ala Tyr Ile Asp
 100 105 110
 Pro Ile Ala Met Glu Tyr Tyr Asn Trp Gly Arg Phe Ser Lys Pro Pro
 115 120 125
 Glu Asp Asp Asp Ala Asn Ser Tyr Glu Asn Val Leu Ile Cys Lys Gln
 130 135 140
 Lys Thr Thr Glu Thr Gly Ala Gln Gln Glu Gly Ile Gly Gly Leu Cys
 145 150 155 160
 Arg Gly Asp Leu Ser Leu Ser Leu Ala Leu Lys Thr Gly Pro Thr Ser
 165 170 175
 Gly Leu Cys Pro Ser Ala Ser Pro Glu Glu Asp Glu Gly Ile Xaa
 180 185 190

<210> 162
 <211> 64
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (64)
 <223> Xaa equals stop translation

<400> 162
 Met Lys His Val Leu Asn Leu Tyr Leu Leu Gly Val Val Leu Thr Leu
 1 5 10 15
 Leu Ser Ile Phe Val Arg Val Met Glu Ser Leu Glu Gly Leu Leu Glu
 20 25 30
 Ser Pro Ser Pro Gly Thr Ser Trp Thr Thr Arg Ser Gln Leu Ala Asn
 35 40 45
 Thr Glu Pro Thr Lys Gly Leu Pro Asp His Pro Ser Arg Ser Met Xaa
 50 55 60

114

<210> 163
 <211> 118
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (118)
 <223> Xaa equals stop translation

<400> 163
 Met Ile Phe Leu Thr Val Leu Pro Leu Ala Phe Leu Phe Leu His Ser
 1 5 10 15
 Gly Phe Tyr His Tyr Ile Ser Phe Ser Cys Leu Phe Ser Leu Ser Leu
 20 25 30
 Ala Leu Phe Phe Phe Leu Asp Val Ala Thr Phe Arg Arg Pro Gly Gln
 35 40 45
 Leu Phe Cys Glu Arg Ser Val Leu Phe Asp Met Phe His Phe Gly Phe
 50 55 60
 Val Ser Leu Phe Leu His Glu Trp Ile Gln Ala Lys His Phe Trp Ala
 65 70 75 80
 Gly Leu Phe Ile Val Leu Pro Ser Asp Val Phe Phe Ser Val His His
 85 90 95
 Leu Glu Ala Pro Asp Gly Ser Phe Pro Asn Ile Ala Lys Leu Ser Leu
 100 105 110
 Ile Ile Leu Leu Arg Xaa
 115

<210> 164
 <211> 43
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (43)
 <223> Xaa equals stop translation

<400> 164
 Met Leu Leu Gln Phe Thr Leu Trp Val Phe Gly Ala Ile His Phe Pro
 1 5 10 15
 Lys Cys Leu Gly Ile Lys Glu Glu Leu Leu Lys Cys Cys Leu Gln Leu
 20 25 30
 Pro Pro Ser Ser Thr Tyr Glu Lys Val Val Xaa
 35 40

115

<210> 165
 <211> 48
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (48)
 <223> Xaa equals stop translation

<400> 165
 Met Leu Ser Arg Arg Leu His Cys Leu Val Leu Tyr Phe Leu Leu Leu
 1 5 10 15
 Leu Leu Ser Phe Ile His Thr Leu Ser Val Ser His Ile Cys Ser Ser
 20 25 30
 Phe Ile Trp Leu Phe Pro Lys Asn Ile Glu Ser Glu Ala Thr Met Xaa
 35 40 45

<210> 166
 <211> 46
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (46)
 <223> Xaa equals stop translation

<400> 166
 Met Glu Lys Met Gly Gln Gly Leu Leu Ser Ser Thr Tyr Leu Thr Val
 1 5 10 15
 Leu His Leu Ile Gln Leu Val Gly Cys Gly Leu Leu Thr Glu Glu Ile
 20 25 30
 Lys Glu Ser Lys Tyr Leu Ile Lys Thr Leu Gly Ser Gly Xaa
 35 40 45

<210> 167
 <211> 207
 <212> PRT
 <213> Homo sapiens

<400> 167
 Met Ile Lys His Val Ala Trp Leu Ile Phe Thr Asn Cys Ile Phe Phe
 1 5 10 15
 Cys Pro Val Ala Phe Phe Ser Phe Ala Pro Leu Ile Thr Ala Ile Ser
 20 25 30

116

Ile Ser Pro Glu Ile Met Lys Ser Val Thr Leu Ile Phe Phe Pro Leu
 35 40 45
 Pro Ala Cys Leu Asn Pro Val Leu Tyr Val Phe Phe Asn Pro Lys Phe
 50 55 60
 Lys Glu Asp Trp Lys Leu Leu Lys Arg Arg Val Thr Lys Lys Ser Gly
 65 70 75 80
 Ser Val Ser Val Ser Ile Ser Ser Gln Gly Gly Cys Leu Glu Gln Asp
 85 90 95
 Phe Tyr Tyr Asp Cys Gly Met Tyr Ser His Leu Gln Gly Asn Leu Thr
 100 105 110
 Val Cys Asp Cys Cys Glu Ser Phe Leu Leu Thr Lys Pro Val Ser Cys
 115 120 125
 Lys His Leu Ile Lys Ser His Ser Cys Pro Ala Leu Ala Val Ala Ser
 130 135 140
 Cys Gln Arg Pro Glu Gly Tyr Trp Ser Asp Cys Gly Thr Gln Ser Ala
 145 150 155 160
 His Ser Asp Tyr Ala Asp Glu Glu Asp Ser Phe Val Ser Asp Ser Ser
 165 170 175
 Asp Gln Val Gln Ala Cys Gly Arg Ala Cys Phe Tyr Gln Ser Arg Gly
 180 185 190
 Phe Pro Leu Val Arg Tyr Ala Tyr Asn Leu Pro Arg Val Lys Asp
 195 200 205

<210> 168

<211> 51

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (51)

<223> Xaa equals stop translation

<400> 168

Met Tyr Ile Phe Glu Leu Ser Leu Tyr Leu Glu Gly Thr Ser Phe Val
 1 5 10 15
 Val Val Leu Leu Phe Leu Leu Ile Ser Val Ser Leu Asp Ser Pro Pro
 20 25 30
 Thr Thr Lys Gly Trp Asp Ser Val Leu His Ile Trp Val Pro Leu Ile
 35 40 45
 Val Gln Xaa
 50

117

<210> 169
 <211> 43
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (43)
 <223> Xaa equals stop translation

<400> 169
 Met Ala His Pro Gly Leu Pro Lys Thr Val Pro Val Tyr Ala Val Val
 1 5 10 15
 Leu Ala Leu Leu Ile Met Thr Leu Pro Leu Thr Leu Thr Ile Asn Leu
 20 25 30
 Asp Asp Asn Leu Tyr Gly Asn Ser Ala Lys Xaa
 35 40

<210> 170
 <211> 56
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (56)
 <223> Xaa equals stop translation

<400> 170
 Met Arg Pro Trp Trp Ser Leu Leu Leu Glu Ala Cys Ala Thr Cys Ala
 1 5 10 15
 Gln Thr Gly Pro Thr Arg Ser Thr Ser Cys Thr Gln Glu Val Ser His
 20 25 30
 Ser Ser Ser Thr Ala Tyr Pro Ala Pro Met Arg Arg Arg Cys Cys Leu
 35 40 45
 Pro Ser Pro Arg Ser Cys Thr Xaa
 50 55

<210> 171
 <211> 109
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (109)
 <223> Xaa equals stop translation

<400> 171
 Met Ala Leu Ala Gly Ser Val Phe Val Leu Gly Gly Val Leu Val Leu

118

1 5 10 15
 Cys Val Glu Arg Asn Gly Glu Gly Glu Met Gly Trp Pro Gln His Leu
 20 25 30
 Pro Lys Ser Gln Pro Leu Ser Pro Pro Val Ala Val Arg Arg Cys Ser
 35 40 45
 Phe Glu Arg Ser Trp Ile Asp Leu Leu Val Glu Thr Ser Ser Ser Met
 50 55 60
 Val Thr Cys Arg Gln Gln Val Gly Thr Pro Asn Gly Met Glu Gly Arg
 65 70 75 80
 Gly Gly Gly Pro Lys Thr Thr Phe Pro Ile Arg Leu Gln Leu Ser Gly
 85 90 95
 Ala Cys Ala Val Arg Pro Glu Ile Gln Trp Glu Val Xaa
 100 105

<210> 172

<211> 51

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (17)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (51)

<223> Xaa equals stop translation

<400> 172

Met Phe Leu Phe Phe Tyr Leu Ser Leu Ala Val Tyr Ala Gln Arg Gln
 1 5 10 15

Xaa Ser Gly Ser Cys Arg Gln Thr Asp His Arg Trp Lys Ser Arg Gly
 20 25 30

Ala Arg Arg Cys Phe Leu Glu Pro Arg Asp Pro Gly Ser Val Pro Gly
 35 40 45

His Pro Xaa
 50

<210> 173

<211> 566

<212> PRT

<213> Homo sapiens

<400> 173

Met Ala Pro Leu Ala Leu His Leu Leu Val Leu Val Pro Ile Leu Leu
 1 5 10 15

119

Ser Leu Val Ala Ser Gln Asp Trp Lys Ala Glu Arg Ser Gln Asp Pro
 20 25 30
 Phe Glu Lys Cys Met Gln Asp Pro Asp Tyr Glu Gln Leu Leu Lys Val
 35 40 45
 Val Thr Trp Gly Leu Asn Arg Thr Leu Lys Pro Gln Arg Val Ile Val
 50 55 60
 Val Gly Ala Gly Val Ala Gly Leu Val Ala Ala Lys Val Leu Ser Asp
 65 70 75 80
 Ala Gly His Lys Val Thr Ile Leu Glu Ala Asp Asn Arg Ile Gly Gly
 85 90 95
 Arg Ile Phe Thr Tyr Arg Asp Gln Asn Thr Gly Trp Ile Gly Glu Leu
 100 105 110
 Gly Ala Met Arg Met Pro Ser Ser His Arg Ile Leu His Lys Leu Cys
 115 120 125
 Gln Gly Leu Gly Leu Asn Leu Thr Lys Phe Thr Gln Tyr Asp Lys Asn
 130 135 140
 Thr Trp Thr Glu Val His Glu Val Lys Leu Arg Asn Tyr Val Val Glu
 145 150 155 160
 Lys Val Pro Glu Lys Leu Gly Tyr Ala Leu Arg Pro Gln Glu Lys Gly
 165 170 175
 His Ser Pro Glu Asp Ile Tyr Gln Met Ala Leu Asn Gln Ala Leu Lys
 180 185 190
 Asp Leu Lys Ala Leu Gly Cys Arg Lys Ala Met Lys Lys Phe Glu Arg
 195 200 205
 His Thr Leu Leu Glu Tyr Leu Leu Gly Glu Gly Asn Leu Ser Arg Pro
 210 215 220
 Ala Val Gln Leu Leu Gly Asp Val Met Ser Glu Asp Gly Phe Phe Tyr
 225 230 235 240
 Leu Ser Phe Ala Glu Ala Leu Arg Ala His Ser Cys Leu Ser Asp Arg
 245 250 255
 Leu Gln Tyr Ser Arg Ile Val Gly Gly Trp Asp Leu Leu Pro Arg Ala
 260 265 270
 Leu Leu Ser Ser Leu Ser Gly Leu Val Leu Leu Asn Ala Pro Val Val
 275 280 285
 Ala Met Thr Gln Gly Pro His Asp Val His Val Gln Ile Glu Thr Ser
 290 295 300
 Pro Pro Ala Arg Asn Leu Lys Val Leu Lys Ala Asp Val Val Leu Leu
 305 310 315 320

120

Thr Ala Ser Gly Pro Ala Val Lys Arg Ile Thr Phe Ser Pro Pro Leu
 325 330 335
 Pro Arg His Met Gln Glu Ala Leu Arg Arg Leu His Tyr Val Pro Ala
 340 345 350
 Thr Lys Val Phe Leu Ser Phe Arg Arg Pro Phe Trp Arg Glu Glu His
 355 360 365
 Ile Glu Gly Gly His Ser Asn Thr Asp Arg Pro Ser Arg Met Ile Phe
 370 375 380
 Tyr Pro Pro Pro Arg Glu Gly Ala Leu Leu Leu Ala Ser Tyr Thr Trp
 385 390 395 400
 Ser Asp Ala Ala Ala Ala Phe Ala Gly Leu Ser Arg Glu Glu Ala Leu
 405 410 415
 Arg Leu Ala Leu Asp Asp Val Ala Ala Leu His Gly Pro Val Val Arg
 420 425 430
 Gln Leu Trp Asp Gly Thr Gly Val Val Lys Arg Trp Ala Glu Asp Gln
 435 440 445
 His Ser Gln Gly Gly Phe Val Val Gln Pro Pro Ala Leu Trp Gln Thr
 450 455 460
 Glu Lys Asp Asp Trp Thr Val Pro Tyr Gly Arg Ile Tyr Phe Ala Gly
 465 470 475 480
 Glu His Thr Ala Tyr Pro His Gly Trp Val Glu Thr Ala Val Lys Leu
 485 490 495
 Leu Arg Ala Ala Ile Lys Ile Asn Ser Arg Lys Gly Pro Ala Ser Asp
 500 505 510
 Thr Ala Ser Pro Glu Gly His Ala Ser Asp Met Glu Gly Gln Gly His
 515 520 525
 Val His Gly Val Ala Ser Ser Pro Ser His Asp Leu Ala Lys Glu Glu
 530 535 540
 Gly Ser His Pro Pro Val Gln Gly Gln Leu Ser Leu Gln Asn Thr Thr
 545 550 555 560
 His Thr Arg Thr Ser His
 565

<210> 174

<211> 224

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (76)

<223> Xaa equals any of the naturally occurring L-amino acids

121

<400> 174

Met Ala Arg Ala Arg Gly Ser Pro Cys Pro Pro Leu Pro Pro Gly Arg
 1 5 10 15

Met Ser Trp Pro His Gly Ala Leu Leu Phe Leu Trp Leu Phe Ser Pro
 20 25 30

Pro Leu Gly Ala Gly Gly Gly Gly Val Ala Val Thr Ser Ala Ala Gly
 35 40 45

Gly Gly Ser Pro Pro Ala Thr Ser Cys Pro Val Ala Cys Ser Cys Ser
 50 55 60

Asn Gln Ala Ser Arg Val Ile Cys Thr Arg Arg Xaa Leu Ala Glu Val
 65 70 75 80

Pro Ala Ser Ile Pro Val Asn Thr Arg Tyr Leu Asn Leu Gln Glu Asn
 85 90 95

Gly Ile Gln Val Ile Arg Thr Asp Thr Phe Lys His Leu Arg His Leu
 100 105 110

Glu Ile Leu Gln Leu Ser Lys Asn Leu Val Arg Lys Ile Glu Val Gly
 115 120 125

Ala Phe Asn Gly Leu Pro Ser Leu Asn Thr Leu Glu Leu Phe Asp Asn
 130 135 140

Arg Leu Thr Thr Val Pro Thr Gln Ala Phe Glu Tyr Leu Ser Lys Leu
 145 150 155 160

Arg Glu Leu Trp Leu Arg Asn Asn Pro Ile Glu Ser Ile Pro Ser Tyr
 165 170 175

Ala Phe Asn Arg Val Pro Ser Leu Arg Arg Leu Asp Leu Gly Glu Leu
 180 185 190

Lys Arg Leu Glu Tyr Ile Ser Glu Ala Ala Phe Glu Gly Leu Val Asn
 195 200 205

Leu Arg Tyr Leu Asn Leu Gly Met Cys Asn Leu Lys Asp Ile Pro Asn
 210 215 220

<210> 175

<211> 123

<212> PRT

<213> Homo sapiens

<400> 175

Met His Asp Gly Ser Lys Pro Phe Pro Arg Tyr Gly Tyr Lys Pro Ser
 1 5 10 15

Pro Pro Asn Gly Cys Gly Ser Pro Leu Phe Gly Val His Leu Asn Ile

122

20	25	30
Gly Ile Pro Ser Leu Thr Lys Cys Cys Asn Gln His Asp Arg Cys Tyr		
35	40	45
Glu Thr Cys Gly Lys Ser Lys Asn Asp Cys Asp Glu Glu Phe Gln Tyr		
50	55	60
Cys Leu Ser Lys Ile Cys Arg Asp Val Gln Lys Thr Leu Gly Leu Thr		
65	70	75
Gln His Val Gln Ala Cys Glu Thr Thr Val Glu Leu Leu Phe Asp Ser		
85	90	95
Val Ile His Leu Gly Cys Lys Pro Tyr Leu Asp Ser Gln Arg Ala Ala		
100	105	110
Cys Arg Cys His Tyr Glu Glu Lys Thr Asp Leu		
115	120	

<210> 176

<211> 60

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (60)

<223> Xaa equals stop translation

<400> 176

Met Gly Leu Ser Val Leu Leu Pro Leu Cys Leu Leu Gly Pro Gly Arg		
1	5	10
Phe Thr Ser Gly Gln Lys Pro Leu Asp Thr Pro Gly Leu Gly Ala Ala		
20	25	30
Val Leu Ser Val Arg Lys Ala Gly Leu Lys Met Arg Ser His Leu Thr		
35	40	45
Pro Ser Val Cys Thr Val Pro Ser Pro Gly Ser Xaa		
50	55	60

<210> 177

<211> 105

<212> PRT

<213> Homo sapiens

<400> 177

Met Asp Thr Val Phe Leu Ile Gln Tyr Leu Phe Leu Thr Phe Pro Arg		
1	5	10
Ile Val Phe Met Leu Gly Phe Val Val Val Leu Ser Phe Leu Leu Gly		
20	25	30
Gly Tyr Leu Leu Phe Val Leu Tyr Leu Ala Ala Thr Asn Gln Thr Thr		

WO 00/04140

PCT/US99/15849

123

35

40

45

Asn Glu Trp Tyr Arg Gly Asp Trp Ala Trp Cys Gln Arg Cys Pro Leu
50 55 60

Val Ala Trp Pro Pro Ser Ala Glu Pro Gln Val His Arg Asn Ile His
65 70 75 80

Ser His Gly Leu Arg Ser Asn Leu Gln Glu Ile Phe Leu Pro Ala Phe
85 90 95

Pro Cys His Glu Arg Lys Lys Gln Glu
100 105

<210> 178

<211> 88

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (88)

<223> Xaa equals stop translation

<400> 178

Met Ala Asp Pro His Val Ser Phe Leu Ser Phe Arg Gln Leu Phe Ser
1 5 10 15

Trp Ala Ala Val Ile Leu Leu Arg Gly Ile Leu Gly Thr Val Ala Pro
20 25 30

Pro Pro Cys Pro Cys Val Leu Asp Leu Ala Val Tyr Pro Leu His Leu
35 40 45

Pro Val Glu Ala Pro Cys Leu Glu Val Val Phe Lys Gln Lys Asn Gly
50 55 60

Lys Asp Asn Cys Leu Val Phe Tyr Pro Asp Pro Ile Pro Leu Arg Gly
65 70 75 80

Ser Leu Leu Gly Pro Phe Ile Xaa
85

<210> 179

<211> 88

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (55)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (66)

124

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (88)

<223> Xaa equals stop translation

<400> 179

Met Ala Asp Pro His Val Ser Phe Leu Ser Phe Arg Gln Leu Phe Ser
 1 5 10 15

Trp Ala Ala Val Ile Leu Leu Arg Gly Ile Leu Gly Thr Val Ala Pro
 20 25 30

Pro Pro Cys Pro Cys Val Leu Asp Leu Ala Val Tyr Pro Leu His Leu
 35 40 45

Pro Val Glu Ala Pro Cys Xaa Glu Val Val Phe Lys Gln Lys Asn Gly
 50 55 60

Lys Xaa Asn Cys Leu Val Phe Tyr Pro Asp Pro Ile Pro Leu Arg Gly
 65 70 75 80

Ser Leu Leu Gly Pro Phe Ile Xaa
 85

<210> 180

<211> 49

<212> PRT

<213> Homo sapiens

<400> 180

Met Asn Leu Leu Gly Met Ile Phe Ser Met Cys Gly Leu Met Leu Lys
 1 5 10 15

Leu Lys Trp Cys Ala Trp Val Ala Val Tyr Cys Ser Phe Ile Ser Phe
 20 25 30

Ala Asn Ser Arg Ser Ser Glu Asp Thr Lys Gln Met Met Ser Ser Phe
 35 40 45

Met

<210> 181

<211> 23

<212> PRT

<213> Homo sapiens

<400> 181

Leu Gly Ser Leu Ser Thr Ala Pro Ser Ser Ala Leu Pro Thr Leu Gly
 1 5 10 15

Ala Arg Arg Thr Arg Ser Lys
 20

125

<210> 182
 <211> 104
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (104)
 <223> Xaa equals stop translation

<400> 182
 Met Leu Thr Phe Phe Met Ala Phe Leu Phe Asn Trp Ile Gly Phe Phe
 1 5 10 15
 Leu Ser Phe Cys Leu Thr Thr Ser Ala Ala Gly Arg Tyr Gly Ala Ile
 20 25 30
 Ser Gly Phe Gly Leu Ser Leu Ile Lys Trp Ile Leu Ile Val Arg Phe
 35 40 45
 Ser Thr Tyr Phe Pro Gly Tyr Phe Asp Gly Gln Tyr Trp Leu Trp Trp
 50 55 60
 Val Phe Leu Val Leu Gly Phe Leu Leu Phe Leu Arg Gly Phe Ile Asn
 65 70 75 80
 Tyr Ala Lys Val Arg Lys Met Pro Glu Thr Phe Ser Asn Leu Pro Arg
 85 90 95
 Thr Arg Val Leu Phe Ile Tyr Xaa
 100

<210> 183
 <211> 198
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (29)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 183
 Met Lys Lys Ser Leu Glu Asn Leu Asn Arg Leu Gln Val Met Leu Leu
 1 5 10 15
 His Leu Thr Ala Ala Phe Leu Gln Arg Ala Gln His Xaa Phe Asp Tyr
 20 25 30
 Lys Asp Glu Ser Gly Phe Pro Lys Pro Pro Ser Tyr Asn Val Ala Thr
 35 40 45
 Thr Leu Pro Ser Tyr Asp Glu Ala Glu Arg Thr Lys Ala Glu Ala Thr
 50 55 60
 Ile Pro Leu Val Pro Gly Arg Asp Glu Asp Phe Val Gly Arg Asp Asp

126

65		70		75		80									
Phe	Asp	Asp	Ala	Asp	Gln	Leu	Arg	Ile	Gly	Asn	Asp	Gly	Ile	Phe	Met
			85						90					95	
Leu	Thr	Phe	Phe	Met	Ala	Phe	Leu	Phe	Asn	Trp	Ile	Gly	Phe	Phe	Leu
			100					105					110		
Ser	Phe	Cys	Leu	Thr	Thr	Ser	Ala	Ala	Gly	Arg	Tyr	Gly	Ala	Ile	Ser
		115					120					125			
Gly	Phe	Gly	Leu	Ser	Leu	Ile	Lys	Trp	Ile	Leu	Ile	Val	Arg	Phe	Ser
	130					135					140				
Thr	Tyr	Phe	Pro	Gly	Tyr	Phe	Asp	Gly	Gln	Tyr	Trp	Leu	Trp	Trp	Val
145					150					155					160
Phe	Leu	Val	Leu	Gly	Phe	Leu	Leu	Phe	Leu	Arg	Gly	Phe	Ile	Asn	Tyr
			165						170					175	
Ala	Lys	Val	Arg	Lys	Met	Pro	Glu	Thr	Phe	Ser	Asn	Leu	Pro	Arg	Thr
			180					185					190		
Arg	Val	Leu	Phe	Ile	Tyr										
		195													

<210> 184
 <211> 70
 <212> PRT
 <213> Homo sapiens

<400> 184
 Met Leu Leu His Leu Thr Ala Ala Phe Leu Gln Arg Ala Gln Phe Ser
 1 5 10 15
 Thr Tyr Phe Pro Gly Tyr Phe Asp Gly Gln Tyr Trp Leu Trp Trp Val
 20 25 30
 Phe Leu Val Leu Gly Phe Leu Leu Phe Leu Arg Gly Phe Ile Asn Tyr
 35 40 45
 Ala Lys Val Arg Lys Met Pro Glu Thr Phe Ser Asn Leu Pro Arg Thr
 50 55 60
 Arg Val Leu Phe Ile Tyr
 65 70

<210> 185
 <211> 82
 <212> PRT
 <213> Homo sapiens

<400> 185
 Met Leu Thr Phe Phe Met Ala Phe Leu Phe Asn Trp Ile Gly Phe Phe
 1 5 10 15

127

Leu Ser Phe Cys Leu Thr Thr Ser Ala Ala Gly Arg Tyr Gly Ala Ile
 20 25 30

Ser Gly Phe Gly Leu Ser Leu Ile Lys Trp Ile Leu Ile Val Arg Phe
 35 40 45

Ser Thr Tyr Phe Pro Ala Phe Met Asn Ser Leu Ser Arg Ser Lys Arg
 50 55 60

Thr Pro Ala Gly Ser Glu Ser Arg Cys Arg Thr Gln Arg Asn Asn His
 65 70 75 80

Leu Leu

<210> 186

<211> 45

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (28)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 186

Met Lys Lys Ser Leu Glu Asn Leu Asn Arg Leu Gln Val Met Leu Leu
 1 5 10 15

His Leu Thr Ala Ala Phe Leu Gln Arg Ala His Xaa Ile Leu Thr Thr
 20 25 30

Arg Met Ser Leu Gly Phe Gln Ser Pro His Leu Thr Met
 35 40 45

<210> 187

<211> 34

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (34)

<223> Xaa equals stop translation

<400> 187

Met Thr Val Met Asp Pro Lys Gln Met Asn Val Ala Ala Val Trp
 1 5 10 15

Ala Val Val Ser Tyr Val Val Ala Asp Met Glu Glu Met Leu Pro Arg
 20 25 30

Ser Xaa

128

<210> 188
<211> 232
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (232)
<223> Xaa equals stop translation

<400> 188
Met Ala Thr Leu Trp Gly Gly Leu Leu Arg Leu Gly Ser Leu Leu Ser
1 5 10 15
Leu Ser Cys Leu Ala Leu Ser Val Leu Leu Leu Ala His Cys Gln Thr
20 25 30
Pro Pro Arg Ile Ser Arg Met Ser Asp Val Asn Val Ser Ala Leu Pro
35 40 45
Ile Lys Lys Asn Ser Gly His Ile Tyr Asn Lys Asn Ile Ser Gln Lys
50 55 60
Asp Cys Asp Cys Leu His Val Val Glu Pro Met Pro Val Arg Gly Pro
65 70 75 80
Asp Val Glu Ala Tyr Cys Leu Arg Cys Glu Cys Lys Tyr Glu Glu Arg
85 90 95
Ser Ser Val Thr Ile Lys Val Thr Ile Ile Ile Tyr Leu Ser Ile Leu
100 105 110
Gly Leu Leu Leu Leu Tyr Met Val Tyr Leu Thr Leu Val Glu Pro Ile
115 120 125
Leu Lys Arg Arg Leu Phe Gly His Ala Gln Leu Ile Gln Ser Asp Asp
130 135 140
Asp Ile Gly Asp His Gln Pro Phe Ala Asn Ala His Asp Val Leu Ala
145 150 155 160
Arg Ser Arg Ser Arg Ala Asn Val Leu Asn Lys Val Glu Tyr Gly Thr
165 170 175
Ala Ala Leu Glu Ala Ser Ser Pro Arg Ala Ala Lys Ser Leu Ser Leu
180 185 190
Thr Gly Met Leu Ser Ser Ala Asn Trp Gly Ile Glu Phe Lys Val Thr
195 200 205
Arg Lys Lys Gln Ala Asp Asn Trp Lys Gly Thr Asp Trp Val Leu Leu
210 215 220
Gly Phe Ile Leu Ile Pro Cys Xaa
225 230

<210> 189

129

<211> 457
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (457)
 <223> Xaa equals stop translation

<400> 189

Met	Ala	Ala	Ala	Gly	Arg	Leu	Pro	Ser	Ser	Trp	Ala	Leu	Phe	Ser	Pro
1				5					10					15	
Leu	Leu	Ala	Gly	Leu	Ala	Leu	Leu	Gly	Val	Gly	Pro	Val	Pro	Ala	Arg
		20						25					30		
Ala	Leu	His	Asn	Val	Thr	Ala	Glu	Leu	Phe	Gly	Ala	Glu	Ala	Trp	Gly
		35					40					45			
Thr	Leu	Ala	Ala	Phe	Gly	Asp	Leu	Asn	Ser	Asp	Lys	Gln	Thr	Asp	Leu
	50					55					60				
Phe	Val	Leu	Arg	Glu	Arg	Asn	Asp	Leu	Ile	Val	Phe	Leu	Ala	Asp	Gln
65					70					75					80
Asn	Ala	Pro	Tyr	Phe	Lys	Pro	Lys	Val	Lys	Val	Ser	Phe	Lys	Asn	His
				85					90					95	
Ser	Ala	Leu	Ile	Thr	Ser	Val	Val	Pro	Gly	Asp	Tyr	Asp	Gly	Asp	Ser
		100						105					110		
Gln	Met	Asp	Val	Leu	Leu	Thr	Tyr	Leu	Pro	Lys	Asn	Tyr	Ala	Lys	Ser
		115					120						125		
Glu	Leu	Gly	Ala	Val	Ile	Phe	Trp	Gly	Gln	Asn	Gln	Thr	Leu	Asp	Pro
	130					135						140			
Asn	Asn	Met	Thr	Ile	Leu	Asn	Arg	Thr	Phe	Gln	Asp	Glu	Pro	Leu	Ile
145					150					155					160
Met	Asp	Phe	Asn	Gly	Asp	Leu	Ile	Pro	Asp	Ile	Phe	Gly	Ile	Thr	Asn
				165					170					175	
Glu	Ser	Asn	Gln	Pro	Gln	Ile	Leu	Leu	Gly	Gly	Asn	Leu	Ser	Trp	His
			180					185					190		
Pro	Ala	Leu	Thr	Thr	Thr	Ser	Lys	Met	Arg	Ile	Pro	His	Ser	His	Ala
		195					200						205		
Phe	Ile	Asp	Leu	Thr	Glu	Asp	Phe	Thr	Ala	Asp	Leu	Phe	Leu	Thr	Thr
	210					215					220				
Leu	Asn	Ala	Thr	Thr	Ser	Thr	Phe	Gln	Phe	Glu	Ile	Trp	Glu	Asn	Leu
225					230					235					240
Asp	Gly	Asn	Phe	Ser	Val	Ser	Thr	Ile	Leu	Glu	Lys	Pro	Gln	Asn	Met
				245					250					255	

130

Met Val Val Gly Gln Ser Ala Phe Ala Asp Phe Asp Gly Asp Gly His
 260 265 270

Met Asp His Leu Leu Pro Gly Cys Glu Asp Lys Asn Cys Gln Lys Ser
 275 280 285

Thr Ile Tyr Leu Val Arg Ser Gly Met Lys Gln Trp Val Pro Val Leu
 290 295 300

Gln Asp Phe Ser Asn Lys Gly Thr Leu Trp Gly Phe Val Pro Phe Val
 305 310 315 320

Asp Glu Gln Gln Pro Thr Glu Ile Pro Ile Pro Ile Thr Leu His Ile
 325 330 335

Gly Asp Tyr Asn Met Asp Gly Tyr Pro Asp Ala Leu Val Ile Leu Lys
 340 345 350

Asn Thr Ser Gly Ser Asn Gln Gln Ala Phe Leu Leu Glu Asn Val Pro
 355 360 365

Cys Asn Asn Ala Ser Cys Glu Glu Ala Arg Arg Met Phe Lys Val Tyr
 370 375 380

Trp Glu Leu Thr Asp Leu Asn Gln Ile Lys Asp Ala Met Val Ala Thr
 385 390 395 400

Phe Phe Asp Ile Tyr Glu Asp Gly Ile Leu Asp Ile Val Val Leu Ser
 405 410 415

Lys Gly Tyr Thr Lys Asn Asp Phe Ala Ile His Thr Leu Lys Asn Asn
 420 425 430

Phe Glu Ala Asp Ala Tyr Phe Val Lys Val Ile Val Leu Ser Gly Leu
 435 440 445

Cys Ser Asn Asp Cys Pro Arg Arg Xaa
 450 455

<210> 190

<211> 185

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (185)

<223> Xaa equals stop translation

<400> 190

Met Leu Phe Leu Phe Ser Met Ala Thr Leu Leu Arg Thr Ser Phe Ser
 1 5 10 15

Asp Pro Gly Val Ile Pro Arg Ala Leu Pro Asp Glu Ala Ala Phe Ile
 20 25 30

Glu Met Glu Ile Glu Ala Thr Asn Gly Ala Val Pro Gln Gly Gln Arg

131

35 40 45
 Pro Pro Pro Arg Ile Lys Asn Phe Gln Ile Asn Asn Gln Ile Val Lys
 50 55 60
 Leu Lys Tyr Cys Tyr Thr Cys Lys Ile Phe Arg Pro Pro Arg Ala Ser
 65 70 75 80
 His Cys Ser Ile Cys Asp Asn Cys Val Glu Arg Phe Asp His His Cys
 85 90 95
 Pro Trp Val Gly Asn Cys Val Gly Lys Arg Asn Tyr Arg Tyr Phe Tyr
 100 105 110
 Leu Phe Ile Leu Ser Leu Ser Leu Leu Thr Ile Tyr Val Phe Ala Phe
 115 120 125
 Asn Ile Val Tyr Val Ala Leu Lys Ser Leu Lys Ile Gly Phe Leu Glu
 130 135 140
 Thr Leu Lys Gly Asn Ser Trp Asn Cys Ser Arg Ser Pro His Leu Leu
 145 150 155 160
 Leu Tyr Thr Leu Val Arg Arg Gly Thr Asp Trp Ile Ser Tyr Phe Pro
 165 170 175
 Arg Gly Ser Gln Pro Asp Asn Gln Xaa
 180 185

<210> 191

<211> 147

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (147)

<223> Xaa equals stop translation

<400> 191

Met Arg Val Leu Val Val Thr Ile Ala Pro Ile Tyr Trp Ala Leu Ala
 1 5 10 15
 Arg Glu Ser Gly Glu Ala Leu Asn Gly His Ser Leu Thr Gly Gly Lys
 20 25 30
 Phe Arg Gln Ser His Thr Trp Ser Leu Leu Gln Gly Ala Ala His Asp
 35 40 45
 Asp Pro Val Ala Arg Gly Leu Asp Pro Asp Gly Leu Leu Leu Asp
 50 55 60
 Val Val Val Asn Gly Val Val Pro Gly Arg Ala Trp Leu Thr Gln Ile
 65 70 75 80
 Phe Lys Cys Arg Thr Leu Lys Lys His Tyr Val Gln Thr Arg Ala Trp
 85 90 95

132

Pro Ala Val Arg Gly Leu His Thr Ala Leu Leu Pro Gly Arg Pro Pro
 100 105 110

Leu Val Pro Thr Leu Gln Pro Gln His Pro Val Gln Arg Gly Pro Gly
 115 120 125

Pro Pro Ala Pro Ala Gly Ala Ala Pro Ala Gly Leu Ser Tyr Gln Leu
 130 135 140

Gly Leu Xaa
 145

<210> 192

<211> 125

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (125)

<223> Xaa equals stop translation

<400> 192

Met Gly Glu Pro Asn Arg His Pro Ser Met Phe Leu Leu Leu Leu Val
 1 5 10 15

Leu Glu Arg Leu Tyr Ala Ser Pro Met Asp Gly Thr Ser Ser Ala Leu
 20 25 30

Ser Met Gly Pro Phe Val Pro Phe Ile Met Arg Cys Gly His Ser Pro
 35 40 45

Val Tyr His Ser Arg Glu Met Ala Ala Arg Ala Leu Val Pro Phe Val
 50 55 60

Met Ile Asp His Ile Pro Asn Thr Ile Arg Thr Leu Leu Ser Thr Leu
 65 70 75 80

Pro Ser Cys Thr Asp Gln Cys Phe Arg Ala Lys Pro His Ser Trp Gly
 85 90 95

His Phe Ser Arg Phe Phe His Leu Leu Gln Ala Tyr Ser Asp Ser Lys
 100 105 110

Thr Arg Asn Glu Phe Arg Leu Pro Ala Arg Ala Asp Xaa
 115 120 125

<210> 193

<211> 52

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (52)

133

<223> Xaa equals stop translation

<400> 193

Met Ile Lys His Val Ala Trp Leu Ile Phe Thr Asn Cys Ile Phe Phe
1 5 10 15

Cys Pro Val Ala Phe Phe Ser Phe Ala Pro Leu Ile Thr Ala Ile Ser
20 25 30

Ile Ser Pro Glu Ile Met Lys Ser Val Thr Leu Ile Phe Phe Pro Cys
35 40 45

Leu Leu Ala Xaa
50

<210> 194

<211> 320

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (68)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (115)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (213)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (320)

<223> Xaa equals stop translation

<400> 194

Met Ala Pro Leu Ala Leu His Leu Leu Val Leu Val Pro Ile Leu Leu
1 5 10 15

Ser Leu Val Ala Ser Gln Asp Trp Lys Ala Glu Arg Ser Gln Asp Pro
20 25 30

Phe Glu Lys Cys Met Gln Asp Pro Asp Tyr Glu Gln Leu Leu Lys Val
35 40 45

Thr Ile Leu Glu Ala Asp Asn Arg Ile Gly Gly Arg Ile Phe Thr Tyr
50 55 60

Arg Asp Gln Xaa Thr Gly Trp Ile Gly Glu Leu Gly Ala Met Arg Met
65 70 75 80

Pro Ser Ser His Arg Ile Leu His Lys Leu Cys Gln Gly Leu Gly Leu

134
 85 90 95
 Asn Leu Thr Lys Phe Thr Gln Tyr Asp Lys Asn Thr Trp Thr Glu Val
 100 105 110
 His Glu Xaa Lys Leu Arg Asn Tyr Val Val Glu Lys Val Pro Glu Lys
 115 120 125
 Leu Gly Tyr Ala Leu Arg Pro Gln Glu Lys Gly His Ser Pro Glu Asp
 130 135 140
 Ile Tyr Gln Met Ala Leu Asn Gln Ala Leu Lys Asp Leu Lys Ala Leu
 145 150 155 160
 Gly Cys Arg Lys Ala Met Lys Lys Phe Glu Arg His Thr Leu Leu Glu
 165 170 175
 Tyr Leu Leu Gly Glu Gly Asn Leu Ser Arg Pro Ala Val Gln Leu Leu
 180 185 190
 Gly Asp Val Met Ser Glu Asp Gly Phe Phe Tyr Leu Ser Phe Ala Glu
 195 200 205
 Ala Leu Arg Ala Xaa Ser Cys Leu Ser Asp Arg Leu Gln Tyr Ser Arg
 210 215 220
 Ile Val Gly Gly Trp Asp Leu Leu Pro Arg Ala Leu Leu Ser Ser Leu
 225 230 235 240
 Ser Gly Leu Val Leu Leu Asn Ala Pro Val Val Ala Met Thr Gln Gly
 245 250 255
 Pro His Asp Val His Val Gln Ile Glu Thr Ser Pro Pro Ala Arg Asn
 260 265 270
 Leu Lys Val Leu Lys Ala Asp Val Val Leu Leu Thr Ala Ser Gly Pro
 275 280 285
 Ala Val Lys Arg Ile Thr Phe Ser Pro Arg Cys Pro Ala Thr Cys Arg
 290 295 300
 Arg Arg Cys Gly Gly Cys Thr Thr Cys Arg Pro Pro Arg Cys Ser Xaa
 305 310 315 320

<210> 195

<211> 130

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (38)

<223> Xaa equals any of the naturally occurring L-amino acids

135

<220>

<221> SITE

<222> (53)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 195

Pro Phe Cys Ser Gly Phe Phe Pro Ser Leu Trp Ile Tyr Leu Pro Phe
1 5 10 15

Ile Phe Asn Val Ser Asp Leu Trp Met Gly Ser Leu Ser Gly Cys Ala
20 25 30

Leu Pro Phe Cys Leu Xaa Val Phe Phe Leu Thr Val Ser Pro Ser Ala
35 40 45

Val Gly Leu Leu Xaa Phe Ala Gly Gly Pro Leu Gln Thr Leu Phe Ala
50 55 60

Trp Val Ser Pro Val Glu Ala Ala Glu Gln Gln Arg Leu Leu Pro Val
65 70 75 80

Leu Ser Ser Gly Ser Phe Val Ser Glu Gly Thr Cys Gln Met Pro Ala
85 90 95

Arg Ala Leu Leu Tyr Glu Val Ser Val Gly Pro Tyr Trp Glu Ile Pro
100 105 110

Pro Ser Gln Asp Thr Arg Arg Ser Gly Thr Tyr Leu Arg Arg Gln Ser
115 120 125

Asp Pro
130

<210> 196

<211> 108

<212> PRT

<213> Homo sapiens

<400> 196

His Glu Gly Ser Cys Arg Ala Pro Gly Phe Ser Ala His Lys Gly Arg
1 5 10 15

Gly Cys Pro Ser Pro Arg Met Thr Leu Pro Ser Arg Ala Leu Ala Ser
20 25 30

Leu Gly Val Gly Val Trp Gly Met Leu Arg Leu Asn Gln Val Thr Val
35 40 45

Ser Cys Gly Gly Ser Arg Trp Ser Ser Arg Val Ala Leu Gly Ala Phe
50 55 60

Ser Trp Val Cys Gly Val Ala Leu Val Leu Gln Pro Ser Gly Gly Gly
65 70 75 80

Leu Gly Leu Thr Ser Pro Ser Glu Gly Cys Trp Glu Gly Glu Leu Ala
85 90 95

136

Leu Ala Val Leu Arg Ala Pro Gly Gly Ser Pro Ser
 100 105

<210> 197
 <211> 104
 <212> PRT
 <213> Homo sapiens

<400> 197
 Ile Pro Leu Thr Leu Pro Gly Ile Phe Leu Leu Ile Arg Leu Phe Trp
 1 5 10 15
 Arg Leu Gly Gln Ser Ile Cys Gly Pro Gly Lys Leu Val Leu Trp Pro
 20 25 30
 Gln Phe Cys Cys Gly Cys Ala Val Ile Ser Gly His Cys Val Pro Arg
 35 40 45
 Gly Met Pro Ser Ser Trp Leu Pro Gly Cys Phe Val Leu Leu Cys Leu
 50 55 60
 Val Ala Val Gly Cys Gln Leu Arg Glu Trp Gly Val Gly Gly Val Ser
 65 70 75 80
 Ala Val Gly Leu Leu Ala Leu Pro His Leu Gln Val Leu Gly Met Arg
 85 90 95
 Gly Arg Gly Leu Ile Ser Gly Gly
 100

<210> 198
 <211> 237
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (142)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 198
 Gly Pro Ala Gly Lys Glu Ala Trp Ile Trp Ser Trp Leu Leu Pro Ser
 1 5 10 15
 Pro Gly Pro Ala Pro Leu Pro Ser Ala Ser Trp Gly Leu Cys Gly Asp
 20 25 30
 Ala Pro Arg Ala Ala Ala Arg Gly Pro Val Glu Pro Gly Ala Ala Arg
 35 40 45
 Met Ala Leu Leu Ser Arg Pro Ala Leu Thr Leu Leu Leu Leu Met
 50 55 60
 Ala Ala Val Val Arg Cys Gln Glu Gln Ala Gln Thr Thr Asp Trp Arg
 65 70 75 80

137

Ala Thr Leu Lys Thr Ile Arg Asn Gly Val His Lys Ile Asp Thr Tyr
85 90 95

Leu Asn Ala Ala Leu Asp Leu Leu Gly Gly Glu Asp Gly Leu Cys Gln
100 105 110

Tyr Lys Cys Ser Asp Gly Ser Lys Pro Phe Pro Arg Tyr Gly Tyr Lys
115 120 125

Pro Ser Pro Pro Asn Gly Cys Gly Ser Pro Leu Phe Gly Xaa His Leu
130 135 140

Asn Ile Gly Ile Pro Ser Leu Thr Lys Cys Cys Asn Gln His Asp Arg
145 150 155 160

Cys Tyr Glu Thr Cys Gly Lys Ser Lys Asn Asp Cys Asp Glu Glu Phe
165 170 175

Gln Tyr Cys Leu Ser Lys Ile Cys Arg Asp Val Gln Lys Thr Leu Gly
180 185 190

Leu Thr Gln His Val Gln Ala Cys Glu Thr Thr Val Glu Leu Leu Phe
195 200 205

Asp Ser Val Ile His Leu Gly Cys Lys Pro Tyr Leu Asp Ser Gln Arg
210 215 220

Ala Ala Cys Arg Cys His Tyr Glu Glu Lys Thr Asp Leu
225 230 235

<210> 199

<211> 8

<212> PRT

<213> Homo sapiens

<400> 199

Cys Cys Asn Gln His Asp Arg Cys

1

5

<210> 200

<211> 15

<212> PRT

<213> Homo sapiens

<400> 200

Ser Leu Thr Lys Cys Cys Asn Gln His Asp Arg Cys Tyr Glu Thr

1

5

10

15

<210> 201

<211> 16

<212> PRT

<213> Homo sapiens

<400> 201

Leu Thr Lys Cys Cys Asn Gln His Asp Arg Cys Tyr Glu Thr Cys Gly

138

1

5

10

15

<210> 202
 <211> 260
 <212> PRT
 <213> Homo sapiens

<400> 202

Gly Thr Ser Ser Ala Arg Pro Arg Gly Ala Leu Pro Gly Gly Ser Ala
 1 5 10 15

Pro Ser Ala Pro His Gly Gln Leu Pro Gly Arg Ala Gln Pro Ala Pro
 20 25 30

Val Ser Gly Pro Pro Pro Thr Ser Gly Leu Cys His Phe Asp Pro Ala
 35 40 45

Ala Pro Trp Pro Leu Trp Pro Gly Pro Trp Gln Leu Pro Pro His Pro
 50 55 60

Gln Asp Trp Pro Ala His Pro Asp Ile Pro Gln Asp Trp Val Ser Phe
 65 70 75 80

Leu Arg Ser Phe Gly Gln Leu Thr Leu Cys Pro Arg Asn Gly Thr Val
 85 90 95

Thr Gly Lys Trp Arg Gly Ser His Val Val Gly Leu Leu Thr Thr Leu
 100 105 110

Asn Phe Gly Asp Gly Pro Asp Arg Asn Lys Thr Arg Thr Phe Gln Ala
 115 120 125

Thr Val Leu Gly Ser Gln Met Gly Leu Lys Gly Ser Ser Ala Gly Gln
 130 135 140

Leu Val Leu Ile Thr Ala Arg Val Thr Thr Glu Arg Thr Ala Gly Thr
 145 150 155 160

Cys Leu Tyr Phe Ser Ala Val Pro Gly Ile Leu Pro Ser Ser Gln Pro
 165 170 175

Pro Ile Ser Cys Ser Glu Glu Gly Ala Gly Asn Ala Thr Leu Ser Pro
 180 185 190

Arg Met Gly Glu Glu Cys Val Ser Val Trp Ser His Glu Gly Leu Val
 195 200 205

Leu Thr Lys Leu Leu Thr Ser Glu Glu Leu Ala Leu Cys Gly Ser Arg
 210 215 220

Leu Leu Val Leu Gly Ser Phe Leu Leu Leu Phe Cys Gly Leu Leu Cys
 225 230 235 240

Cys Val Thr Ala Met Cys Phe His Pro Arg Arg Glu Ser His Trp Ser

139
245 250 255
Arg Thr Arg Leu
260

<210> 203
<211> 80
<212> PRT
<213> Homo sapiens

<400> 203
Ala Arg Ala Pro Pro Gly Pro Glu Gly Leu Ser Pro Glu Ala Gln Pro
1 5 10 15
Pro Leu Leu Pro Met Gly Asn Cys Gln Ala Gly His Asn Leu His Leu
20 25 30
Cys Leu Ala His His Pro Pro Leu Val Cys Ala Thr Leu Ile Leu Leu
35 40 45
Leu Leu Gly Leu Ser Gly Leu Gly Leu Gly Ser Phe Leu Leu Thr His
50 55 60
Arg Thr Gly Leu Arg Thr Leu Thr Ser Pro Arg Thr Gly Ser Leu Phe
65 70 75 80

<210> 204
<211> 224
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (6)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (9)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (22)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (143)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE

140

<222> (186)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 204

```

Arg Phe Leu Ser Val Xaa Pro Gln Xaa Glu Val Pro Phe Leu Leu His
  1           5           10           15

Pro Cys Val Cys Phe Xaa Gly Gly His Pro Ser Leu Leu Pro Asp Pro
          20           25           30

Cys Arg Ala Val Gly Gly Gly Trp Glu Ala Pro Arg Cys Cys Leu His
          35           40           45

Glu Ala Leu Cys Gln Ser Leu Gly Cys Lys Ala Glu Glu Ile Val Ser
  50           55           60

Val Ser Glu Ser Ser Ser Ala Gln Arg Cys Trp Tyr Leu Leu Arg Gly
  65           70           75           80

Arg Lys Ala Gly Gly Arg Gly Pro Ala Ser Pro Val Leu Phe Ala Leu
          85           90           95

Met Arg Leu Glu Ser Leu Cys His Leu Cys Leu Ala Cys Leu Phe Phe
          100          105          110

Arg Leu Pro Ala Thr Arg Thr Val Tyr Cys Met Asn Glu Ala Glu Ile
          115          120          125

Val Asp Val Ala Leu Gly Ile Leu Ile Glu Ser Arg Lys Gln Xaa Lys
          130          135          140

Ala Cys Glu Gln Pro Ala Leu Ala Gly Ala Asp Asn Pro Glu His Ser
          145          150          155          160

Pro Pro Cys Ser Val Ser Pro His Thr Ser Ser Gly Ser Ser Ser Glu
          165          170          175

Glu Glu Asp Ser Gly Lys Gln Ala Leu Xaa Pro Gly Leu Ser Pro Ser
          180          185          190

Gln Arg Pro Gly Gly Ser Ser Ser Ala Cys Ser Arg Ser Pro Glu Glu
          195          200          205

Glu Glu Glu Glu Asp Val Leu Lys Tyr Val Arg Glu Ile Phe Phe Ser
          210          215          220

```

<210> 205

<211> 199

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (35)

141

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (103)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (191)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 205

Val	Pro	Gly	Trp	Pro	Arg	Ala	Cys	Ser	Pro	Cys	Gln	Ala	Asp	Ser	Pro
1				5					10					15	

Arg	Ala	His	Pro	Pro	Lys	Leu	Arg	Gly	Ile	Leu	Arg	Trp	Ala	Pro	Val
			20					25					30		

Pro	Leu	Xaa	Cys	Ala	Ala	Leu	Cys	Pro	Pro	Leu	Asp	Ser	Gly	Met	Ser
	35						40						45		

Met	Ala	Ala	Cys	Pro	Glu	Ala	Pro	Glu	Pro	Ser	Phe	Leu	Arg	Glu	Val
	50					55					60				

Pro	Ser	Ser	Pro	Ala	Ser	Thr	Gln	Trp	His	Arg	Pro	Cys	Asn	Phe	Arg
	65				70					75					80

Gln	Val	Glu	Ala	Asn	Pro	Arg	Lys	Glu	Pro	Lys	Asn	Leu	Val	Trp	Arg
				85					90					95	

Asp	Val	Ser	Leu	Gly	Gln	Xaa	Ser	Arg	Thr	Pro	Arg	Gly	Ser	Gly	Leu
			100					105					110		

Glu	Leu	Val	Arg	Val	Cys	Gly	Gly	Gly	Met	Gln	Arg	Asp	Lys	Thr	Val
		115					120					125			

Val	Glu	Glu	Arg	Val	Gly	Glu	Glu	Arg	Glu	Arg	Glu	Arg	Glu	Arg	Glu
		130				135					140				

Ser	Leu	Gly	Gly	Ala	Gly	Lys	His	Gly	Glu	Met	Arg	Cys	Val	Tyr	Val
	145				150					155					160

Arg	Glu	Ser	Val	Gly	Ala	Pro	Gly	Arg	Ala	Gly	Gly	Gly	Gly	Asn	Gly
				165					170					175	

Val	Asn	Ser	Val	Gly	Cys	Val	Arg	Thr	Val	His	Ser	Gly	Ser	Xaa	Pro
			180					185					190		

Pro	Pro	Ser	Ala	Gly	Val	Ser
						195

<210> 206

<211> 174

<212> PRT

<213> Homo sapiens

142

<400> 206

```

Thr Arg Pro Gly Lys Glu Leu Asn Leu Val Phe Gly Leu Gln Leu Ser
  1               5               10               15

Met Ala Arg Ile Gly Ser Thr Val Asn Met Asn Leu Met Gly Trp Leu
      20               25               30

Tyr Ser Lys Ile Glu Ala Leu Leu Gly Ser Ala Gly His Thr Thr Leu
      35               40               45

Gly Ile Thr Leu Met Ile Gly Gly Ile Thr Cys Ile Leu Ser Leu Ile
      50               55               60

Cys Ala Leu Ala Leu Ala Tyr Leu Asp Gln Arg Ala Glu Arg Ile Leu
      65               70               75               80

His Lys Glu Gln Gly Lys Thr Gly Glu Val Ile Lys Leu Thr Asp Val
      85               90               95

Lys Asp Phe Ser Leu Pro Leu Trp Leu Ile Phe Ile Ile Cys Val Cys
      100               105               110

Tyr Tyr Val Ala Val Phe Pro Phe Ile Gly Leu Gly Lys Val Phe Phe
      115               120               125

Thr Glu Lys Phe Gly Phe Ser Ser Gln Ala Ala Ser Ala Ile Asn Ser
      130               135               140

Val Val Tyr Val Ile Ser Ala Pro Met Ser Pro Val Phe Gly Leu Leu
      145               150               155               160

Val Asp Lys Thr Gly Lys Asn Ile Ile Trp Val Leu Cys Ala
      165               170

```

<210> 207

<211> 31

<212> PRT

<213> Homo sapiens

<400> 207

```

Cys Lys Asp Leu Cys Ser Arg Val Tyr Leu Leu Thr Leu Ser Pro Leu
  1               5               10               15

Leu Ser Tyr Asp Pro Ala Thr Ser His Ser Pro Arg Asn Thr Gln
      20               25               30

```

<210> 208

<211> 369

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (78)

<223> Xaa equals any of the naturally occurring L-amino acids

143

<400> 208

Ile Ile Cys Glu Cys Trp Glu Glu Glu Cys Gln Ser Cys Arg Leu Lys
 1 5 10 15
 Ile Thr Gln Pro Arg Glu Ile Cys Arg Met Asp Phe Leu Val Leu Phe
 20 25 30
 Leu Phe Tyr Leu Ala Ser Val Leu Met Gly Leu Val Leu Ile Cys Val
 35 40 45
 Cys Ser Lys Thr His Ser Leu Lys Gly Leu Ala Arg Gly Gly Ala Gln
 50 55 60
 Ile Phe Ser Cys Ile Ile Pro Glu Cys Leu Gln Arg Ala Xaa His Gly
 65 70 75 80
 Leu Leu His Tyr Leu Phe His Thr Arg Asn His Thr Phe Ile Val Leu
 85 90 95
 His Leu Val Leu Gln Gly Met Val Tyr Thr Glu Tyr Thr Trp Glu Val
 100 105 110
 Phe Gly Tyr Cys Gln Glu Leu Glu Leu Ser Leu His Tyr Leu Leu Leu
 115 120 125
 Pro Tyr Leu Leu Leu Gly Val Asn Leu Phe Phe Phe Thr Leu Thr Cys
 130 135 140
 Gly Thr Asn Pro Gly Ile Ile Thr Lys Ala Asn Glu Leu Leu Phe Leu
 145 150 155 160
 His Val Tyr Glu Phe Asp Glu Val Met Phe Pro Lys Asn Val Arg Cys
 165 170 175
 Ser Thr Cys Asp Leu Arg Lys Pro Ala Arg Ser Lys His Cys Ser Val
 180 185 190
 Cys Asn Trp Cys Val His Arg Phe Asp His His Cys Val Trp Val Asn
 195 200 205
 Asn Cys Ile Gly Ala Trp Asn Ile Arg Tyr Phe Leu Ile Tyr Val Leu
 210 215 220
 Thr Leu Thr Ala Ser Ala Ala Thr Val Ala Ile Val Ser Thr Thr Phe
 225 230 235 240
 Leu Val His Leu Val Val Met Ser Asp Leu Tyr Gln Glu Thr Tyr Ile
 245 250 255
 Asp Asp Leu Gly His Leu His Val Met Asp Thr Val Phe Leu Ile Gln
 260 265 270
 Tyr Leu Phe Leu Thr Phe Pro Arg Ile Val Phe Met Leu Gly Phe Val
 275 280 285
 Val Val Leu Ser Phe Leu Leu Gly Gly Tyr Leu Leu Phe Val Leu Tyr
 290 295 300

144

Leu Ala Ala Thr Asn Gln Thr Thr Asn Glu Trp Tyr Arg Gly Asp Trp
 305 310 315 320

Ala Trp Cys Gln Arg Cys Pro Leu Val Ala Trp Pro Pro Ser Ala Glu
 325 330 335

Pro Gln Val His Arg Asn Ile His Ser His Gly Leu Arg Ser Asn Leu
 340 345 350

Gln Glu Ile Phe Leu Pro Ala Phe Pro Cys His Glu Arg Lys Lys Gln
 355 360 365

Glu

<210> 209

<211> 147

<212> PRT

<213> Homo sapiens

<400> 209

Leu Leu Ser Phe Lys Ile Arg Gly Leu Arg Thr Glu Asp Ala Gly Trp
 1 5 10 15

Ala Gln Ser Ser Ser Gly Gly Leu Cys Val Arg Gly Asp Ala Phe Trp
 20 25 30

Met Pro Ser Ser Ser Ser Gly Leu Gly Ser Pro Ser Arg Pro Pro Ser
 35 40 45

Ser Phe Leu Cys Leu Leu Leu Leu Leu Leu Pro Pro Ala Ala Leu Ala
 50 55 60

Leu Leu Leu Phe Phe Leu Asp Phe Phe Pro Pro Arg Ala Ala Val Ser
 65 70 75 80

Pro Phe Leu Pro Asp His Cys Ser Ala Arg Gln Pro Arg Val Trp Arg
 85 90 95

Arg Glu Thr Leu Asn Arg Ser Ala Ser Gly Leu Gly Cys Trp Ala Arg
 100 105 110

Ser Thr Glu Gln Gly Ala Val Gly Val Ala Thr Gly Thr Val Leu Asp
 115 120 125

Ile Ser Leu Pro Ala Ser Cys Leu Ser Leu Trp Pro Pro Gly Pro Ser
 130 135 140

Gly Gly Ile
 145

<210> 210

<211> 143

<212> PRT

<213> Homo sapiens

145

<400> 210

Gln Leu Gly Leu Cys Leu Thr Ser Ala Ser Leu Pro Pro Ala Ser Arg
 1 5 10 15
 Cys Gly His Gln Ala Pro Leu Gly Ala Ser Asp Leu Ser Ala His His
 20 25 30
 Ser Ala Pro Gly Phe Ser Asp Ser Tyr Phe Thr Met Ser Cys Gln Ser
 35 40 45
 Ser Leu Ser Arg Ala Glu Ile Leu Gln Cys Pro Leu Val Pro Ser Val
 50 55 60
 Ser Pro Pro Thr His Leu Pro Gln Gly Arg Ala Asn Lys Ser Ser Arg
 65 70 75 80
 Ala Ser Leu Pro Leu Leu Pro Gln Thr His Trp Cys Leu Phe Pro Ser
 85 90 95
 Ala Arg Gly Trp Arg Arg Gly Ile Gln Ser Gly Leu Pro Pro Gly Gly
 100 105 110
 Ser Cys Thr Ser Pro Arg Ser Pro Pro Gln Thr Leu His Gln His Ile
 115 120 125
 Thr Leu Val Asn His Asn Thr Ser Tyr Trp Gln Ser Pro Ser Thr
 130 135 140

<210> 211

<211> 160

<212> PRT

<213> Homo sapiens

<400> 211

His Gln Pro Pro Cys Leu Leu Pro Leu Ala Val Ala Thr Arg Pro Leu
 1 5 10 15
 Trp Gly His Leu Thr Cys Leu Pro Ile Ile Leu His Leu Val Ser Val
 20 25 30
 Thr Leu Thr Ser Pro Cys Leu Ala Asn Gln Ala Phe Gln Gly Gln Arg
 35 40 45
 Ser Tyr Asn Ala Leu Trp Cys Pro Leu Phe Leu Leu Leu Pro Thr Ser
 50 55 60
 Pro Lys Gly Glu Gln Thr Asn His Pro Glu Pro Ala Cys Pro Cys Phe
 65 70 75 80
 Pro Lys Leu Thr Gly Val Phe Ser Leu Gln His Val Val Gly Ala Glu
 85 90 95
 Glu Phe Ser Gln Val Phe Leu Leu Val Asp Pro Val Pro Val Leu Asp
 100 105 110
 His Leu Leu Lys Leu Phe Thr Ser Thr Ser His Leu Leu Ile Ile Ile
 115 120 125

146

Pro His Ile Gly Lys Ala Pro Ala Pro Asp Ser Leu Leu Glu Glu Leu
 130 135 140

Ser Leu Ser Leu Ala Thr His Cys Lys Val Ala Val Ala Arg Phe Thr
 145 150 155 160

<210> 212
 <211> 157
 <212> PRT
 <213> Homo sapiens

<400> 212
 Met Ala Ala Glu Gly Ser Arg Phe Ser Ser Gln Ser Pro Gly Leu Val
 1 5 10 15
 Asp Arg Gln Gly Pro Lys Cys Asp Pro Ser Arg Leu Val Ser Pro Trp
 20 25 30
 Gly Arg His Gly Leu Arg Ile Leu Gln Ile Gly His His His Gly Arg
 35 40 45
 Asp Gly Gln His Glu Ala Thr His His Leu Leu Arg Val Leu Arg Ala
 50 55 60
 Pro Arg Val Gly Lys Ala Asp Glu Gly Ala Val Asp Ser Asp Pro Ser
 65 70 75 80
 Thr Pro Leu Gln Leu Lys His Glu Ala Ala His Ala Glu Asp His Ala
 85 90 95
 Gln Gln Val His Val Val Arg Arg Arg Val Val Gln Gly Arg Val Thr
 100 105 110
 Phe Ala Arg Arg Gly Leu Val Pro Gln His Phe Val Arg Pro Pro Trp
 115 120 125
 Val Arg His Ile Val Ser Gly His Ser Glu Ser Lys Ala Arg Ser Arg
 130 135 140
 Leu Phe Arg Cys Arg Asn Arg Ser Phe Arg Arg Ala Ser
 145 150 155

<210> 213
 <211> 38
 <212> PRT
 <213> Homo sapiens

<400> 213
 Arg Leu Val Ser Pro Trp Gly Arg His Gly Leu Arg Ile Leu Gln Ile
 1 5 10 15
 Gly His His His Gly Arg Asp Gly Gln His Glu Ala Thr His His Leu

147

20

25

30

Leu Arg Val Leu Arg Ala
35

<210> 214
<211> 12
<212> PRT
<213> Homo sapiens

<400> 214
Pro Thr Asp Val Leu Lys Ile Arg Met Gln Ala Gln
1 5 10

<210> 215
<211> 7
<212> PRT
<213> Homo sapiens

<400> 215
Thr Tyr Glu Gln Leu Lys Arg
1 5

<210> 216
<211> 137
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (22)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (33)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (71)
<223> Xaa equals any of the naturally occurring L-amino acids

<400> 216
Arg Pro Arg Pro Ser Ala Ser Ser Leu Ala Arg Ser Ala Ser Leu Leu
1 5 10 15

Pro Ala Ala His Gly Xaa Gly Val Gly Gly Ala Gly Gly Gly Ser Ser
20 25 30

Xaa Leu Arg Ser Arg Tyr Gln Gln Leu Gln Asn Glu Glu Glu Ser Gly
35 40 45

Glu Pro Glu Gln Ala Ala Gly Asp Ala Pro Pro Pro Tyr Ser Ser Ile
50 55 60

148

Ser Ala Glu Ser Ala His Xaa Phe Asp Tyr Lys Asp Glu Ser Gly Phe
 65 70 75 80

Pro Lys Pro Pro Ser Tyr Asn Val Ala Thr Thr Leu Pro Ser Tyr Asp
 85 90 95

Glu Ala Glu Arg Thr Lys Ala Glu Ala Thr Ile Pro Leu Val Pro Gly
 100 105 110

Arg Asp Glu Asp Phe Val Gly Arg Asp Asp Phe Asp Asp Ala Asp Gln
 115 120 125

Leu Arg Ile Gly Asn Asp Gly Ile Phe
 130 135

<210> 217
 <211> 20
 <212> PRT
 <213> Homo sapiens

<400> 217
 Arg Tyr Gln Gln Leu Gln Asn Glu Glu Glu Ser Gly Glu Pro Glu Gln
 1 5 10 15

Ala Ala Gly Asp
 20

<210> 218
 <211> 22
 <212> PRT
 <213> Homo sapiens

<400> 218
 Pro Gly Arg Asp Glu Asp Phe Val Gly Arg Asp Asp Phe Asp Ala
 1 5 10 15

Asp Gln Leu Arg Ile Gly
 20

<210> 219
 <211> 103
 <212> PRT
 <213> Homo sapiens

<400> 219
 Met Leu Thr Phe Phe Met Ala Phe Leu Phe Asn Trp Ile Gly Phe Phe
 1 5 10 15

Leu Ser Phe Cys Leu Thr Thr Ser Ala Ala Gly Arg Tyr Gly Ala Ile
 20 25 30

Ser Gly Phe Gly Leu Ser Leu Ile Lys Trp Ile Leu Ile Val Arg Phe
 35 40 45

149

Ser Thr Tyr Phe Pro Gly Tyr Phe Asp Gly Gln Tyr Trp Leu Trp Trp
 50 55 60
 Val Phe Leu Val Leu Gly Phe Leu Leu Phe Leu Arg Gly Phe Ile Asn
 65 70 75 80
 Tyr Ala Lys Val Arg Lys Met Pro Glu Thr Phe Ser Asn Leu Pro Arg
 85 90 95
 Thr Arg Val Leu Phe Ile Tyr
 100

<210> 220

<211> 198

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (29)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 220

Met Lys Lys Ser Leu Glu Asn Leu Asn Arg Leu Gln Val Met Leu Leu
 1 5 10 15
 His Leu Thr Ala Ala Phe Leu Gln Arg Ala Gln His Xaa Phe Asp Tyr
 20 25 30
 Lys Asp Glu Ser Gly Phe Pro Lys Pro Pro Ser Tyr Asn Val Ala Thr
 35 40 45
 Thr Leu Pro Ser Tyr Asp Glu Ala Glu Arg Thr Lys Ala Glu Ala Thr
 50 55 60
 Ile Pro Leu Val Pro Gly Arg Asp Glu Asp Phe Val Gly Arg Asp Asp
 65 70 75 80
 Phe Asp Asp Ala Asp Gln Leu Arg Ile Gly Asn Asp Gly Ile Phe Met
 85 90 95
 Leu Thr Phe Phe Met Ala Phe Leu Phe Asn Trp Ile Gly Phe Phe Leu
 100 105 110
 Ser Phe Cys Leu Thr Thr Ser Ala Ala Gly Arg Tyr Gly Ala Ile Ser
 115 120 125
 Gly Phe Gly Leu Ser Leu Ile Lys Trp Ile Leu Ile Val Arg Phe Ser
 130 135 140
 Thr Tyr Phe Pro Gly Tyr Phe Asp Gly Gln Tyr Trp Leu Trp Trp Val
 145 150 155 160
 Phe Leu Val Leu Gly Phe Leu Leu Phe Leu Arg Gly Phe Ile Asn Tyr
 165 170 175
 Ala Lys Val Arg Lys Met Pro Glu Thr Phe Ser Asn Leu Pro Arg Thr

150
180 185 190
Arg Val Leu Phe Ile Tyr
195

<210> 221
<211> 70
<212> PRT
<213> Homo sapiens

<400> 221
Met Leu Leu His Leu Thr Ala Ala Phe Leu Gln Arg Ala Gln Phe Ser
1 5 10 15
Thr Tyr Phe Pro Gly Tyr Phe Asp Gly Gln Tyr Trp Leu Trp Trp Val
20 25 30
Phe Leu Val Leu Gly Phe Leu Leu Phe Leu Arg Gly Phe Ile Asn Tyr
35 40 45
Ala Lys Val Arg Lys Met Pro Glu Thr Phe Ser Asn Leu Pro Arg Thr
50 55 60
Arg Val Leu Phe Ile Tyr
65 70

<210> 222
<211> 82
<212> PRT
<213> Homo sapiens

<400> 222
Met Leu Thr Phe Phe Met Ala Phe Leu Phe Asn Trp Ile Gly Phe Phe
1 5 10 15
Leu Ser Phe Cys Leu Thr Thr Ser Ala Ala Gly Arg Tyr Gly Ala Ile
20 25 30
Ser Gly Phe Gly Leu Ser Leu Ile Lys Trp Ile Leu Ile Val Arg Phe
35 40 45
Ser Thr Tyr Phe Pro Ala Phe Met Asn Ser Leu Ser Arg Ser Lys Arg
50 55 60
Thr Pro Ala Gly Ser Glu Ser Arg Cys Arg Thr Gln Arg Asn Asn His
65 70 75 80
Leu Leu

<210> 223
<211> 45
<212> PRT
<213> Homo sapiens

151

<220>

<221> SITE

<222> (28)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 223

Met	Lys	Lys	Ser	Leu	Glu	Asn	Leu	Asn	Arg	Leu	Gln	Val	Met	Leu	Leu
1				5					10					15	

His	Leu	Thr	Ala	Ala	Phe	Leu	Gln	Arg	Ala	His	Xaa	Ile	Leu	Thr	Thr
			20					25					30		

Arg	Met	Ser	Leu	Gly	Phe	Gln	Ser	Pro	His	Leu	Thr	Met
		35					40					45

<210> 224

<211> 33

<212> PRT

<213> Homo sapiens

<400> 224

Met	Thr	Val	Met	Asp	Pro	Lys	Gln	Met	Asn	Val	Ala	Ala	Ala	Val	Trp
1				5					10					15	

Ala	Val	Val	Ser	Tyr	Val	Val	Ala	Asp	Met	Glu	Glu	Met	Leu	Pro	Arg
			20					25					30		

Ser

<210> 225

<211> 189

<212> PRT

<213> Homo sapiens

<400> 225

Pro	Arg	Val	Arg	Ser	Arg	Glu	Pro	Val	Ala	Gly	Ala	Pro	Gly	Cys	Gly
1				5					10					15	

Thr	Ala	Gly	Pro	Pro	Ala	Met	Ala	Thr	Leu	Trp	Gly	Gly	Leu	Leu	Arg
			20					25					30		

Leu	Gly	Ser	Leu	Leu	Ser	Leu	Ser	Cys	Leu	Ala	Leu	Ser	Val	Leu	Leu
		35					40					45			

Leu	Ala	His	Cys	Gln	Thr	Pro	Pro	Ser	Asp	Cys	Leu	His	Val	Val	Glu
	50						55				60				

Pro	Met	Pro	Val	Arg	Gly	Pro	Asp	Val	Glu	Ala	Tyr	Cys	Leu	Arg	Cys
	65				70					75				80	

Glu	Cys	Lys	Tyr	Glu	Glu	Arg	Ser	Ser	Val	Thr	Ile	Lys	Val	Thr	Ile
			85						90					95	

Ile	Ile	Tyr	Leu	Ser	Ile	Leu	Gly	Leu	Leu	Leu	Tyr	Met	Val	Tyr
			100					105						110

152

Leu Thr Leu Val Glu Pro Ile Leu Lys Arg Arg Leu Phe Gly His Ala
 115 120 125

Gln Leu Ile Gln Ser Asp Asp Asp Ile Gly Asp His Gln Pro Phe Ala
 130 135 140

Asn Ala His Asp Val Leu Ala Arg Ser Arg Ser Arg Ala Asn Val Leu
 145 150 155 160

Asn Lys Val Glu Tyr Ala Gln Gln Arg Trp Lys Leu Gln Val Gln Glu
 165 170 175

Gln Arg Lys Ser Val Phe Asp Arg His Val Val Leu Ser
 180 185

<210> 226

<211> 231

<212> PRT

<213> Homo sapiens

<400> 226

Met Ala Thr Leu Trp Gly Gly Leu Leu Arg Leu Gly Ser Leu Leu Ser
 1 5 10 15

Leu Ser Cys Leu Ala Leu Ser Val Leu Leu Leu Ala His Cys Gln Thr
 20 25 30

Pro Pro Arg Ile Ser Arg Met Ser Asp Val Asn Val Ser Ala Leu Pro
 35 40 45

Ile Lys Lys Asn Ser Gly His Ile Tyr Asn Lys Asn Ile Ser Gln Lys
 50 55 60

Asp Cys Asp Cys Leu His Val Val Glu Pro Met Pro Val Arg Gly Pro
 65 70 75 80

Asp Val Glu Ala Tyr Cys Leu Arg Cys Glu Cys Lys Tyr Glu Glu Arg
 85 90 95

Ser Ser Val Thr Ile Lys Val Thr Ile Ile Ile Tyr Leu Ser Ile Leu
 100 105 110

Gly Leu Leu Leu Leu Tyr Met Val Tyr Leu Thr Leu Val Glu Pro Ile
 115 120 125

Leu Lys Arg Arg Leu Phe Gly His Ala Gln Leu Ile Gln Ser Asp Asp
 130 135 140

Asp Ile Gly Asp His Gln Pro Phe Ala Asn Ala His Asp Val Leu Ala
 145 150 155 160

Arg Ser Arg Ser Arg Ala Asn Val Leu Asn Lys Val Glu Tyr Gly Thr
 165 170 175

Ala Ala Leu Glu Ala Ser Ser Pro Arg Ala Ala Lys Ser Leu Ser Leu
 180 185 190

153

Thr Gly Met Leu Ser Ser Ala Asn Trp Gly Ile Glu Phe Lys Val Thr
 195 200 205

Arg Lys Lys Gln Ala Asp Asn Trp Lys Gly Thr Asp Trp Val Leu Leu
 210 215 220

Gly Phe Ile Leu Ile Pro Cys
 225 230

<210> 227

<211> 456

<212> PRT

<213> Homo sapiens

<400> 227

Met Ala Ala Ala Gly Arg Leu Pro Ser Ser Trp Ala Leu Phe Ser Pro
 1 5 10 15

Leu Leu Ala Gly Leu Ala Leu Leu Gly Val Gly Pro Val Pro Ala Arg
 20 25 30

Ala Leu His Asn Val Thr Ala Glu Leu Phe Gly Ala Glu Ala Trp Gly
 35 40 45

Thr Leu Ala Ala Phe Gly Asp Leu Asn Ser Asp Lys Gln Thr Asp Leu
 50 55 60

Phe Val Leu Arg Glu Arg Asn Asp Leu Ile Val Phe Leu Ala Asp Gln
 65 70 75 80

Asn Ala Pro Tyr Phe Lys Pro Lys Val Lys Val Ser Phe Lys Asn His
 85 90 95

Ser Ala Leu Ile Thr Ser Val Val Pro Gly Asp Tyr Asp Gly Asp Ser
 100 105 110

Gln Met Asp Val Leu Leu Thr Tyr Leu Pro Lys Asn Tyr Ala Lys Ser
 115 120 125

Glu Leu Gly Ala Val Ile Phe Trp Gly Gln Asn Gln Thr Leu Asp Pro
 130 135 140

Asn Asn Met Thr Ile Leu Asn Arg Thr Phe Gln Asp Glu Pro Leu Ile
 145 150 155 160

Met Asp Phe Asn Gly Asp Leu Ile Pro Asp Ile Phe Gly Ile Thr Asn
 165 170 175

Glu Ser Asn Gln Pro Gln Ile Leu Leu Gly Gly Asn Leu Ser Trp His
 180 185 190

Pro Ala Leu Thr Thr Thr Ser Lys Met Arg Ile Pro His Ser His Ala
 195 200 205

Phe Ile Asp Leu Thr Glu Asp Phe Thr Ala Asp Leu Phe Leu Thr Thr
 210 215 220

154

Leu Asn Ala Thr Thr Ser Thr Phe Gln Phe Glu Ile Trp Glu Asn Leu
 225 230 235 240
 Asp Gly Asn Phe Ser Val Ser Thr Ile Leu Glu Lys Pro Gln Asn Met
 245 250 255
 Met Val Val Gly Gln Ser Ala Phe Ala Asp Phe Asp Gly Asp Gly His
 260 265 270
 Met Asp His Leu Leu Pro Gly Cys Glu Asp Lys Asn Cys Gln Lys Ser
 275 280 285
 Thr Ile Tyr Leu Val Arg Ser Gly Met Lys Gln Trp Val Pro Val Leu
 290 295 300
 Gln Asp Phe Ser Asn Lys Gly Thr Leu Trp Gly Phe Val Pro Phe Val
 305 310 315 320
 Asp Glu Gln Gln Pro Thr Glu Ile Pro Ile Pro Ile Thr Leu His Ile
 325 330 335
 Gly Asp Tyr Asn Met Asp Gly Tyr Pro Asp Ala Leu Val Ile Leu Lys
 340 345 350
 Asn Thr Ser Gly Ser Asn Gln Gln Ala Phe Leu Leu Glu Asn Val Pro
 355 360 365
 Cys Asn Asn Ala Ser Cys Glu Glu Ala Arg Arg Met Phe Lys Val Tyr
 370 375 380
 Trp Glu Leu Thr Asp Leu Asn Gln Ile Lys Asp Ala Met Val Ala Thr
 385 390 395 400
 Phe Phe Asp Ile Tyr Glu Asp Gly Ile Leu Asp Ile Val Val Leu Ser
 405 410 415
 Lys Gly Tyr Thr Lys Asn Asp Phe Ala Ile His Thr Leu Lys Asn Asn
 420 425 430
 Phe Glu Ala Asp Ala Tyr Phe Val Lys Val Ile Val Leu Ser Gly Leu
 435 440 445
 Cys Ser Asn Asp Cys Pro Arg Arg
 450 455

<210> 228

<211> 282

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (144)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

155

<221> SITE

<222> (168)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 228

Met	Thr	Lys	Arg	Glu	Asp	Gly	Gly	Tyr	Thr	Phe	Thr	Ala	Thr	Pro	Glu
1				5					10					15	

Asp	Phe	Pro	Lys	Lys	His	Lys	Ala	Pro	Val	Ile	Asp	Ile	Gly	Ile	Ala
			20					25					30		

Asn	Thr	Gly	Lys	Phe	Ile	Met	Thr	Ala	Ser	Ser	Asp	Thr	Thr	Val	Leu
		35					40					45			

Ile	Trp	Ser	Leu	Lys	Gly	Gln	Val	Leu	Ser	Thr	Ile	Asn	Thr	Asn	Gln
	50					55					60				

Met	Asn	Asn	Thr	His	Ala	Ala	Val	Ser	Pro	Cys	Gly	Arg	Phe	Val	Ala
	65				70					75					80

Ser	Cys	Gly	Phe	Thr	Pro	Asp	Val	Lys	Val	Trp	Glu	Val	Cys	Phe	Gly
				85					90					95	

Lys	Lys	Gly	Glu	Phe	Gln	Glu	Val	Val	Arg	Ala	Phe	Glu	Leu	Lys	Gly
			100					105					110		

His	Ser	Ala	Ala	Val	His	Ser	Phe	Ala	Phe	Ser	Asn	Asp	Ser	Arg	Arg
		115					120					125			

Met	Ala	Ser	Val	Ser	Lys	Asp	Gly	Thr	Trp	Lys	Leu	Trp	Asp	Thr	Xaa
	130					135					140				

Val	Glu	Tyr	Lys	Lys	Lys	Gln	Asp	Pro	Tyr	Leu	Leu	Lys	Thr	Gly	Arg
	145				150					155					160

Phe	Glu	Glu	Ala	Ala	Gly	Ala	Xaa	Pro	Cys	Arg	Leu	Ala	Leu	Ser	Pro
			165						170					175	

Asn	Ala	Gln	Val	Leu	Ala	Leu	Ala	Ser	Gly	Ser	Ser	Ile	His	Leu	Tyr
		180						185					190		

Asn	Thr	Arg	Arg	Gly	Glu	Lys	Glu	Glu	Cys	Phe	Glu	Arg	Val	His	Gly
		195				200						205			

Glu	Cys	Ile	Ala	Asn	Leu	Ser	Phe	Asp	Ile	Thr	Gly	Arg	Phe	Leu	Ala
	210					215					220				

Ser	Cys	Gly	Asp	Arg	Ala	Val	Arg	Leu	Phe	His	Asn	Thr	Pro	Gly	His
	225				230					235				240	

Arg	Ala	Met	Val	Glu	Glu	Met	Gln	Gly	His	Leu	Lys	Arg	Ala	Ser	Asn
			245						250					255	

Glu	Ser	Thr	Arg	Gln	Arg	Leu	Gln	Gln	Gln	Leu	Thr	Gln	Ala	Gln	Glu
			260				265						270		

Thr	Leu	Lys	Ser	Leu	Gly	Ala	Leu	Lys	Lys
		275					280		

<210> 229
 <211> 456
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (17)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (37)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (318)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (342)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 229
 Val Ile Arg His Glu Gly Ser Thr Asn Met Glu Leu Ser Gln Met Ser
 1 5 10 15
 Xaa Leu Met Gly Leu Ser Val Leu Leu Gly Leu Leu Ala Leu Met Ala
 20 25 30
 Thr Ala Ala Val Xaa Arg Gly Trp Leu Arg Ala Gly Glu Glu Arg Ser
 35 40 45
 Gly Arg Pro Ala Cys Gln Lys Ala Asn Gly Phe Pro Pro Asp Lys Ser
 50 55 60
 Ser Gly Ser Lys Lys Gln Lys Gln Tyr Gln Arg Ile Arg Lys Glu Lys
 65 70 75 80
 Pro Gln Gln His Asn Phe Thr His Arg Leu Leu Ala Ala Ala Leu Lys
 85 90 95
 Ser His Ser Gly Asn Ile Ser Cys Met Asp Phe Ser Ser Asn Gly Lys
 100 105 110
 Tyr Leu Ala Thr Cys Ala Asp Asp Arg Thr Ile Arg Ile Trp Ser Thr
 115 120 125
 Lys Asp Phe Leu Gln Arg Glu His Arg Ser Met Arg Ala Asn Val Glu
 130 135 140
 Leu Asp His Ala Thr Leu Val Arg Phe Ser Pro Asp Cys Arg Ala Phe
 145 150 155 160

157

Ile Val Trp Leu Ala Asn Gly Asp Thr Leu Arg Val Phe Lys Met Thr
 165 170 175
 Lys Arg Glu Asp Gly Gly Tyr Thr Phe Thr Ala Thr Pro Glu Asp Phe
 180 185 190
 Pro Lys Lys His Lys Ala Pro Val Ile Asp Ile Gly Ile Ala Asn Thr
 195 200 205
 Gly Lys Phe Ile Met Thr Ala Ser Ser Asp Thr Thr Val Leu Ile Trp
 210 215 220
 Ser Leu Lys Gly Gln Val Leu Ser Thr Ile Asn Thr Asn Gln Met Asn
 225 230 235 240
 Asn Thr His Ala Ala Val Ser Pro Cys Gly Arg Phe Val Ala Ser Cys
 245 250 255
 Gly Phe Thr Pro Asp Val Lys Val Trp Glu Val Cys Phe Gly Lys Lys
 260 265 270
 Gly Glu Phe Gln Glu Val Val Arg Ala Phe Glu Leu Lys Gly His Ser
 275 280 285
 Ala Ala Val His Ser Phe Ala Phe Ser Asn Asp Ser Arg Arg Met Ala
 290 295 300
 Ser Val Ser Lys Asp Gly Thr Trp Lys Leu Trp Asp Thr Xaa Val Glu
 305 310 315 320
 Tyr Lys Lys Lys Gln Asp Pro Tyr Leu Leu Lys Thr Gly Arg Phe Glu
 325 330 335
 Glu Ala Ala Gly Ala Xaa Pro Cys Arg Leu Ala Leu Ser Pro Asn Ala
 340 345 350
 Gln Val Leu Ala Leu Ala Ser Gly Ser Ser Ile His Leu Tyr Asn Thr
 355 360 365
 Arg Arg Gly Glu Lys Glu Glu Cys Phe Glu Arg Val His Gly Glu Cys
 370 375 380
 Ile Ala Asn Leu Ser Phe Asp Ile Thr Gly Arg Phe Leu Ala Ser Cys
 385 390 395 400
 Gly Asp Arg Ala Val Arg Leu Phe His Asn Thr Pro Gly His Arg Ala
 405 410 415
 Met Val Glu Glu Met Gln Gly His Leu Lys Arg Ala Ser Asn Glu Ser
 420 425 430
 Thr Arg Gln Arg Leu Gln Gln Gln Leu Thr Gln Ala Gln Glu Thr Leu
 435 440 445
 Lys Ser Leu Gly Ala Leu Lys Lys
 450 455

158

<210> 230

<211> 363

<212> PRT

<213> Homo sapiens

<400> 230

```

Met Ser Val Met Val Val Arg Lys Lys Val Thr Arg Lys Trp Glu Lys
  1              5              10              15

Leu Pro Gly Arg Asn Thr Phe Cys Cys Asp Gly Arg Val Met Met Ala
      20              25              30

Arg Gln Lys Gly Ile Phe Tyr Leu Thr Leu Phe Leu Ile Leu Gly Thr
      35              40              45

Cys Thr Leu Phe Phe Ala Phe Glu Cys Arg Tyr Leu Ala Val Gln Leu
      50              55              60

Ser Pro Ala Ile Pro Val Phe Ala Ala Met Leu Phe Leu Phe Ser Met
      65              70              75              80

Ala Thr Leu Leu Arg Thr Ser Phe Ser Asp Pro Gly Val Ile Pro Arg
      85              90              95

Ala Leu Pro Asp Glu Ala Ala Phe Ile Glu Met Glu Ile Glu Ala Thr
      100              105              110

Asn Gly Ala Val Pro Gln Gly Gln Arg Pro Pro Pro Arg Ile Lys Asn
      115              120              125

Phe Gln Ile Asn Asn Gln Ile Val Lys Leu Lys Tyr Cys Tyr Thr Cys
      130              135              140

Lys Ile Phe Arg Pro Pro Arg Ala Ser His Cys Ser Ile Cys Asp Asn
      145              150              155              160

Cys Val Glu Arg Phe Asp His His Cys Pro Trp Val Gly Asn Cys Val
      165              170              175

Gly Lys Arg Asn Tyr Arg Tyr Phe Tyr Leu Phe Ile Leu Ser Leu Ser
      180              185              190

Leu Leu Thr Ile Tyr Val Phe Ala Phe Asn Ile Val Tyr Val Ala Leu
      195              200              205

Lys Ser Leu Lys Ile Gly Phe Leu Glu Thr Leu Lys Glu Thr Pro Gly
      210              215              220

Thr Val Leu Glu Val Leu Ile Cys Phe Phe Thr Leu Trp Ser Val Val
      225              230              235              240

Gly Leu Thr Gly Phe His Thr Phe Leu Val Ala Leu Asn Gln Thr Thr
      245              250              255

Asn Glu Asp Ile Lys Gly Ser Trp Thr Gly Lys Asn Arg Val Gln Asn
      260              265              270

Pro Tyr Ser His Gly Asn Ile Val Lys Asn Cys Cys Glu Val Leu Cys

```

159

275 280 285
 Gly Pro Leu Pro Pro Ser Val Leu Asp Arg Arg Gly Ile Leu Pro Leu
 290 295 300
 Glu Glu Ser Gly Ser Arg Pro Pro Ser Thr Gln Glu Thr Ser Ser Ser
 305 310 315 320
 Leu Leu Pro Gln Ser Pro Ala Pro Thr Glu Leu Asn Ser Asn Glu Met
 325 330 335
 Pro Glu Asp Ser Ser Thr Pro Glu Glu Met Pro Pro Pro Glu Pro Pro
 340 345 350
 Glu Pro Pro Gln Glu Ala Ala Glu Ala Glu Lys
 355 360

<210> 231

<211> 184

<212> PRT

<213> Homo sapiens

<400> 231

Met Leu Phe Leu Phe Ser Met Ala Thr Leu Leu Arg Thr Ser Phe Ser
 1 5 10 15
 Asp Pro Gly Val Ile Pro Arg Ala Leu Pro Asp Glu Ala Ala Phe Ile
 20 25 30
 Glu Met Glu Ile Glu Ala Thr Asn Gly Ala Val Pro Gln Gly Gln Arg
 35 40 45
 Pro Pro Pro Arg Ile Lys Asn Phe Gln Ile Asn Asn Gln Ile Val Lys
 50 55 60
 Leu Lys Tyr Cys Tyr Thr Cys Lys Ile Phe Arg Pro Pro Arg Ala Ser
 65 70 75 80
 His Cys Ser Ile Cys Asp Asn Cys Val Glu Arg Phe Asp His His Cys
 85 90 95
 Pro Trp Val Gly Asn Cys Val Gly Lys Arg Asn Tyr Arg Tyr Phe Tyr
 100 105 110
 Leu Phe Ile Leu Ser Leu Ser Leu Leu Thr Ile Tyr Val Phe Ala Phe
 115 120 125
 Asn Ile Val Tyr Val Ala Leu Lys Ser Leu Lys Ile Gly Phe Leu Glu
 130 135 140
 Thr Leu Lys Gly Asn Ser Trp Asn Cys Ser Arg Ser Pro His Leu Leu
 145 150 155 160
 Leu Tyr Thr Leu Val Arg Arg Gly Thr Asp Trp Ile Ser Tyr Phe Pro
 165 170 175
 Arg Gly Ser Gln Pro Asp Asn Gln

160

180

<210> 232
 <211> 52
 <212> PRT
 <213> Homo sapiens

<400> 232
 Leu His Glu Cys Leu Pro Gly Ser Ile Ser Tyr Leu His Pro Arg Thr
 1 5 10 15
 Pro Trp Leu Cys Leu Pro Pro Gln His Leu Ser Phe Ser Thr Phe Ser
 20 25 30
 Pro Pro Trp Gln Pro Ala Met Ser Pro Val Pro Gly Thr Gly Gly Pro
 35 40 45
 Pro Cys Gly Leu
 50

<210> 233
 <211> 177
 <212> PRT
 <213> Homo sapiens

<400> 233
 Met Leu Pro Leu Leu Ile Ile Cys Leu Leu Pro Ala Ile Glu Gly Lys
 1 5 10 15
 Asn Cys Leu Arg Cys Trp Pro Glu Leu Ser Ala Leu Ile Asp Tyr Asp
 20 25 30
 Leu Gln Ile Leu Trp Val Thr Pro Gly Pro Pro Thr Glu Leu Ser Gln
 35 40 45
 Ser Ile His Ser Leu Phe Leu Glu Asp Asn Asn Phe Leu Lys Pro Trp
 50 55 60
 Tyr Leu Asp Arg Asp His Leu Glu Glu Glu Thr Ala Lys Phe Phe Thr
 65 70 75 80
 Gln Val His Gln Ala Ile Lys Thr Leu Arg Asp Asp Lys Thr Val Leu
 85 90 95
 Leu Glu Glu Ile Tyr Thr His Lys Asn Leu Phe Thr Glu Arg Leu Asn
 100 105 110
 Lys Ile Ser Asp Gly Leu Lys Glu Lys Gly Ala Pro Pro Leu His Glu
 115 120 125
 Cys Leu Pro Gly Ser Ile Ser Tyr Leu His Pro Arg Thr Pro Trp Leu
 130 135 140
 Cys Leu Pro Pro Gln His Leu Ser Phe Ser Thr Phe Ser Pro Pro Trp
 145 150 155 160

161

Gln Pro Ala Met Ser Pro Val Pro Gly Thr Gly Gly Pro Pro Cys Gly
 165 170 175

Leu

<210> 234
 <211> 95
 <212> PRT
 <213> Homo sapiens

<400> 234
 Pro Pro Val Pro Pro Trp Ile Ser Leu Pro Leu Thr Gly Ser Pro Pro
 1 5 10 15
 Arg Pro Gly Phe Val Pro Val Ser Pro Phe Cys Phe Ser Pro Met Thr
 20 25 30
 Asn Gly His Gln Val Leu Leu Leu Leu Leu Thr Ser Ala Val Ala
 35 40 45
 Ala Gly Pro Trp Pro Gln Val His Ala Gly Gln Trp Gly Trp Met Cys
 50 55 60
 Leu Pro Pro Gly Leu Pro Ser Val Gln Ala Arg Ser Gly Leu Gly Gly
 65 70 75 80
 Leu Pro Gly Gly Pro Gln Trp Val Pro Gly Gly Ala Arg Gly Tyr
 85 90 95

<210> 235
 <211> 404
 <212> PRT
 <213> Homo sapiens

<400> 235
 Ile Gln Gln Trp Gly Asp Ser Val Leu Gly Arg Arg Cys Arg Asp Leu
 1 5 10 15
 Leu Leu Gln Leu Tyr Leu Gln Arg Pro Glu Leu Arg Val Pro Val Pro
 20 25 30
 Glu Val Leu Leu His Ser Glu Gly Ala Ala Ser Ser Ser Val Cys Lys
 35 40 45
 Leu Asp Gly Leu Ile His Arg Phe Ile Thr Leu Leu Ala Asp Thr Ser
 50 55 60
 Asp Ser Arg Ala Leu Glu Asn Arg Gly Ala Asp Ala Ser Met Ala Cys
 65 70 75 80
 Arg Lys Leu Ala Val Ala His Pro Leu Leu Leu Leu Arg His Leu Pro
 85 90 95
 Met Ile Ala Ala Leu Leu His Gly Arg Thr His Leu Asn Phe Gln Glu
 100 105 110

162

Phe Arg Gln Gln Asn His Leu Ser Cys Phe Leu His Val Leu Gly Leu
 115 120 125
 Leu Glu Leu Leu Gln Pro His Val Phe Arg Ser Glu His Gln Gly Ala
 130 135 140
 Leu Trp Asp Cys Leu Leu Ser Phe Ile Arg Leu Leu Leu Asn Tyr Arg
 145 150 155 160
 Lys Ser Ser Arg His Leu Ala Ala Phe Ile Asn Lys Phe Val Gln Phe
 165 170 175
 Ile His Lys Tyr Ile Thr Tyr Asn Ala Pro Ala Ala Ile Ser Phe Leu
 180 185 190
 Gln Lys His Ala Asp Pro Leu His Asp Leu Ser Phe Asp Asn Ser Asp
 195 200 205
 Leu Val Met Leu Lys Ser Leu Leu Ala Gly Leu Ser Leu Pro Ser Arg
 210 215 220
 Asp Asp Arg Thr Asp Arg Gly Leu Asp Glu Glu Gly Glu Glu Glu Ser
 225 230 235 240
 Ser Ala Gly Ser Leu Pro Leu Val Ser Val Ser Leu Phe Thr Pro Leu
 245 250 255
 Thr Ala Ala Glu Met Ala Pro Tyr Met Lys Arg Leu Ser Arg Gly Gln
 260 265 270
 Thr Val Glu Asp Leu Leu Glu Val Leu Ser Asp Ile Asp Glu Met Ser
 275 280 285
 Arg Arg Arg Pro Glu Ile Leu Ser Phe Phe Ser Thr Asn Leu Gln Arg
 290 295 300
 Leu Met Ser Ser Ala Glu Glu Cys Cys Arg Asn Leu Ala Phe Ser Leu
 305 310 315 320
 Ala Leu Arg Ser Met Gln Asn Ser Pro Ser Ile Ala Ala Ala Phe Leu
 325 330 335
 Pro Thr Phe Met Tyr Cys Leu Gly Ser Gln Asp Phe Glu Val Val Gln
 340 345 350
 Thr Ala Leu Arg Asn Leu Pro Glu Tyr Ala Leu Leu Cys Gln Glu His
 355 360 365
 Ala Ala Val Leu Leu His Arg Ala Phe Leu Val Gly Met Tyr Gly Gln
 370 375 380
 Met Asp Pro Ser Ala Gln Ile Ser Glu Ala Leu Arg Ile Leu His Met
 385 390 395 400
 Glu Ala Val Met

163

<210> 236
 <211> 361
 <212> PRT
 <213> Homo sapiens

<400> 236

```

Met Leu Leu Lys His Leu Gln Arg Met Val Ser Val Pro Gln Val Lys
  1              5              10              15

Ala Ser Ala Leu Lys Val Val Thr Leu Thr Ala Asn Asp Lys Thr Ser
      20              25              30

Val Ser Phe Ser Ser Leu Pro Gly Gln Gly Val Ile Tyr Asn Val Ile
      35              40              45

Val Trp Asp Pro Phe Leu Asn Thr Ser Ala Ala Tyr Ile Pro Ala His
      50              55              60

Thr Tyr Ala Cys Ser Phe Glu Ala Gly Glu Gly Ser Cys Ala Ser Leu
      65              70              75              80

Gly Arg Val Ser Ser Lys Val Phe Phe Thr Leu Phe Ala Leu Leu Gly
      85              90              95

Phe Phe Ile Cys Phe Phe Gly His Arg Phe Trp Lys Thr Glu Leu Phe
      100             105             110

Phe Ile Gly Phe Ile Ile Met Gly Phe Phe Phe Tyr Ile Leu Ile Thr
      115             120             125

Arg Leu Thr Pro Ile Lys Tyr Asp Val Asn Leu Ile Leu Thr Ala Val
      130             135             140

Thr Gly Ser Val Gly Gly Met Phe Leu Val Ala Val Trp Trp Arg Phe
      145             150             155             160

Gly Ile Leu Ser Ile Cys Met Leu Cys Val Gly Leu Val Leu Gly Phe
      165             170             175

Leu Ile Ser Ser Val Thr Phe Phe Thr Pro Leu Gly Asn Leu Lys Ile
      180             185             190

Phe His Asp Asp Gly Val Phe Trp Val Thr Phe Ser Cys Ile Ala Ile
      195             200             205

Leu Ile Pro Val Val Phe Met Gly Cys Leu Arg Ile Leu Asn Ile Leu
      210             215             220

Thr Cys Gly Val Ile Gly Ser Tyr Ser Val Val Leu Ala Ile Asp Ser
      225             230             235             240

Tyr Trp Ser Thr Ser Leu Ser Tyr Ile Thr Leu Asn Val Leu Lys Arg
      245             250             255

Ala Leu Asn Lys Asp Phe His Arg Ala Phe Thr Asn Val Pro Phe Gln
      260             265             270

```

164

Thr Asn Asp Phe Ile Ile Leu Ala Val Trp Gly Met Leu Ala Val Ser
 275 280 285

Gly Ile Thr Leu Gln Ile Arg Arg Glu Arg Gly Arg Pro Phe Phe Pro
 290 295 300

Pro His Pro Tyr Lys Leu Trp Lys Gln Glu Arg Glu Arg Arg Val Thr
 305 310 315 320

Asn Ile Leu Asp Pro Ser Tyr His Ile Pro Pro Leu Arg Glu Arg Leu
 325 330 335

Tyr Gly Arg Leu Thr Gln Ile Lys Gly Leu Phe Gln Lys Glu Gln Pro
 340 345 350

Ala Gly Glu Arg Thr Pro Leu Leu Leu
 355 360

<210> 237

<211> 116

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (37)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (40)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 237

Trp Ala Arg Leu Arg Gly Pro Gly Ala His Ala Arg Thr Ser Pro Gln
 1 5 10 15

Pro Trp Arg Gly Pro Ser Pro Ala Gln Ala Ala Met Gly Phe Leu Gln
 20 25 30

Leu Leu Val Val Xaa Val Leu Xaa Ser Glu His Arg Val Ala Gly Ala
 35 40 45

Ala Glu Val Phe Gly Asn Ser Ser Glu Gly Leu Ile Glu Phe Ser Val
 50 55 60

Gly Lys Phe Arg Tyr Phe Glu Leu Asn Arg Pro Phe Pro Glu Glu Ala
 65 70 75 80

Ile Leu His Asp Ile Ser Ser Asn Val Thr Phe Leu Ile Phe Gln Ile
 85 90 95

His Ser Gln Tyr Gln Asn Thr Thr Val Ser Phe Ser Pro Arg Arg Arg
 100 105 110

Ser Pro Thr Met
 115

165

<210> 238
 <211> 166
 <212> PRT
 <213> Homo sapiens

<400> 238

Pro Arg Val Arg Pro Ala Ser Pro Pro Val Arg Ser Pro Ala Arg Trp
 1 5 10 15
 Gly Ser Met Ala Gly Ser Pro Leu Leu Trp Gly Pro Arg Ala Gly Gly
 20 25 30
 Val Gly Leu Leu Val Leu Leu Leu Gly Leu Phe Arg Pro Pro Pro
 35 40 45
 Ala Leu Cys Ala Arg Pro Val Lys Glu Pro Arg Gly Leu Ser Ala Ala
 50 55 60
 Ser Pro Pro Leu Ala Arg Leu Ala Leu Leu Ala Ala Ser Gly Gly Gln
 65 70 75 80
 Cys Pro Glu Val Arg Arg Arg Gly Arg Cys Arg Pro Gly Ala Gly Ala
 85 90 95
 Gly Ala Ser Ala Gly Ala Glu Arg Gln Glu Arg Ala Arg Ala Glu Ala
 100 105 110
 Gln Arg Leu Arg Ile Ser Arg Arg Ala Ser Trp Arg Ser Cys Cys Ala
 115 120 125
 Ser Gly Ala Pro Pro Ala Thr Leu Ile Arg Leu Trp Ala Trp Thr Thr
 130 135 140
 Thr Pro Thr Arg Leu Gln Arg Ser Ser Leu Ala Leu Cys Ser Ala Pro
 145 150 155 160
 Ala Leu Thr Leu Pro Pro
 165

<210> 239
 <211> 414
 <212> PRT
 <213> Homo sapiens

<400> 239

Pro Arg Val Arg Leu Ala Thr Pro Asn Ile Trp Asp Leu Ser Met Leu
 1 5 10 15
 Phe Ala Phe Ile Ser Leu Leu Val Met Leu Pro Thr Trp Trp Ile Val
 20 25 30
 Ser Ser Trp Leu Val Trp Gly Val Ile Leu Phe Val Tyr Leu Val Ile
 35 40 45
 Arg Ala Leu Arg Leu Trp Arg Thr Ala Lys Leu Gln Val Thr Leu Lys

166

50	55	60
Lys Tyr Ser Val His Leu Glu Asp Met Ala Thr Asn Ser Arg Ala Phe		
65	70	75 80
Thr Asn Leu Val Arg Lys Ala Leu Arg Leu Ile Gln Glu Thr Glu Val		
	85	90 95
Ile Ser Arg Gly Phe Thr Leu Val Ser Ala Ala Cys Pro Phe Asn Lys		
	100	105 110
Ala Gly Gln His Pro Ser Gln His Leu Ile Gly Leu Arg Lys Ala Val		
	115	120 125
Tyr Arg Thr Leu Arg Ala Asn Phe Gln Ala Ala Arg Leu Ala Thr Leu		
	130	135 140
Tyr Met Leu Lys Asn Tyr Pro Leu Asn Ser Glu Ser Asp Asn Val Thr		
	145	150 155 160
Asn Tyr Ile Cys Val Val Pro Phe Lys Glu Leu Gly Leu Gly Leu Ser		
	165	170 175
Glu Glu Gln Ile Ser Glu Glu Glu Ala His Asn Phe Thr Asp Gly Phe		
	180	185 190
Ser Leu Pro Ala Leu Lys Val Leu Phe Gln Leu Trp Val Ala Gln Ser		
	195	200 205
Ser Glu Phe Phe Arg Arg Leu Ala Leu Leu Leu Ser Thr Ala Asn Ser		
	210	215 220
Pro Pro Gly Pro Leu Leu Thr Pro Ala Leu Leu Pro His Arg Ile Leu		
	225	230 235 240
Ser Asp Val Thr Gln Gly Leu Pro His Ala His Ser Ala Cys Leu Glu		
	245	250 255
Glu Leu Lys Arg Ser Tyr Glu Phe Tyr Arg Tyr Phe Glu Thr Gln His		
	260	265 270
Gln Ser Val Pro Gln Cys Leu Ser Lys Thr Gln Gln Lys Ser Arg Glu		
	275	280 285
Leu Asn Asn Val His Thr Ala Val Arg Ser Leu Gln Leu His Leu Lys		
	290	295 300
Ala Leu Leu Asn Glu Val Ile Ile Leu Glu Asp Glu Leu Glu Lys Leu		
	305	310 315 320
Val Cys Thr Lys Glu Thr Gln Glu Leu Val Ser Glu Ala Tyr Pro Ile		
	325	330 335
Leu Glu Gln Lys Leu Lys Leu Ile Gln Pro His Val Gln Ala Ser Asn		
	340	345 350
Asn Cys Trp Glu Glu Ala Ile Ser Gln Val Asp Lys Leu Leu Arg Arg		
	355	360 365

167

Asn Thr Asp Lys Lys Gly Lys Pro Glu Ile Ala Cys Glu Asn Pro His
370 375 380

Cys Thr Val Ser Thr Phe Glu Ala Ala Tyr Ser Thr His Cys Arg Gln
385 390 395 400

Arg Ser Asn Pro Arg Gly Ala Gly Ile Arg Ser Leu Cys Arg
405 410

<210> 240

<211> 145

<212> PRT

<213> Homo sapiens

<400> 240

Ala Ala Pro His Pro Pro Leu Leu Arg Pro Leu Cys Leu Trp Cys Pro
1 5 10 15

Leu Trp Pro Ala Trp Pro Leu Arg Gly Arg Pro Arg Ser Ala Trp Lys
20 25 30

Arg Trp Pro Pro Leu Pro Val Gly Pro Ala Lys Leu Gly Cys Ser Met
35 40 45

Thr Thr Arg Gln Pro Thr Ala Val Ser Trp Pro Cys Trp Leu Met Ser
50 55 60

Ser Ser Leu Ser Thr Ala Cys Leu Ala Trp Thr Leu Thr Gly Ser Leu
65 70 75 80

Ala Arg Glu Ala Thr Arg Arg Ala Arg Ser Leu Ser Pro Thr Trp Asn
85 90 95

Cys Ser Ala Arg Gln Val Pro Pro Ser Pro Pro His Ser Gly Leu Gly
100 105 110

Arg Arg Gly Trp Ala His Cys His Leu Thr Cys Leu Leu Val Thr Gln
115 120 125

Leu Phe Arg Val Gly Arg Ile His Pro Ile Leu Ser Leu Pro Leu Val
130 135 140

Thr

145

<210> 241

<211> 72

<212> PRT

<213> Homo sapiens

<400> 241

Leu Gln Leu Ala Ser Gln Ser Ala Gly Ile Lys Gly Met Ser His Cys
1 5 10 15

Ala Arg Pro Thr Phe Leu Thr Leu Leu Leu Ala Ser Cys Phe Trp Ala

168

20 25 30
 Ala Ala Ile Pro Asn Arg Asn Val Ile Leu Ser Val Ser Phe Arg Pro
 35 40 45
 Leu His Met Gln Phe Thr Leu Ser Ile Leu Val Phe Ile Leu Arg Ile
 50 55 60
 Leu Ile Leu Leu Arg Ser Phe Leu
 65 70

<210> 242
 <211> 140
 <212> PRT
 <213> Homo sapiens

<400> 242
 Met Val Leu Val Leu Arg His Pro Leu Cys Ala Arg Glu Arg Ala Phe
 1 5 10 15
 Arg Glu Pro Gly Arg Gly Leu Leu Thr Arg Thr Gly Gln His Asp Gly
 20 25 30
 Ala Pro Ala Val Thr Ala Val Pro Gly Pro Leu Gly Ala Val Ala Ala
 35 40 45
 Ala Glu Gly Arg Arg Ser Ala Trp Gly Ala Gly Gly Ser Ser Pro Pro
 50 55 60
 Arg Lys Val Leu Trp Gly Asp Met Arg Gly Arg Arg Ala Gly Val Asp
 65 70 75 80
 Val Leu Gly Pro Ala Leu Ser Ser Glu Ala Ala Gly Ala Glu Ala Arg
 85 90 95
 Gly Trp Gly Met Pro Gly Met Gly Val Gly Val Gly Ala Ser Glu Thr
 100 105 110
 Arg Gly Ala Leu Phe Leu Gly Arg Glu Gly Val His Gly Pro Cys Pro
 115 120 125
 Met Asp Gly Leu Gly Pro Trp Pro Trp Gly Pro Trp
 130 135 140

<210> 243
 <211> 353
 <212> PRT
 <213> Homo sapiens

<400> 243
 Met Gly Pro Ala Val Lys Met Trp Thr Asn Ala Trp Lys Gly Leu Asp
 1 5 10 15
 Asp Cys His Tyr Asn Gln Leu Cys Glu Asn Thr Pro Gly Gly His Arg
 20 25 30

169

Cys Ser Cys Pro Arg Gly Tyr Arg Met Gln Gly Pro Ser Leu Pro Cys
 35 40 45
 Leu Asp Val Asn Glu Cys Leu Gln Leu Pro Lys Ala Cys Ala Tyr Gln
 50 55 60
 Cys His Asn Leu Gln Gly Ser Tyr Arg Cys Leu Cys Pro Pro Gly Gln
 65 70 75 80
 Thr Leu Leu Arg Asp Gly Lys Ala Cys Thr Ser Leu Glu Arg Asn Gly
 85 90 95
 Gln Asn Val Thr Thr Val Ser His Arg Gly Pro Leu Leu Pro Trp Leu
 100 105 110
 Arg Pro Trp Ala Ser Ile Pro Gly Thr Ser Tyr His Ala Trp Val Ser
 115 120 125
 Leu Arg Pro Gly Pro Met Ala Leu Ser Ser Val Gly Arg Ala Trp Cys
 130 135 140
 Pro Pro Gly Phe Ile Arg Gln Asn Gly Val Cys Thr Asp Leu Asp Glu
 145 150 155 160
 Cys Arg Val Arg Asn Leu Cys Gln His Ala Cys Arg Asn Thr Glu Gly
 165 170 175
 Ser Tyr Gln Cys Leu Cys Pro Ala Gly Tyr Arg Leu Leu Pro Ser Gly
 180 185 190
 Lys Asn Cys Gln Asp Ile Asn Glu Cys Glu Glu Glu Ser Ile Glu Cys
 195 200 205
 Gly Pro Gly Gln Met Cys Phe Asn Thr Arg Gly Ser Tyr Gln Cys Val
 210 215 220
 Asp Thr Pro Cys Pro Ala Thr Tyr Arg Gln Gly Pro Ser Pro Gly Thr
 225 230 235 240
 Cys Phe Arg Arg Cys Ser Gln Asp Cys Gly Thr Gly Gly Pro Ser Thr
 245 250 255
 Leu Gln Tyr Arg Leu Leu Pro Leu Pro Leu Gly Val Arg Ala His His
 260 265 270
 Asp Val Ala Arg Leu Thr Ala Phe Ser Glu Val Gly Val Pro Ala Asn
 275 280 285
 Arg Thr Glu Leu Ser Met Leu Glu Pro Asp Pro Arg Ser Pro Phe Ala
 290 295 300
 Leu Arg Pro Leu Arg Ala Gly Leu Gly Ala Val Tyr Thr Arg Arg Ala
 305 310 315 320
 Leu Thr Arg Ala Gly Leu Tyr Arg Leu Thr Val Arg Ala Ala Ala Pro
 325 330 335
 Arg His Gln Ser Val Phe Val Leu Leu Ile Ala Val Ser Pro Tyr Pro

170

340

345

350

Tyr

<210> 244

<211> 146

<212> PRT

<213> Homo sapiens

<400> 244

Met Arg Val Leu Val Val Thr Ile Ala Pro Ile Tyr Trp Ala Leu Ala
 1 5 10 15

Arg Glu Ser Gly Glu Ala Leu Asn Gly His Ser Leu Thr Gly Gly Lys
 20 25 30

Phe Arg Gln Ser His Thr Trp Ser Leu Leu Gln Gly Ala Ala His Asp
 35 40 45

Asp Pro Val Ala Arg Gly Leu Asp Pro Asp Gly Leu Leu Leu Asp
 50 55 60

Val Val Val Asn Gly Val Val Pro Gly Arg Ala Trp Leu Thr Gln Ile
 65 70 75 80

Phe Lys Cys Arg Thr Leu Lys Lys His Tyr Val Gln Thr Arg Ala Trp
 85 90 95

Pro Ala Val Arg Gly Leu His Thr Ala Leu Leu Pro Gly Arg Pro Pro
 100 105 110

Leu Val Pro Thr Leu Gln Pro Gln His Pro Val Gln Arg Gly Pro Gly
 115 120 125

Pro Pro Ala Pro Ala Gly Ala Ala Pro Ala Gly Leu Ser Tyr Gln Leu
 130 135 140

Gly Leu
 145

<210> 245

<211> 638

<212> PRT

<213> Homo sapiens

<400> 245

His Ala Ser Gly Ala Phe Leu Val Val Arg Gly Glu Pro Gln Gly Ser
 1 5 10 15

Trp Gly Ser Met Thr Gly Val Ile Asn Gly Arg Lys Phe Gly Val Ala
 20 25 30

Thr Leu Asn Thr Ser Val Met Gln Glu Ala His Ser Gly Val Ser Ser
 35 40 45

171

Ile His Ser Ser Ile Arg His Val Pro Ala Asn Val Gly Pro Leu Met
 50 55 60

Arg Val Leu Val Val Thr Ile Ala Pro Ile Tyr Trp Ala Leu Ala Arg
 65 70 75 80

Glu Ser Gly Glu Ala Leu Asn Gly His Ser Leu Thr Gly Gly Lys Phe
 85 90 95

Arg Gln Glu Ser His Val Glu Phe Ala Thr Gly Glu Leu Leu Thr Met
 100 105 110

Thr Gln Trp Pro Gly Val Trp Ile Pro Met Ala Ser Cys Ser Ser Thr
 115 120 125

Trp Trp Ser Met Ala Leu Ser Pro Asp Ser Leu Ala Asp Ala Asp Leu
 130 135 140

Gln Val Gln Asp Phe Glu Glu His Tyr Val Gln Thr Gly Pro Gly Gln
 145 150 155 160

Leu Phe Val Gly Ser Thr Gln Arg Phe Phe Gln Gly Gly Leu Pro Ser
 165 170 175

Phe Leu Arg Cys Asn His Ser Ile Gln Tyr Asn Ala Ala Arg Gly Pro
 180 185 190

Gln Pro Gln Leu Val Gln His Leu Arg Ala Ser Ala Ile Ser Ser Ala
 195 200 205

Phe Asp Pro Glu Ala Glu Ala Leu Arg Phe Gln Leu Ala Thr Ala Leu
 210 215 220

Gln Ala Glu Glu Asn Glu Val Gly Cys Pro Glu Gly Phe Glu Leu Asp
 225 230 235 240

Ser Gln Gly Ala Phe Cys Val Asp Val Asp Glu Cys Ala Trp Asp Ala
 245 250 255

His Leu Cys Arg Glu Gly Gln Arg Cys Val Asn Leu Leu Gly Ser Tyr
 260 265 270

Arg Cys Leu Pro Asp Cys Gly Pro Gly Phe Arg Val Ala Asp Gly Ala
 275 280 285

Gly Cys Glu Asp Val Asp Glu Cys Leu Glu Gly Leu Asp Asp Cys His
 290 295 300

Tyr Asn Gln Leu Cys Glu Asn Thr Pro Gly Gly His Arg Cys Ser Cys
 305 310 315 320

Pro Arg Gly Tyr Arg Met Gln Gly Pro Ser Leu Pro Cys Leu Asp Val
 325 330 335

Asn Glu Cys Leu Gln Leu Pro Lys Ala Cys Ala Tyr Gln Cys His Asn
 340 345 350

Leu Gln Gly Ser Tyr Arg Cys Leu Cys Pro Pro Gly Gln Thr Leu Leu

172

355 360 365
 Arg Asp Gly Lys Ala Cys Thr Ser Leu Glu Arg Asn Gly Gln Asn Val
 370 375 380
 Thr Thr Val Ser His Arg Gly Pro Leu Leu Pro Trp Leu Arg Pro Trp
 385 390 395 400
 Ala Ser Ile Pro Gly Thr Ser Tyr His Ala Trp Val Ser Leu Arg Pro
 405 410 415
 Gly Pro Met Ala Leu Ser Ser Val Gly Arg Ala Trp Cys Pro Pro Gly
 420 425 430
 Phe Ile Arg Gln Asn Gly Val Cys Thr Asp Leu Asp Glu Cys Arg Val
 435 440 445
 Arg Asn Leu Cys Gln His Ala Cys Arg Asn Thr Glu Gly Ser Tyr Gln
 450 455 460
 Cys Leu Cys Pro Ala Gly Tyr Arg Leu Leu Pro Ser Gly Lys Asn Cys
 465 470 475 480
 Gln Asp Ile Asn Glu Cys Glu Glu Glu Ser Ile Glu Cys Gly Pro Gly
 485 490 495
 Gln Met Cys Phe Asn Thr Arg Gly Ser Tyr Gln Cys Val Asp Thr Pro
 500 505 510
 Cys Pro Ala Thr Tyr Arg Gln Gly Pro Ser Pro Gly Thr Cys Phe Arg
 515 520 525
 Arg Cys Ser Gln Asp Cys Gly Thr Gly Gly Pro Ser Thr Leu Gln Tyr
 530 535 540
 Arg Leu Leu Pro Leu Pro Leu Gly Val Arg Ala His His Asp Val Ala
 545 550 555 560
 Arg Leu Thr Ala Phe Ser Glu Val Gly Val Pro Ala Asn Arg Thr Glu
 565 570 575
 Leu Ser Met Leu Glu Pro Asp Pro Arg Ser Pro Phe Ala Leu Arg Pro
 580 585 590
 Leu Arg Ala Gly Leu Gly Ala Val Tyr Thr Arg Arg Ala Leu Thr Arg
 595 600 605
 Ala Gly Leu Tyr Arg Leu Thr Val Arg Ala Ala Ala Pro Arg His Gln
 610 615 620
 Ser Val Phe Val Leu Leu Ile Ala Val Ser Pro Tyr Pro Tyr
 625 630 635

<210> 246

<211> 367

<212> PRT

<213> Homo sapiens

173

<400> 246

```

Met Gly Glu Lys Phe Leu Leu Leu Ala Met Lys Glu Asn His Pro Glu
 1              5              10              15

Cys Phe Cys Lys Ile Leu Lys Ile Leu His Cys Met Asp Pro Gly Glu
      20              25              30

Trp Leu Pro Gln Thr Glu His Cys Val His Leu Thr Pro Lys Glu Phe
      35              40              45

Leu Ile Trp Thr Met Asp Ile Ala Ser Asn Glu Arg Ser Glu Ile Gln
      50              55              60

Ser Val Ala Leu Arg Leu Ala Ser Lys Val Ile Ser His His Met Gln
      65              70              75              80

Thr Cys Val Glu Asn Arg Glu Leu Ile Ala Ala Glu Leu Lys Gln Trp
      85              90              95

Val Gln Leu Val Ile Leu Ser Cys Glu Asp His Leu Pro Thr Glu Ser
      100              105              110

Arg Leu Ala Val Val Glu Val Leu Thr Ser Thr Thr Pro Leu Phe Leu
      115              120              125

Thr Asn Pro His Pro Ile Leu Glu Leu Gln Asp Thr Leu Ala Leu Trp
      130              135              140

Lys Cys Val Leu Thr Leu Leu Gln Ser Glu Glu Gln Ala Val Arg Asp
      145              150              155              160

Ala Ala Thr Glu Thr Val Thr Thr Ala Met Ser Gln Glu Asn Thr Cys
      165              170              175

Gln Ser Thr Glu Phe Ala Phe Cys Gln Val Asp Ala Ser Ile Ala Leu
      180              185              190

Ala Leu Ala Leu Ala Val Leu Cys Asp Leu Leu Gln Gln Trp Asp Gln
      195              200              205

Leu Ala Pro Gly Leu Pro Ile Leu Leu Gly Trp Leu Leu Gly Glu Ser
      210              215              220

Asp Asp Leu Val Ala Cys Val Glu Ser Met His Gln Val Glu Glu Asp
      225              230              235              240

Tyr Leu Phe Glu Lys Ala Glu Val Asn Phe Trp Ala Glu Thr Leu Ile
      245              250              255

Phe Val Lys Tyr Leu Cys Lys His Leu Phe Cys Leu Leu Ser Lys Ser
      260              265              270

Gly Trp Arg Pro Pro Ser Pro Glu Met Leu Cys His Leu Gln Arg Met
      275              280              285

Val Ser Glu Gln Cys His Leu Leu Ser Gln Phe Phe Arg Glu Leu Pro
      290              295              300

```

174

Pro Ala Ala Glu Phe Val Lys Thr Val Glu Phe Thr Arg Leu Arg Ile
 305 310 315 320

Gln Glu Glu Arg Thr Leu Ala Cys Leu Arg Leu Leu Ala Phe Leu Glu
 325 330 335

Gly Lys Glu Gly Glu Asp Thr Leu Val Leu Ser Val Trp Asp Ser Tyr
 340 345 350

Ala Glu Ser Arg Gln Leu Thr Leu Pro Arg Thr Glu Ala Ala Cys
 355 360 365

<210> 247
 <211> 124
 <212> PRT
 <213> Homo sapiens

<400> 247
 Met Gly Glu Pro Asn Arg His Pro Ser Met Phe Leu Leu Leu Leu Val
 1 5 10 15

Leu Glu Arg Leu Tyr Ala Ser Pro Met Asp Gly Thr Ser Ser Ala Leu
 20 25 30

Ser Met Gly Pro Phe Val Pro Phe Ile Met Arg Cys Gly His Ser Pro
 35 40 45

Val Tyr His Ser Arg Glu Met Ala Ala Arg Ala Leu Val Pro Phe Val
 50 55 60

Met Ile Asp His Ile Pro Asn Thr Ile Arg Thr Leu Leu Ser Thr Leu
 65 70 75 80

Pro Ser Cys Thr Asp Gln Cys Phe Arg Ala Lys Pro His Ser Trp Gly
 85 90 95

His Phe Ser Arg Phe Phe His Leu Leu Gln Ala Tyr Ser Asp Ser Lys
 100 105 110

Thr Arg Asn Glu Phe Arg Leu Pro Ala Arg Ala Asp
 115 120

<210> 248
 <211> 674
 <212> PRT
 <213> Homo sapiens

<400> 248
 Met Thr Gly Arg Glu Phe Phe Ser Arg Phe Pro Glu Leu Tyr Pro Phe
 1 5 10 15

Leu Leu Lys Gln Leu Glu Thr Val Ala Asn Thr Val Asp Ser Asp Met
 20 25 30

Gly Glu Pro Asn Arg His Pro Ser Met Phe Leu Leu Leu Leu Val Leu

175

35	40	45
Glu Arg Leu Tyr Ala Ser Pro Met Asp Gly Thr Ser Ser Ala Leu Ser		
50	55	60
Met Gly Pro Phe Val Pro Phe Ile Met Arg Cys Gly His Ser Pro Val		
65	70	75
Tyr His Ser Arg Glu Met Ala Ala Arg Ala Leu Val Pro Phe Val Met		
	85	90
Ile Asp His Ile Pro Asn Thr Ile Arg Thr Leu Leu Ser Thr Leu Pro		
	100	105
Ser Cys Thr Asp Gln Cys Phe Arg Gln Asn His Ile His Gly Thr Leu		
	115	120
Leu Gln Val Phe His Leu Leu Gln Ala Tyr Ser Asp Ser Lys His Gly		
	130	135
Thr Asn Ser Asp Phe Gln His Glu Leu Thr Asp Ile Thr Val Cys Thr		
	145	150
Lys Ala Lys Leu Trp Leu Ala Lys Arg Gln Asn Pro Cys Leu Val Thr		
	165	170
Arg Ala Val Tyr Ile Asp Ile Leu Phe Leu Leu Thr Cys Cys Leu Asn		
	180	185
Arg Ser Ala Lys Asp Asn Gln Pro Val Leu Glu Ser Leu Gly Phe Trp		
	195	200
Glu Glu Val Arg Gly Ile Ile Ser Gly Ser Glu Leu Ile Thr Gly Phe		
	210	215
Pro Trp Ala Phe Lys Val Pro Gly Leu Pro Gln Tyr Leu Gln Ser Leu		
	225	230
Thr Arg Leu Ala Ile Ala Ala Val Trp Ala Ala Ala Ala Lys Ser Gly		
	245	250
Glu Arg Glu Thr Asn Val Pro Ile Ser Phe Ser Gln Leu Leu Glu Ser		
	260	265
Ala Phe Pro Glu Val Arg Ser Leu Thr Leu Glu Ala Leu Leu Glu Lys		
	275	280
Phe Leu Ala Ala Ala Ser Gly Leu Gly Glu Lys Gly Val Pro Pro Leu		
	290	295
Leu Cys Asn Met Gly Glu Lys Phe Leu Leu Leu Ala Met Lys Glu Asn		
	305	310
His Pro Glu Cys Phe Cys Lys Ile Leu Lys Ile Leu His Cys Met Asp		
	325	330
Pro Gly Glu Trp Leu Pro Gln Thr Glu His Cys Val His Leu Thr Pro		
	340	345
		350

176

Lys Glu Phe Leu Ile Trp Thr Met Asp Ile Ala Ser Asn Glu Arg Ser
 355 360 365
 Glu Ile Gln Ser Val Ala Leu Arg Leu Ala Ser Lys Val Ile Ser His
 370 375 380
 His Met Gln Thr Cys Val Glu Asn Arg Glu Leu Ile Ala Ala Glu Leu
 385 390 395 400
 Lys Gln Trp Val Gln Leu Val Ile Leu Ser Cys Glu Asp His Leu Pro
 405 410 415
 Thr Glu Ser Arg Leu Ala Val Val Glu Val Leu Thr Ser Thr Thr Pro
 420 425 430
 Leu Phe Leu Thr Asn Pro His Pro Ile Leu Glu Leu Gln Asp Thr Leu
 435 440 445
 Ala Leu Trp Lys Cys Val Leu Thr Leu Leu Gln Ser Glu Glu Gln Ala
 450 455 460
 Val Arg Asp Ala Ala Thr Glu Thr Val Thr Thr Ala Met Ser Gln Glu
 465 470 475 480
 Asn Thr Cys Gln Ser Thr Glu Phe Ala Phe Cys Gln Val Asp Ala Ser
 485 490 495
 Ile Ala Leu Ala Leu Ala Leu Ala Val Leu Cys Asp Leu Leu Gln Gln
 500 505 510
 Trp Asp Gln Leu Ala Pro Gly Leu Pro Ile Leu Leu Gly Trp Leu Leu
 515 520 525
 Gly Glu Ser Asp Asp Leu Val Ala Cys Val Glu Ser Met His Gln Val
 530 535 540
 Glu Glu Asp Tyr Leu Phe Glu Lys Ala Glu Val Asn Phe Trp Ala Glu
 545 550 555 560
 Thr Leu Ile Phe Val Lys Tyr Leu Cys Lys His Leu Phe Cys Leu Leu
 565 570 575
 Ser Lys Ser Gly Trp Arg Pro Pro Ser Pro Glu Met Leu Cys His Leu
 580 585 590
 Gln Arg Met Val Ser Glu Gln Cys His Leu Leu Ser Gln Phe Phe Arg
 595 600 605
 Glu Leu Pro Pro Ala Ala Glu Phe Val Lys Thr Val Glu Phe Thr Arg
 610 615 620
 Leu Arg Ile Gln Glu Glu Arg Thr Leu Ala Cys Leu Arg Leu Leu Ala
 625 630 635 640
 Phe Leu Glu Gly Lys Glu Gly Glu Asp Thr Leu Val Leu Ser Val Trp
 645 650 655

177

Asp Ser Tyr Ala Glu Ser Arg Gln Leu Thr Leu Pro Arg Thr Glu Ala
 660 665 670

Ala Cys

<210> 249
 <211> 10
 <212> PRT
 <213> Homo sapiens

<400> 249
 Ile Ile Ser Gly Ser Glu Leu Ile Thr Gly
 1 5 10

<210> 250
 <211> 230
 <212> PRT
 <213> Homo sapiens

<400> 250
 Val Asp Gly Ile Asp Lys Leu Asp Ile Glu Phe Leu Gln Gln Phe Leu
 1 5 10 15

Glu Thr His Ser Arg Gly Pro Arg Leu His Ser Pro Gly His Ala Ser
 20 25 30

Gln Glu Ala Thr Pro Gly Ala Asn Met Ser Ser Gly Thr Glu Leu Leu
 35 40 45

Trp Pro Gly Ala Ala Leu Leu Val Leu Leu Gly Val Ala Ala Ser Leu
 50 55 60

Cys Val Arg Cys Ser Arg Pro Gly Ala Lys Arg Ser Glu Lys Ile Tyr
 65 70 75 80

Gln Gln Arg Ser Leu Arg Glu Asp Gln Gln Ser Phe Thr Gly Ser Arg
 85 90 95

Thr Tyr Ser Leu Val Gly Gln Ala Trp Pro Gly Pro Leu Ala Asp Met
 100 105 110

Ala Pro Thr Arg Lys Asp Lys Leu Leu Gln Phe Tyr Pro Ser Leu Glu
 115 120 125

Asp Pro Ala Ser Ser Arg Tyr Gln Asn Phe Ser Lys Gly Ser Arg His
 130 135 140

Gly Ser Glu Glu Ala Tyr Ile Asp Pro Ile Ala Met Glu Tyr Tyr Asn
 145 150 155 160

Trp Gly Arg Phe Ser Lys Pro Pro Glu Asp Asp Ala Asn Ser Tyr
 165 170 175

Glu Asn Val Leu Ile Cys Lys Gln Lys Thr Thr Glu Thr Gly Ala Gln
 180 185 190

178

Gln Glu Gly Ile Gly Gly Leu Cys Arg Gly Asp Leu Ser Leu Ser Leu
 195 200 205

Ala Leu Lys Thr Gly Pro Thr Ser Gly Leu Cys Pro Ser Ala Ser Pro
 210 215 220

Glu Glu Asp Glu Gly Ile
 225 230

<210> 251
 <211> 122
 <212> PRT
 <213> Homo sapiens

<400> 251
 Val Leu Trp Arg Glu Ala Ser Ala Leu Val Leu Ser Asn Arg Leu Ser
 1 5 10 15

Ser Gly Leu Leu His Asp Leu Leu Leu Gln Pro Ala Ile His Ser Arg
 20 25 30

Leu Phe Pro Arg Arg Ser Arg Gly Leu Ser Glu Gly Glu Gly Ser Ser
 35 40 45

Val Ser Leu Gln Arg Ser Arg Val Leu Ser Ala Met Lys His Val Leu
 50 55 60

Asn Leu Tyr Leu Leu Gly Val Val Leu Thr Leu Leu Ser Ile Phe Val
 65 70 75 80

Arg Val Met Glu Ser Leu Glu Gly Leu Leu Glu Ser Pro Ser Pro Gly
 85 90 95

Thr Ser Trp Thr Thr Arg Ser Gln Leu Ala Asn Thr Glu Pro Thr Lys
 100 105 110

Gly Leu Pro Asp His Pro Ser Arg Ser Met
 115 120

<210> 252
 <211> 129
 <212> PRT
 <213> Homo sapiens

<400> 252
 Tyr Thr Phe His Thr Gln Ile Phe Leu Asp Phe Pro Met Ile Phe Leu
 1 5 10 15

Thr Val Leu Pro Leu Ala Phe Leu Phe Leu His Ser Gly Phe Tyr His
 20 25 30

Tyr Ile Ser Phe Ser Cys Leu Phe Ser Leu Ser Leu Ala Leu Phe Phe
 35 40 45

Phe Leu Asp Val Ala Thr Phe Arg Arg Pro Gly Gln Leu Phe Cys Glu

179

50 55 60
 Arg Ser Val Leu Phe Asp Met Phe His Phe Gly Phe Val Ser Leu Phe
 65 70 75 80
 Leu His Glu Trp Ile Gln Ala Lys His Phe Trp Ala Gly Leu Phe Ile
 85 90 95
 Val Leu Pro Ser Asp Val Phe Phe Ser Val His His Leu Glu Ala Pro
 100 105 110
 Asp Gly Ser Phe Pro Asn Ile Ala Lys Leu Ser Leu Ile Ile Leu Leu
 115 120 125
 Arg

<210> 253
 <211> 99
 <212> PRT
 <213> Homo sapiens

<400> 253
 Gly Thr Arg Phe Pro Thr Gly Glu Thr Pro Ser Leu Gly Phe Thr Val
 1 5 10 15
 Thr Leu Val Leu Leu Asn Ser Leu Ala Phe Leu Leu Met Ala Val Ile
 20 25 30
 Tyr Thr Lys Leu Tyr Cys Asn Leu Glu Lys Glu Asp Leu Ser Glu Asn
 35 40 45
 Ser Gln Ser Ser Met Ile Lys His Val Ala Trp Leu Ile Phe Thr Asn
 50 55 60
 Cys Ile Phe Phe Cys Pro Val Ala Phe Phe Ser Phe Ala Pro Leu Ile
 65 70 75 80
 Thr Ala Ile Ser Ile Ser Pro Glu Ile Met Lys Ser Val Thr Leu Ile
 85 90 95
 Phe Phe Pro

<210> 254
 <211> 51
 <212> PRT
 <213> Homo sapiens

<400> 254
 Met Ile Lys His Val Ala Trp Leu Ile Phe Thr Asn Cys Ile Phe Phe
 1 5 10 15
 Cys Pro Val Ala Phe Phe Ser Phe Ala Pro Leu Ile Thr Ala Ile Ser
 20 25 30

180

Ile Ser Pro Glu Ile Met Lys Ser Val Thr Leu Ile Phe Phe Pro Cys
 35 40 45

Leu Leu Ala
 50

<210> 255

<211> 259

<212> PRT

<213> Homo sapiens

<400> 255

Gly Thr Arg Phe Pro Thr Gly Glu Thr Pro Ser Leu Gly Phe Thr Val
 1 5 10 15

Thr Leu Val Leu Leu Asn Ser Leu Ala Phe Leu Leu Met Ala Val Ile
 20 25 30

Tyr Thr Lys Leu Tyr Cys Asn Leu Glu Lys Glu Asp Leu Ser Glu Asn
 35 40 45

Ser Gln Ser Ser Met Ile Lys His Val Ala Trp Leu Ile Phe Thr Asn
 50 55 60

Cys Ile Phe Phe Cys Pro Val Ala Phe Phe Ser Phe Ala Pro Leu Ile
 65 70 75 80

Thr Ala Ile Ser Ile Ser Pro Glu Ile Met Lys Ser Val Thr Leu Ile
 85 90 95

Phe Phe Pro Leu Pro Ala Cys Leu Asn Pro Val Leu Tyr Val Phe Phe
 100 105 110

Asn Pro Lys Phe Lys Glu Asp Trp Lys Leu Leu Lys Arg Arg Val Thr
 115 120 125

Lys Lys Ser Gly Ser Val Ser Val Ser Ile Ser Ser Gln Gly Gly Cys
 130 135 140

Leu Glu Gln Asp Phe Tyr Tyr Asp Cys Gly Met Tyr Ser His Leu Gln
 145 150 155 160

Gly Asn Leu Thr Val Cys Asp Cys Cys Glu Ser Phe Leu Leu Thr Lys
 165 170 175

Pro Val Ser Cys Lys His Leu Ile Lys Ser His Ser Cys Pro Ala Leu
 180 185 190

Ala Val Ala Ser Cys Gln Arg Pro Glu Gly Tyr Trp Ser Asp Cys Gly
 195 200 205

Thr Gln Ser Ala His Ser Asp Tyr Ala Asp Glu Glu Asp Ser Phe Val
 210 215 220

Ser Asp Ser Ser Asp Gln Val Gln Ala Cys Gly Arg Ala Cys Phe Tyr
 225 230 235 240

181

Gln Ser Arg Gly Phe Pro Leu Val Arg Tyr Ala Tyr Asn Leu Pro Arg
 245 250 255

Val Lys Asp

<210> 256
 <211> 22
 <212> PRT
 <213> Homo sapiens

<400> 256
 Cys Asp Cys Cys Glu Ser Phe Leu Leu Thr Lys Pro Val Ser Cys Lys
 1 5 10 15

His Leu Ile Lys Ser His
 20

<210> 257
 <211> 81
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (20)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 257
 Ala Leu Glu Asn Ser Gly Ser Pro Gly Leu Gln Asp Ser Ala Arg Ala
 1 5 10 15

His Phe Asn Xaa Ser Leu Arg Ser Phe Ser Phe Leu Arg Asn Gln Met
 20 25 30

Tyr Ile Phe Glu Leu Ser Leu Tyr Leu Glu Gly Thr Ser Phe Val Val
 35 40 45

Val Leu Leu Phe Leu Leu Ile Ser Val Ser Leu Asp Ser Pro Pro Thr
 50 55 60

Thr Lys Gly Trp Asp Ser Val Leu His Ile Trp Val Pro Leu Ile Val
 65 70 75 80

Gln

<210> 258
 <211> 77
 <212> PRT
 <213> Homo sapiens

<400> 258
 Gly His Glu Ser Ile Cys Gly Ser Cys Arg Ser Trp Ile Tyr Phe Ser
 1 5 10 15

182

Ile Arg Cys Arg Arg Arg Met Arg Pro Trp Trp Ser Leu Leu Leu Glu
 20 25 30
 Ala Cys Ala Thr Cys Ala Gln Thr Gly Pro Thr Arg Ser Thr Ser Cys
 35 40 45
 Thr Gln Glu Val Ser His Ser Ser Ser Thr Ala Tyr Pro Ala Pro Met
 50 55 60
 Arg Arg Arg Cys Cys Leu Pro Ser Pro Arg Ser Cys Thr
 65 70 75

<210> 259
 <211> 119
 <212> PRT
 <213> Homo sapiens

<400> 259
 Lys Arg Ala Gly Val Glu Val Gly Gly Leu Val Met Ala Leu Ala Gly
 1 5 10 15
 Ser Val Phe Val Leu Gly Gly Val Leu Val Leu Cys Val Glu Arg Asn
 20 25 30
 Gly Glu Gly Glu Met Gly Trp Pro Gln His Leu Pro Lys Ser Gln Pro
 35 40 45
 Leu Ser Pro Pro Val Ala Val Arg Arg Cys Ser Phe Glu Arg Ser Trp
 50 55 60
 Ile Asp Leu Leu Val Glu Thr Ser Ser Ser Met Val Thr Cys Arg Gln
 65 70 75 80
 Gln Val Gly Thr Pro Asn Gly Met Glu Gly Arg Gly Gly Gly Pro Lys
 85 90 95
 Thr Thr Phe Pro Ile Arg Leu Gln Leu Ser Gly Ala Cys Ala Val Arg
 100 105 110
 Pro Glu Ile Gln Trp Glu Val
 115

<210> 260
 <211> 275
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (47)
 <223> Xaa equals any of the naturally occurring L-amino acids

 <220>
 <221> SITE
 <222> (94)

183

<223> Xaa equals any of the naturally occurring L-amino acids

<220> .

<221> SITE

<222> (192)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 260

Gln	Asp	Trp	Lys	Ala	Glu	Arg	Ser	Gln	Asp	Pro	Phe	Glu	Lys	Cys	Met
1				5					10					15	
Gln	Asp	Pro	Asp	Tyr	Glu	Gln	Leu	Leu	Lys	Val	Thr	Ile	Leu	Glu	Ala
			20					25					30		
Asp	Asn	Arg	Ile	Gly	Gly	Arg	Ile	Phe	Thr	Tyr	Arg	Asp	Gln	Xaa	Thr
		35					40					45			
Gly	Trp	Ile	Gly	Glu	Leu	Gly	Ala	Met	Arg	Met	Pro	Ser	Ser	His	Arg
	50					55					60				
Ile	Leu	His	Lys	Leu	Cys	Gln	Gly	Leu	Gly	Leu	Asn	Leu	Thr	Lys	Phe
	65				70					75					80
Thr	Gln	Tyr	Asp	Lys	Asn	Thr	Trp	Thr	Glu	Val	His	Glu	Xaa	Lys	Leu
				85					90					95	
Arg	Asn	Tyr	Val	Val	Glu	Lys	Val	Pro	Glu	Lys	Leu	Gly	Tyr	Ala	Leu
			100					105					110		
Arg	Pro	Gln	Glu	Lys	Gly	His	Ser	Pro	Glu	Asp	Ile	Tyr	Gln	Met	Ala
		115					120					125			
Leu	Asn	Gln	Ala	Leu	Lys	Asp	Leu	Lys	Ala	Leu	Gly	Cys	Arg	Lys	Ala
	130					135					140				
Met	Lys	Lys	Phe	Glu	Arg	His	Thr	Leu	Leu	Glu	Tyr	Leu	Leu	Gly	Glu
145					150					155				160	
Gly	Asn	Leu	Ser	Arg	Pro	Ala	Val	Gln	Leu	Leu	Gly	Asp	Val	Met	Ser
			165						170					175	
Glu	Asp	Gly	Phe	Phe	Tyr	Leu	Ser	Phe	Ala	Glu	Ala	Leu	Arg	Ala	Xaa
			180					185					190		
Ser	Cys	Leu	Ser	Asp	Arg	Leu	Gln	Tyr	Ser	Arg	Ile	Val	Gly	Gly	Trp
	195						200						205		
Asp	Leu	Leu	Pro	Arg	Ala	Leu	Leu	Ser	Ser	Leu	Ser	Gly	Leu	Val	Leu
	210					215						220			
Leu	Asn	Ala	Pro	Val	Val	Ala	Met	Thr	Gln	Gly	Pro	His	Asp	Val	His
225					230					235				240	
Val	Gln	Ile	Glu	Thr	Ser	Pro	Pro	Ala	Arg	Asn	Leu	Lys	Val	Leu	Lys
				245					250					255	
Ala	Asp	Val	Val	Leu	Leu	Thr	Ala	Ser	Gly	Pro	Ala	Val	Lys	Arg	Ile
			260					265					270		

184

Thr Phe Ser
275

<210> 261
<211> 212
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (123)
<223> Xaa equals any of the naturally occurring L-amino acids

<400> 261
Leu Pro Arg His Met Gln Glu Ala Leu Arg Arg Leu His Tyr Val Pro
1 5 10 15
Ala Thr Lys Val Phe Leu Ser Phe Arg Arg Pro Phe Trp Arg Glu Glu
20 25 30
His Ile Glu Gly Gly His Ser Asn Thr Asp Arg Pro Ser Arg Met Ile
35 40 45
Phe Tyr Pro Pro Pro Arg Glu Gly Ala Leu Leu Leu Ala Ser Tyr Thr
50 55 60
Trp Ser Asp Ala Ala Ala Ala Phe Ala Gly Leu Ser Arg Glu Glu Ala
65 70 75 80
Leu Arg Leu Ala Leu Asp Asp Val Ala Ala Leu His Gly Pro Val Val
85 90 95
Arg Gln Leu Trp Asp Gly Thr Gly Val Val Lys Arg Trp Ala Glu Asp
100 105 110
Gln His Ser Gln Gly Gly Phe Val Val Gln Xaa Pro Ala Leu Trp Gln
115 120 125
Thr Glu Lys Asp Asp Trp Thr Val Pro Tyr Gly Arg Ile Tyr Phe Ala
130 135 140
Gly Glu His Thr Ala Tyr Pro His Gly Trp Val Glu Thr Ala Val Lys
145 150 155 160
Ser Ala Leu Arg Ala Ala Ile Lys Ile Asn Ser Arg Lys Gly Pro Ala
165 170 175
Ser Asp Thr Ala Ser Pro Glu Gly His Ala Ser Asp Met Glu Gly Gln
180 185 190
Gly His Val His Gly Val Ala Ser Ser Pro Ser His Asp Leu Ala Lys
195 200 205
Glu Glu Gly Ser
210

185

<210> 262
 <211> 319
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (68)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (115)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (213)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 262
 Met Ala Pro Leu Ala Leu His Leu Leu Val Leu Val Pro Ile Leu Leu
 1 5 10 15
 Ser Leu Val Ala Ser Gln Asp Trp Lys Ala Glu Arg Ser Gln Asp Pro
 20 25 30
 Phe Glu Lys Cys Met Gln Asp Pro Asp Tyr Glu Gln Leu Leu Lys Val
 35 40 45
 Thr Ile Leu Glu Ala Asp Asn Arg Ile Gly Gly Arg Ile Phe Thr Tyr
 50 55 60
 Arg Asp Gln Xaa Thr Gly Trp Ile Gly Glu Leu Gly Ala Met Arg Met
 65 70 75 80
 Pro Ser Ser His Arg Ile Leu His Lys Leu Cys Gln Gly Leu Gly Leu
 85 90 95
 Asn Leu Thr Lys Phe Thr Gln Tyr Asp Lys Asn Thr Trp Thr Glu Val
 100 105 110
 His Glu Xaa Lys Leu Arg Asn Tyr Val Val Glu Lys Val Pro Glu Lys
 115 120 125
 Leu Gly Tyr Ala Leu Arg Pro Gln Glu Lys Gly His Ser Pro Glu Asp
 130 135 140
 Ile Tyr Gln Met Ala Leu Asn Gln Ala Leu Lys Asp Leu Lys Ala Leu
 145 150 155 160
 Gly Cys Arg Lys Ala Met Lys Lys Phe Glu Arg His Thr Leu Leu Glu
 165 170 175
 Tyr Leu Leu Gly Glu Gly Asn Leu Ser Arg Pro Ala Val Gln Leu Leu
 180 185 190

186

Gly Asp Val Met Ser Glu Asp Gly Phe Phe Tyr Leu Ser Phe Ala Glu
195 200 205

Ala Leu Arg Ala Xaa Ser Cys Leu Ser Asp Arg Leu Gln Tyr Ser Arg
210 215 220

Ile Val Gly Gly Trp Asp Leu Leu Pro Arg Ala Leu Leu Ser Ser Leu
225 230 235 240

Ser Gly Leu Val Leu Leu Asn Ala Pro Val Val Ala Met Thr Gln Gly
245 250 255

Pro His Asp Val His Val Gln Ile Glu Thr Ser Pro Pro Ala Arg Asn
260 265 270

Leu Lys Val Leu Lys Ala Asp Val Val Leu Leu Thr Ala Ser Gly Pro
275 280 285

Ala Val Lys Arg Ile Thr Phe Ser Pro Arg Cys Pro Ala Thr Cys Arg
290 295 300

Arg Arg Cys Gly Gly Cys Thr Thr Cys Arg Pro Pro Arg Cys Ser
305 310 315

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/15849**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C12N 15/11, 15/00, 15/63; C07H 21/02, 21/04

US CL : 536/23.1, 23.4; 435/320.1, 69.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.4; 435/320.1, 69.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Sequence searched SEQ ID NO:11, SEQ ID NO: 103, APS, STN, CAPLUS, terms nucleic acid, express, vector, pancreas islet cell tumors.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	US 5,849,498 A (BANDMAN et al) 15 December 1998, sequence listing.	1-10, 14-15, and 21
A	US 5,670,367 A (DORNER et al) 23 September 1997, especially sequence listing.	1-10, 14-15, and 21

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

B earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

30 SEPTEMBER 1999

Date of mailing of the international search report

21 OCT 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

LI LEE

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/15849

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-10, 14-15, and 21

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/15849

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-10, 14-15, and 21, drawn to isolated nucleic acid and expression system.

Group II, claim(s) 11-12 and 16, drawn to isolated polypeptide.

Group III, claim(s) 13, drawn to antibody.

Group IV, claim(s) 17, drawn to method for preventing a medical condition.

Group V, claim(s) 18-19, drawn to method of diagnosing a disease.

Group VI, claim(s) 20, 22, drawn to method for identifying a binding partner.

Group VII, claim 23, drawn to product produced by method of claim 20.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

The inventions listed as Groups I-VII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The special technical feature of Group I is considered to be isolated nucleic acid and expression system.

The special technical feature of Group II is considered to be isolated polypeptide.

The special technical feature of Group III is considered to be antibody.

The special technical feature of Group IV is considered to be a method for preventing a medical condition..

The special technical feature of Group V is considered to be method of diagnosing a disease.

The special technical feature of Group VI is considered to be a method for identifying a binding partner..

The special technical feature of Group VII is considered to be product produced by method of claim 20.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

There are 71 genes, from gene Nos 1-71.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: The 71 genes have different nucleic acid sequences and they are from different cell. Therefore they lack same special technical features.